ECOLOGICAL SOCIETY OF AMERICA

Ecology/Ecological Monographs/Ecological Applications

PREPRINT

This preprint is a PDF of a manuscript that has been accepted for publication in an ESA journal. It is the final version that was uploaded and approved by the author(s). While the paper has been through the usual rigorous peer review process of ESA journals, it has not been copy-edited, nor have the graphics and tables been modified for final publication. Also note that the paper may refer to online Appendices and/or Supplements that are not yet available. We have posted this preliminary version of the manuscript online in the interest of making the scientific findings available for distribution and citation as quickly as possible following acceptance. However, readers should be aware that the final, published version will look different from this version and may also have some differences in content.

The doi for this manuscript and the correct format for citing the paper are given at the top of the online (html) abstract.

Once the final published version of this paper is posted online, it will replace the preliminary version at the specified doi.

Neglected role of fungal community composition in explaining variation in wood decay rates

A. van der Wal^{1*}, E. Ottosson¹, W. de Boer^{1,2}

¹ Department of Microbial Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, 6708 PB, the Netherlands

² Department of Soil Quality, Wageningen University, Wageningen, 6708 PB, the Netherlands

Email addresses:

a.vanderwal@nioo.knaw.nl (* corresponding author)

e.ottosson@nioo.knaw.nl

w.deboer@nioo.knaw.nl

Running title: Wood decay rate varies with fungal composition

Abstract

Decomposition of wood is an important component of global carbon cycling. Most wood decomposition models are based on tree characteristics and environmental conditions; however, they do not include community dynamics of fungi which are the major wood decomposers. We examined the factors explaining variation in sapwood decay in oak tree stumps 2 and 5 years after cutting. Wood moisture content was significantly correlated with sapwood decay in younger stumps, whereas ITS-based composition and species richness of the fungal community were the best predictors for mass loss in the older stumps. Co-occurrence analysis showed that in freshly cut trees and in younger stumps fungal communities were non-randomly structured, whereas fungal communities in old stumps could not be separated from a randomly assembled community. These results indicate that the most important factors explaining variation in wood decay rates can change over time and that the strength of competitive interactions between fungi in decaying tree stumps may level off with increased wood decay. Our field analysis further suggests that ascomycetes may have a prominent role in wood decay, but their wood-degrading abilities need to be further tested under controlled conditions. The next challenging step will be to integrate fungal community assembly processes in wood decay models to improve carbon sequestration estimates of forests.

Key words: 454 pyrosequencing of ITS, assembly, fungal interactions, local scale, moisture content, saprotrophic fungi, sapwood, wood decomposition, *Quercus robur*.

39 **Introduction**

40 Dead wood is an important component in the functioning of forest ecosystems. It plays a major
41 role in nutrient cycling as a temporary storage stock of carbon and macronutrients which only
42 become available again during decomposition (Cornelissen et al. 2012). Therefore, a better
43 understanding of factors influencing the rate of wood decomposition can aid in estimating the
44 carbon sequestration capacity of forests under climate change.

45 To date, most wood decay models are based on wood properties (physical and chemical
46 characteristics of the tree species), moisture and temperature (Yin et al. 1999, Radtke et al. 2009,
47 Zell et al. 2009) and are used to predict wood decay rates over large temporal and spatial scales.
48 These models do not, however, account for the variation that is found at smaller temporal and
49 spatial scales (Palviainen et al. 2010, Woodall 2010). Most carbon is lost during the first decade
50 of wood decomposition, the period for which predictions by current models have the lowest
51 accuracy (Fahey et al. 2005). This hampers the extrapolation of short-term, site-based
52 measurements to larger temporal and spatial scales, thereby reducing the reliability of carbon
53 sequestration estimates of forests. The gap between observed and predicted decay rates could be
54 due to the fact that fungal community dynamics are not taken into account in current wood decay
55 models.

56 In terrestrial ecosystems higher fungi are the main decomposers of the major wood
57 polymers (cellulose, hemi-cellulose and lignin) (van der Wal et al. 2013). White rot fungi are the
58 only organisms known to be able to completely decompose lignin, whereas brown rot fungi only
59 modify lignin during decomposition of cellulose and hemi-cellulose. Soft rot occurs in wet wood,
60 making wood soft by hydrolysis of part of the cellulose, but with little or no effect on lignin.

Experiments have shown that the type of wood-rot and fungal identity can have a strong impact on wood decay rates (Boddy 2001).

When two or more fungal species are present in a woody resource, interactions between fungal species may occur that also affect the rate of decay. Freshly fallen wood may already contain established fungal species, or latently present fungal propagules which will be amongst the earliest colonizers (Boddy 2001, Parfitt et al. 2010). Furthermore, a number of wood-rot fungi have the ability to colonize living trees for instance by pathogenesis through the roots (Stokland et al. 2012) or vectoring by insects (Persson et al. 2011). These parasitic and / or endophytic fungi may continue to live as decomposers in the fallen tree and hence, have a head start in the competition for available resources in the wood. Other early colonizers include opportunistic fungi and bacteria that grow on easily accessible (hemi-)cellulose and simple soluble substrates (Van der Wal et al. 2007). An already established fungal species may inhibit but also promote the colonization of successor species (Heilmann-Clausen & Boddy 2005). The positive or negative effects on later establishing fungi may depend on alternation of the chemical environment through the production of antibiotics (composition, amount) as well as physical modification of the wood (Niemelä et al. 1995). Pre-emptive competition through the consumption of easily degradable substrates as well as the occupation of space inside the wood also results in a limited availability of substrates for secondary colonizing fungi (Boddy 2001, Payne et al. 2000). Hence, the identity and interactions of species that colonize first, may affect colonization success of later arriving species. This effect is often referred to as a priority effect (Fukami et al. 2010).

In the next phase of wood decay, when two or more wood-decaying higher fungi have been able to colonize wood, competitive interactions continue. This can also affect decay rates e.g. fungi can invest more resources in the production of secondary metabolites than in growth

and decomposition (Woodward & Boddy 2008). Thus, fungal species composition and interactions may have a strong impact on wood decay during all stages of decomposition.

The aim of this study was to assess the importance of fungal community composition to explain local variation in decay rates of naturally colonized woody resources. A few studies have indicated a possible relation between variation in wood decomposition rate and fungal community composition in naturally colonized logs. These studies used traditional methods to describe the fungal community such as isolation of mycelia by plating wood pieces on agar (Chapela et al. 1988, Boddy et al. 1989). Nowadays fungal communities can be described at much higher resolution using high-throughput DNA sequencing methods which are not biased by morphological or growth characteristics of the fungi. In this field study, we make use of a chronosequence of naturally decaying oak tree stumps in adjacent small-sized forest plots to minimize the variation in abiotic conditions.

Material and methods

Site description and field sampling

A chronosequence of decaying tree stumps of *Quercus robur* (English oak) was established in a forest stand on a sandy soil near Bergharen, The Netherlands (51°51'39 N, 5°40'15 E). The study stand consisted of *Q. robur* (about 70% of the vegetation) mixed with *Rubus fruticosus*, *Sorbus aucuparia*, *Betula pendula*, *Pteridium aquilinum* and *Amelanchier lamarckii*. At this location, three plots were chosen where oak trees had been cut in January 2007, November 2010 and March 2012, hereafter referred to as “old”, “young” and “fresh” samples, respectively. Plots were situated next to each other and plot sizes were about 1 ha. At each tree harvest, all trees were cut within a plot. In April 2012, 20 randomly selected stumps (stumps with diameters of <15 cm

were excluded), were sampled from the 2007 plot, 20 stumps from the 2010 plot, and 6 stumps were sampled from the 2012 plot to represent the starting point of decay. The average height of stumps in the 2007 and 2010 plots was 50 ± 10 cm, and the average height of stumps in the 2012 plot was 27 ± 7 cm. There was no significant relationship between stem height and sapwood or heartwood densities of stumps in the 2012 plots ($P > 0.4$), so we assumed that small differences in stem height of individual stumps were not affecting initial wood densities. The upper 5 cm of the stump was removed with a chain saw to avoid sampling mosses and fungal propagules present on the outer part, and a disc containing the next 3 cm was collected. Diameter of wood discs was 20 ± 2 cm in the 2007 plots, 21 ± 4 cm in the 2010 plots and 22 ± 3 cm in the 2012 plots. Discs were stored in plastic bags at -20°C until analyses.

Wood density and moisture content analyses

For each disc, a wedge-shaped piece (1/8 of the total disc) representing as much as possible all fungal decay patterns (e.g. interactions zones, type of wood decay) present in the whole disc was cut out and separated into sapwood, heartwood and, if still present, bark (Fig. 1). Volumes of each segment were calculated using Archimedes' volume displacement method. All samples were then oven dried at 70°C for three days and the density of each segment was calculated as dry weight per unit volume (g/cm^3). Moisture content (%) was calculated as $((\text{mass wet wood} - \text{mass dry wood}) / \text{dry wood}) * 100\%$.

Sample preparation

From each disc, sawdust samples were taken using an electric drill (bit diameter 8 mm). Sawdust from sapwood and heartwood were separately collected and the drill bit was sterilized between samples with ethanol. At least 15 drilled holes were made in both heartwood and sapwood. The

resulting sawdust samples were pooled resulting in two samples per disc: one from heartwood and one from sapwood. Samples were stored at -20°C until further analyses (Fig. 1).

DNA extraction, amplification and sequencing

Sapwood sawdust samples were frozen in liquid nitrogen and ground into a fine powder. Heartwood samples were excluded for further analyses (see Results section below). DNA was isolated from 0.15 g fresh weight of sapwood samples using the PowerSoil DNA Isolation kit according to the manufacturer's instructions (MO BIO Laboratories, Inc.), with some modifications: after adding solution C1 (causing cell lysis), samples were incubated at 60°C for 30 min., and after adding solution C6 (releasing DNA from spin filter), samples were incubated at 30°C for 10 min. The nuclear rDNA internal transcribed spacer (ITS) region was amplified using the fungal-specific primer pair fITS9 and ITS4 (Ihrmark et al. 2012). Adapter sequences were added to the primers as recommended by Roche as well as 6-bp tags specific for each sample. PCR reactions were performed in 25 µl reaction mixtures and contained 400 µM of each dNTP, 1 U of FastStart Expand High Fidelity polymerase (Roche Applied Sciences, Indianapolis, IN), 2.5 µl 10 x PCR buffer with MgCl₂, 10 µM of each of the two primers and 1 µl DNA (1-10 ng). The temperature cycling PCR conditions were: denaturation at 95°C for 5 min, followed by 30 cycles of 95°C for 30 s, 58°C for 30 s and 72°C for 1 min. The final extension step was 72°C for 10 min. After confirming the presence of expected sizes of PCR products by agarose- gel electrophoresis, PCR products from 4 reactions were pooled per sample and purified using a QIAquick PCR Purification Kit (Qiagen). DNA in samples was quantified by a fluorescence-based method (Pico Green assay) and the samples were sequenced (Macrogen Inc. Company, South Korea) on a Roche 454 automated sequencer and GS FLX system using titanium chemistry (454 Life Sciences, Branford, CT).

Bioinformatics

Sequences and quality information were extracted from the Standard Flowgram Format (SFF) files using the SFF converter tool in the Galaxy interface (Goecks et al. 2010). The 454 SFF files are deposited in the European Nucleotide Archive (<http://www.ebi.ac.uk/ena/data/view/PRJEB4497>). Sequences were analyzed using the Qiime version 1.2.1 scripts (Caporaso et al. 2010), which were made available in the Galaxy interface. Quality filtering of the sequences involved the removal of short sequences (< 200 bp), sequences with low read quality, and sequences containing homopolymers or ambiguous characters exceeding six nucleotides. The sequences were also checked for PCR chimeras using UCHIME version 4.2.40 (Edgar et al. 2011). The sequences passing the quality control thresholds were clustered into operational taxonomic units (OTUs) using USEARCH version 5.2.236 (Edgar, 2010) with a minimum sequence identity cutoff of 97%. Sequences within clusters of dominant OTUs (accounting for $\geq 10\%$ of all the sequences in each sample) were grouped based on % identity scores in ClustalX v.2.1 and manually checked and blasted in the UNITE database (Abarenkov et al., 2010) to confirm that sequences in each OTU resulted in the same taxonomic identity. The average length of the ITS sequences passing the filtering step was 380 bp. For each OTU, the most abundant sequence was selected as a representative for all sequences within an OTU. Taxonomy was assigned to representative sequences by comparing them with known reference sequences in the UNITE and GenBank (NCBI) database using the Blastn algorithm. Sequences were, whenever possible, identified to the species (> 98% similarity) or genus (94-97% similarity) level. The relative abundance of each OTU was calculated by dividing the number of sequences per OTU by the total number of sequences per sample.

Enzyme assays

Enzyme activities (laccase, manganese peroxidase, cellulase and hemicellulase) were assayed spectrophotometrically in the same extracts according to Van der Wal et al. (2007). Briefly, 8 ml of milliQ water was added to 1 g of sawdust and shaken for 1 h at room temperature, and then the slurry was pressed over a stainless steel filter (containing pores with a diameter of 2 mm). The supernatants were kept at -20°C until analysis of enzyme activities. Laccase activity was measured via oxidation of ABTS (2,2'-azinobis(3-ethylbenzthiazoline-6-sulfonic acid)), and manganese peroxidase activity was measured via the oxidative coupling of DMAB (3-dimethylaminobenzoic acid) and MBTH (3-methyl-2-benzothiazolinone hydrazine hydrochloride) in the presence of Mn^{2+} and H_2O_2 . The activity of endo-1,4-β-glucanase as an indicator of cellulase was estimated using carboxymethyl cellulose linked with Remazol brilliant blue R (Azo-CMCellulose, Megazyme, Bray, Ireland), and the activity of endo-1,4-β-xylanase (xylanase) as indicator of hemicellulase was estimated by using birchwood xylan linked with Remazol brilliant blue (Azo-xylan, Megazyme, Bray, Ireland) as a substrate.

Data analysis and statistics

The relation between the percentage of mass loss, wood moisture content, diameter, OTU richness, fungal diversity and fungal evenness was calculated by linear regression using SPSS (version 20.0.0). Statistical significance was assumed at $P < 0.05$.

Ordination analyses were performed in Canoco version 4.5. To identify if samples with the same “stump age” grouped together based on the composition of their fungal communities, a Detrended Correspondence Analysis (DCA) was performed. Canonical Correspondence Analysis (CCA) was used to test if the composition of the fungal community is related to the percentage of

mass loss, tree diameter or wood moisture content, and to extracellular enzyme activities.

Significance of canonical axes was assessed by the forward approach using Monte Carlo

permutation tests under the reduced model. CCA was performed separately on the samples

collected from the 2007 and 2010 plots.

Fungal diversity was calculated as the Shannon index $H' = -\sum(p_i \times \ln p_i)$, where p_i represents the relative abundance of species (OTU) i , and the Shannon's equitability (evenness) $E_H = H' / \ln S$, where S represents the total number of species (OTUs) present in the community.

Co-occurrence analysis of fungal communities were performed by a C-score analysis as well as a checkerboard index using default settings in EcoSim version 7.0 (Gotelli & Entsminger 2009). In a competitively structured community, both the observed C-score and checkerboard index are significantly larger than values derived from a community assembled in a random pattern. Samples with the same "stump age" were grouped and each group was analyzed separately. Species that occur in less than three samples (stumps) were removed from each matrix since they are unlikely to provide biologically meaningful information about co-occurrence.

Results

Wood densities

Heartwood density of old samples of oak stumps was almost the same as that of fresh samples (Fig. 2, Welch's test, $P = 0.16$). In contrast, the sapwood density of fresh samples was significantly higher than in young and old samples (Fig. 2, Welch's test, $P < 0.001$). Remarkably, sapwood densities between young and old samples did not significantly differ ($P = 0.57$). Since heartwood was apparently hardly degraded in 5 years, we focused our measurements and data analyses only on sapwood samples.

225

226 Relation between sapwood decay, diameter and moisture content

227 For both young and old samples, there was no significant relationship between sapwood decay
 228 and diameter of the tree stump (Appendix A, Fig. A1A & A1C). In young samples, we found a
 229 significant positive relation between sapwood moisture content and wood decay (Appendix A,
 230 Fig. A1B, $P = 0.02$) but we did not find this relation for old stumps (Appendix A, Fig. A1D, $P =$
 231 0.45).

232

233 Fungal community composition and stump age

234 In total 425,766 quality sequences were obtained from 42 sapwood samples, and the mean
 235 number of sequences per sample was 10,198. We identified in total 447 different OTUs
 236 (Supplement 1) of which 262 were ascomycetes, 148 basidiomycetes, 7 zygomycetes, 3
 237 glomeromycetes, 1 chytridiomycete and 26 could not be identified. In fresh samples 231 OTUs
 238 were identified and the most abundant OTUs consisted of only 10 OTUs of which most had
 239 sequence similarities with early successional fungal species such as sugar fungi (yeasts),
 240 endophytes and plant parasites. Only one OTU had sequence similarities with a white rot fungus
 241 (Appendix A, Fig. A2; Appendix B & C). In young samples, 270 OTUs were identified of which
 242 26 OTUs comprised the most abundant ones. Eleven of these OTUs had the highest match with
 243 saprotrophic fungi, of which ten were assigned to white rot fungi, two OTUs were assigned to
 244 parasites, and two OTUs were assigned to fungi with both saprotrophic and parasitic abilities
 245 (Fig. 3, Appendix B & D). In old samples, 243 OTUs were identified and the most abundant
 246 OTUs consisted of 18 OTUs. Eight of these OTUs had the highest match with saprotrophic fungi,

of which seven were assigned to white rot fungi. One OTU was assigned to a fungus with both saprotrophic and parasitic abilities (Fig. 4, Appendix B & E). In fresh, young and old sapwood samples 30%, 23% and 44% of the abundant OTUs could not be identified to the genus level. Most of the dominant OTUs were found in only one or two samples, showing that the abundant OTUs were not equally distributed among stumps. However, two OTUs were dominantly present in all stumps (fresh, young and old), i.e. *Mollisia* sp. and an unidentified *Helotiales* (Appendix B).

The fungal community composition of fresh samples grouped together on the left side in the DCA plot (Fig. 5). Young samples showed overlap in species composition both with fresh and old samples; old samples tended to group together on the right side of the DCA biplot. Therefore, the first DCA axis may represent a successional gradient of fungal communities. On the vertical axis young and old samples are much more spread out than fresh samples, and this can be interpreted as an increase in the variation in fungal community composition among young and old stumps.

Both the C-score and checkerboard pair analysis from the fresh and young samples showed that the fungal communities were non-randomly structured since both values were significantly higher compared to the mean values derived from the randomized communities. In old samples the observed community matrix could not be separated from the values derived from the randomized community (Appendix F).

Effect of fungal community composition on sapwood decay rates

In old samples the number of OTUs (OTU richness) per sample was positively related with sapwood decay (Appendix A, Fig. A3, $P = 0.04$), which was not explained by a significant relationship between number of reads per sample and OTU richness ($P > 0.05$). For both young and old samples, there was no relation between fungal diversity and / or fungal evenness with sapwood decay ($P > 0.05$).

Forward selection in CCA showed that fungal community composition was significantly correlated with sapwood decay in old samples ($P = 0.02$), explaining 10% of the total variance in species composition. Fig. 4 reveals that the most abundant OTUs in the samples that were hardly decomposed, consisted of *Phlebia radiata*, and in the samples that had strongly decomposed an two unidentified *Sordariales* and one unidentified *Heliotales* both belonging to the *Ascomycota* were most abundant. One OTU identified as *Trametes versicolor* was almost exclusively present in samples with similar mass loss (47 and 54%). In some samples showing different mass losses the same OTU, identified as *Mollisia sp.*, was among the dominant ones, but then always in combination with another fungal species. In young samples forward selection in CCA did not reveal a relationship between fungal community composition and wood decay ($P > 0.89$). In accordance, Fig. 3 shows that *Panellus stipticus* was abundant throughout several samples showing different mass losses, all in combination with an OTU identified as *Trametes versicolor*. Hence, in young samples, a similar composition of abundant fungal species in different stumps did not result in similar sapwood decay rates.

Other variables (moisture content and tree diameter) were not significantly related to fungal community composition (CCA, $P > 0.17$). Enzyme activities were also not significantly related to fungal community composition or wood decay (CCA, $P > 0.2$). Laccase activity was in general low in most samples (ranging from 0 to $0.43 \text{ nmol g}^{-1} \text{ h}^{-1}$). Remarkably, the highest laccase activity was found in the least decayed old sample ($5.61 \text{ nmol g}^{-1} \text{ h}^{-1}$).

292

293 Discussion

294 The variation in initial sapwood decay was significantly correlated with wood moisture content,
 295 whereas variation in later stages of sapwood decay was significantly related to the composition
 296 and OTU richness of the fungal community. This study is the first to indicate that the most
 297 important factors explaining variation in wood decay rates may change over time. Earlier studies
 298 focused only on the effect of abiotic factors and differences in traits of tree species (Mackensen et
 299 al. 2003, Brischke & Rapp 2008, Cornwell et al. 2009) In addition, other studies did not include a
 300 time series (Chapela et al. 1988, Boddy et al. 1989, Lindner et al. 2011), concentrated on
 301 decomposition stages instead of time of decomposition (Rajala et al. 2012) or used only fruiting
 302 body data and culture methods for detection of fungi (Chapela et al. 1988, Boddy et al. 1989).
 303 Despite the fact that early-successional fungal communities differed across individual young
 304 stumps, there was no indication that differences in community composition had an impact on
 305 initial sapwood decay. Instead, wood moisture content could partly (27%) explain variation in
 306 wood decay. However, the importance of moisture as explaining factor for longer term wood
 307 decay is unclear as there was no significant relationship with decay in older stumps.

308 As wood decomposition proceeded, we observed a significant relationship between the
 309 composition of the fungal community and mass loss. Potential mechanisms explaining this
 310 relation could be based on specific traits of fungal species, e.g. the presence or absence of a
 311 strong decay species (Lindner et al. 2011) or interactions between fungal species during
 312 community assembly processes (Fukami et al. 2010, Van der Wal et al. 2013). For instance,
 313 *Phlebia radiata* was the dominant fungus in samples that were hardly decomposed (Fig. 4). This
 314 fungus is known for its replacing ability at later stages in community development by lysing other

wood-decaying fungi (Rayner & Todd 1979). This may indicate that fungi that colonized first caused hardly any mass loss (i.e. consuming only the easily available carbon) and during replacement by *P. radiata*, fungal hyphae instead of wood are consumed. Alternatively, during competition fungi may invest more in defensive metabolites such as melanin than in growth and decomposition. This idea is supported by the high laccase activity found in the least decayed sample as laccases are thought to be involved in the formation of melanin (Baldrian 2006).

Most of the abundant OTUs were only found in one or two samples, showing that dominant fungal species were unique on individual stumps. This reflects the stochastic nature of fungal colonization (see discussion below). However, two OTUs, both assigned to the order *Helotiales*, were dominantly present in fresh, young and old stumps. Members of the *Helotiales* include saprotrophic-, parasitic- and endophytic fungi (Wang et al. 2006, Tedersoo et al. 2009). Since these fungi were already present in freshly cut stumps, it may indicate that they first displayed the endophytic lifestyle. This lifestyle will put these fungi in a good position to shift to saprophytism after the tree died, given them a competitive advantage over other later-arriving fungi for available resources in the wood (the so-called priority effect). In addition, their presence may influence the success of later establishing fungi by altering the chemical environment through the production of antibiotics or by physically modifying the wood (Niemelä et al. 1995, Heilmann-Claussen & Boddy 2005). The effect of relative abundance of the *Helotiales* species on wood decay did not show a clear pattern. The OTU assigned as *Mollisia sp.* was present in quickly as well as in slowly decaying stumps, whereas the OTU assigned as *unidentified Helotiales (4)* was present in stumps that lost already at least 40% of their mass in both young and old stumps. Therefore, we cannot draw any conclusions about the effect of *Helotiales* fungi on wood decay rate, but because of their abundance throughout all decay stages it is worthwhile to include them in controlled wood decay experiments.

A positive relation between OTU richness and mass loss was observed in old stumps (Fig. S3). We only found this relation based on richness, indicating that the presence of a certain species is more important for this relation than its abundance. Also Rajala et al. (2011) observed an increase of rRNA-based fungal richness with wood decay in naturally decaying spruce logs, but also the opposite has been found in wood that was artificially inoculated with different fungal species (Fukami et al. 2010). A positive relation between species richness and wood decay may point to additive or synergistic activities of fungal species (Fukasawa et al. 2011, Hättenschwiler et al. 2011). In wood, however, competitive interactions between wood-rot fungi are very common, can take place at all stages of wood decomposition and have a strong impact on fungal community composition (Boddy 2001). However, the fact that the fungal communities in old stumps could not be separated from a randomly assembled community in the co-occurrence analysis may indicate that competitive interactions are not the most dominant type of interaction in the old stumps (Appendix F). Indeed, in fresh and young samples, both the C-score and the checkerboard index showed that fungal communities were non-randomly structured, which could point to competitive species interactions (Gotelli & Entsminger 2009). Therefore, the strength of competitive interactions between fungi in decaying tree stumps may level off with increased wood decay. Alternatively, advanced decayed wood may cause an increase in the heterogeneity of wood polymer structures, creating a greater range of fungal microhabitats.

In wood that most quickly decomposed, 95% of the fungal community consisted of taxa belonging to the Ascomycota (Appendix G). It is generally assumed that basidiomycete fungi have greater enzyme diversity than ascomycetes and are thus causing more weight loss (Osono et al. 2003). Our results may imply that also ascomycete fungi can generate fast decomposition rates. Only one other study addressed the possible importance of ascomycete fungi in natural wood decay (Boddy et al. 1989). Another possibility is that these ascomycete fungi have replaced

the basidiomycetes that first decayed the wood rapidly. Wood decay abilities of ascomycetes need to be further investigated under controlled conditions.

Usually the differentiation between sapwood and heartwood is not made to determine the decay rate. Here we observed a strong decomposition resistance of heartwood. This is probably due to the presence of organic toxic compounds in oak heartwood, which inhibits microbial growth (Schmidt 2006). In contrast, the sapwood was quickly decaying during the first year and thereafter the average sapwood decay leveled off. Lumping the decay rate of heartwood and sapwood may thus result in an underestimation of the decay rate of sapwood.

The relation between fungal community composition and wood mass loss was significant, but only explained 10% of the total variance. This is very likely due to the huge differences of fungal communities among individual tree stumps (Appendix B, Fig. 3 & 4). Large differences of fungal communities across dead wood units have been reported before on spruce logs (Kubartová et al. 2012). The large variation among tree stumps could be due to stochastic fungal colonization processes of fungi via for instance dead tree roots, infection by wind- or insect-dispersed spores or mycelial fragments (Persson et al. 2011) or, as indicated above, some fungi may be already present in living trees (Parfitt et al. 2010). A further increase in variation among fungal communities during later stages of decay can be expected due to subsequent community assembly processes (Van der Wal et al. 2013).

In summary, current wood decay models do not explain the variation in wood mass loss that is found on a local scale, consequently restricting the possibility to reliably estimate carbon sequestration rates at larger temporal and spatial scales. Our results show that moisture content may be a reliable predictor for wood mass loss during the first years of wood decay, but after this period, the fungal community composition is contributing to explain the variance in wood mass

loss. The identity of fungal species and their richness seem to affect decomposition. The effect of ecologically relevant combinations of fungi on wood decay rates should be assessed under controlled conditions to formulate rules that can be used to optimize current wood decay models.

Acknowledgements

We thank Hans Hengeveld, district manager of Geldersch Landschap & Kasteelen, for permission to sample the stumps, Henk Eshuis for helping with wood sawing and Mattias de Hollander for providing help with bioinformatics issues. Funding was provided by the Netherlands Organisation for Scientific Research (NWO) in the form of a personal Veni grant to A.v.d.W and E.O was supported by the foundation of Anna and Gunnar Vidfeldt. This is publication number 5626 of the NIOO-KNAW Netherlands Institute of Ecology.

Literature cited

- Abarenkov, K., R. Henrik Nilsson, K.-H. Larsson, I. J. Alexander, U. Eberhardt, S. Erland, K. Høiland, R. Kjølner, E. Larsson, T. Pennanen, R. Sen, A. F. S. Taylor, L. Tedersoo, B. M. Ursing, T. Vrålstad, K. Liimatainen, U. Peintner, and U. Kõljalg. 2010. The UNITE database for molecular identification of fungi – recent updates and future perspectives. *New Phytologist* **186**:281-285.
- Baldrian, P. 2006. Fungal laccases – occurrence and properties. *FEMS Microbiology Reviews* **30**:215-242.

- 406 Boddy, L. 2001. Fungal community ecology and wood decomposition processes in angiosperms:
407 from standing tree to complete decay of coarse woody debris. *Ecological Bulletins* **49**:43-
408 56.
- 409 Boddy, L., E. M. Owens, and I. H. Chapela. 1989. Small-scale variation in decay rate within logs
410 one year after felling- effect of fungal community structure and moisture content. *FEMS*
411 *Microbiology Ecology* **62**:173-184.
- 412 Brischke, C. and A. Rapp. 2008. Dose-response relationships between wood moisture content,
413 wood temperature and fungal decay determined for 23 European field test sites. *Wood*
414 *Science and Technology* **42**:507-518.
- 415 Caporaso, J. G., J. Kuczynski, J. Stombaugh, K. Bittinger, F. D. Bushman, E. K. Costello, N.
416 Fierer, A. G. Pena, J. K. Goodrich, J. I. Gordon, G. A. Huttley, S. T. Kelley, D. Knights,
417 J. E. Koenig, R. E. Ley, C. A. Lozupone, D. McDonald, B. D. Muegge, M. Pirrung, J.
418 Reeder, J. R. Sevinsky, P. J. Turnbaugh, W. A. Walters, J. Widmann, T. Yatsunenko, J.
419 Zaneveld, and R. Knight. 2010. QIIME allows analysis of high-throughput community
420 sequencing data. *Nat Meth* **7**:335-336.
- 421 Chapela, I. H., L. Boddy, and A. D. M. Rayner. 1988. Structure and development of fungal
422 communities in beech logs four and a half years after felling. *FEMS Microbiology Letters*
423 **53**:59-70.
- 424 Cornelissen, J. H. C., U. Sass-Klaassen, L. Poorter, K. van Geffen, R. S. P. van Logtestijn, J. van
425 Hal, L. Goudzwaard, F. J. Sterck, R. Klaassen, G. T. Freschet, A. van der Wal, H. Eshuis,
426 J. Zuo, W. de Boer, T. Lamers, M. Weemstra, V. Cretin, R. Martin, J. den Ouden, M. P.
427 Berg, R. Aerts, G. M. J. Mohren, and M. M. Hefting. 2012. Controls on coarse wood
428 decay in temperate tree species: birth of the LOGLIFE experiment. *Ambio* **41**:231-245.

- 429 Cornwell, W. K., J. H. C. Cornelissen, S. D. Allison, J. Bauhus, P. Eggleton, C. M. Preston, F.
 430 Scarff, J. T. Weedon, C. Wirth, and A. E. Zanne. 2009. Plant traits and wood fates across
 431 the globe: rotted, burned, or consumed? *Global Change Biology* **15**:2431-2449.
- 432 Edgar, R. C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
 433 **26**: 2460-2461.
- 434 Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., and Knight R. 2011. UCHIME improves
 435 sensitivity and speed of chimera detection. *Bioinformatics* **27**: 1–7.
- 436 Fahey, T. J., T. G. Siccama, C. T. Driscoll, G. E. Likens, J. Campbell, C. E. Johnson, J. J. Battles,
 437 J. D. Aber, J. J. Cole, M. C. Fisk, P. M. Groffman, S. P. Hamburg, R. T. Holmes, P. A.
 438 Schwarz, and R. D. Yanai. 2005. The biogeochemistry of carbon at Hubbard Brook.
 439 *Biogeochemistry* **75**:109-176.
- 440 Fukami, T., I. A. Dickie, J. P. Wilkie, B. C. Paulus, D. Park, A. Roberts, P. K. Buchanan, and R.
 441 B. Allen. 2010. Assembly history dictates ecosystem functioning: evidence from wood
 442 decomposer communities. *Ecology Letters* **13**:675-684.
- 443 Fukasawa, Y., Osono, T., and Takeda H. 2011. Wood decomposing abilities of diverse
 444 lignicolous fungi on non-decayed and decayed beech wood. *Mycologia* **103**: 474–482.
- 445 Goecks, J., Nekrutenko, A., and Taylor, J. 2010. Galaxy: a comprehensive approach for
 446 supporting accessible, reproducible, and transparent computational research in the life
 447 sciences. *Genome Biology* **11**: R86.
- 448 Gotelli, N.J., and Entsminger, G.L. 2009. EcoSim: Null Models Software for Ecology. Version 7.
 449 Acquired Intelligence Inc. & Ke-sey-Bear, Jericho, VT. Available at:
 450 <http://garyentsminger.com/ecosim.htm>.
- 451 Hättenschwiler, S., N. Fromin, and S. Barantal. 2011. Functional diversity of terrestrial microbial
 452 decomposers and their substrates. *Comptes Rendus Biologies* **334**:393-402.

- 453 Heilmann-Clausen, J and Boddy, L. 2005. Inhibition and stimulation effects in communities of
454 wood decay fungi: Exudates from colonized wood influence growth by other species.
455 Microbial Ecology **49**:399–406.
- 456 Ihrmark, K., I. T. M. Bodeker, K. Cruz-Martinez, H. Friberg, A. Kubartova, J. Schenck, Y. Strid,
457 J. Stenlid, M. Brandström-Durling, K. E. Clemmensen, and B. D. Lindahl. 2012. New
458 primers to amplify the fungal ITS2 region – evaluation by 454-sequencing of artificial and
459 natural communities. FEMS Microbiology Ecology **82**:666-677.
- 460 Kubartová, A., E. Ottosson, A. Dahlberg, and J. Stenlid. 2012. Patterns of fungal communities
461 among and within decaying logs, revealed by 454 sequencing. Molecular Ecology
462 **21**:4514-4532.
- 463 Lindner, D. L., R. Vasaitis, A. Kubartová, J. Allmér, H. Johannesson, M. T. Banik, and J. Stenlid.
464 2011. Initial fungal colonizer affects mass loss and fungal community development in
465 *Picea abies* logs 6 yr after inoculation. Fungal Ecology **4**:449-460.
- 466 Mackensen, J., J. Bauhus, and E. Webber. 2003. Decomposition rates of coarse woody debris - A
467 review with particular emphasis on Australian tree species. Australian Journal of Botany
468 **51**:27-37.
- 469 Niemelä, T., P. Renvall, and R. Penttilä. 1995. Interactions of fungi at late stages of wood
470 decomposition. Annales Botanici Fennici **32**:141–152.
- 471 Osono, T, Fukasawa, Y., and Takeda, H. 2003. Roles of diverse fungi in larch needle-litter
472 decomposition. Mycologia **95**:820–826.
- 473 Palviainen, M., Finer, L., Laiho, R., Shorohova, E., Kapitsa, E., and Vanha-Majamaa, I. 2010.
474 Carbon and nitrogen release from decomposing Scots pine, Norway spruce and silver
475 birch stumps. Forest Ecology Management **259**:390–398.

- 476 Parfitt, D., J. Hunt, D. Dockrell, H. J. Rogers, and L. Boddy. 2010. Do all trees carry the seeds of
477 their own destruction? PCR reveals numerous wood decay fungi latently present in
478 sapwood of a wide range of angiosperm trees. *Fungal Ecology* **3**:338-346.
- 479 Payne, C., A. Bruce, and H.Y. Staines. 2000. Yeast and bacteria as biological control agents
480 against fungal discolouration of *Pinus sylvestris* blocks in laboratory-based tests and the
481 role of antifungal volatiles. *Holzforschung* **54**:563-569.
- 482 Persson, Y., Ihrmark, K., and Stenlid, J. 2011. Do bark beetles facilitate the establishment of rot
483 fungi in Norway spruce? *Fungal Ecology* **4**:262–269.
- 484 Radtke, P. J., R. L. Amateis, S. P. Prisley, C. A. Copenheaver, D. C. Chojnacky, J. R. Pittman,
485 and H. E. Burkhart. 2009. Modeling production and decay of coarse woody debris in
486 loblolly pine plantations. *Forest Ecology and Management* **257**:790-799.
- 487 Rajala, T., M. Peltoniemi, J. Hantula, R. Mäkipää, and T. Pennanen. 2011. RNA reveals a
488 succession of active fungi during the decay of Norway spruce logs. *Fungal Ecology*
489 **4**:437-448.
- 490 Rajala, T., Peltoniemi, M., Pennanen, T., and Mäkipää, R. 2012. Fungal community dynamics in
491 relation to substrate quality of decaying Norway spruce (*Picea abies* [L.] Karst.) logs in
492 boreal forests. *FEMS Microbiology Ecology* **81**:494–505.
- 493 Rayner, A.D.M., and Todd, N.K. 1979. Population and community structure and dynamics of
494 fungi in decaying wood. *Advances in Botanical Research* **7**:334-420.
- 495 Schmidt, O. 2006. Wood and tree fungi. Biology, damage, protection, and use. Springer-Verlag,
496 Heidelberg.
- 497 Stokland, J.N., Siitonen, J., and Jonsson, B.G. 2012. Biodiversity in dead wood. Cambridge
498 University Press, New York.

- 499 Tedersoo, L., K. Pärtel, T. Jairus, G. Gates, K. Põldmaa, and H. Tamm. 2009. Ascomycetes
500 associated with ectomycorrhizas: molecular diversity and ecology with particular
501 reference to the Helotiales. *Environmental Microbiology* **11**:3166-3178.
- 502 Van der Wal, A., W. de Boer, W. Smant, and J. van Veen. 2007. Initial decay of woody
503 fragments in soil is influenced by size, vertical position, nitrogen availability and soil
504 origin. *Plant and Soil* **301**:189-201.
- 505 Van der Wal, A., T. D. Geydan, T. W. Kuyper, and W. de Boer. 2013. A thready affair: linking
506 fungal diversity and community dynamics to terrestrial decomposition processes. *FEMS*
507 *Microbiology Reviews* **37**:477-494.
- 508 Wang, Z., M. Binder, C. L. Schoch, P. R. Johnston, J. W. Spatafora, and D. S. Hibbett. 2006.
509 Evolution of helotialean fungi (Leotiomycetes, Pezizomycotina): A nuclear rDNA
510 phylogeny. *Molecular Phylogenetics and Evolution* **41**:295-312.
- 511 Woodall, C.W. 2010. Carbon flux of down woody materials in forests of the North Central
512 United States. *International Journal of Forestry Research* **2010**:1-9.
- 513 Woodward, S. and L. Boddy. 2008. Chapter 7 Interactions between saprotrophic fungi. Pages
514 125-141 *in* J. C. F. Lynne Boddy and W. Pieter van, editors. British Mycological Society
515 Symposia Series. Academic Press.
- 516 Yin, X.W. 1999. The decay of forest woody debris: numerical modeling and implications based
517 on some 300 data cases from North America. *Oecologia* **121**:81-98.
- 518 Zell, J., Kandler, G., and M. Hanewinkel. 2009. Predicting constant decay rates of coarse woody
519 debris- A meta-analysis approach with a mixed model. *Ecological Modelling* **220**:904-
520 912.

522 **Description of Ecological Archives material**

523 **Appendices**

524 **Appendix A:** Three figures showing 1) relationships between sapwood decay, wood moisture
525 content and stump diameter, 2) percentage of most abundant OTUs in freshly cut trees and 3)
526 relationship between sapwood decay and OTU richness in old stumps.

527 **Figure A1:** Relation between sapwood decay in oak stumps (presented as % of wood mass loss),
528 wood moisture content and tree stump diameter. A & B: measured in young samples (1.4 years
529 since cutting), C & D: measured in old samples (5.3 years since cutting).

530 **Figure A2:** Percentage of most abundant operational taxonomic units (OTUs) in sapwood
531 samples collected from freshly cut oak trees. (A): Ascomycota, (B): Basidiomycota. Numbers
532 between brackets indicate OTU number (see Appendix B) of OTUs that could not be identified to
533 the species level.

534 **Figure A3:** Relation between sapwood decay (presented as % of wood mass loss) and the
535 number of OTUs (OTU richness) measured in old samples (5.3 years since cutting) taken from
536 oak stumps.

537 **Appendix B:** Most abundant OTUs (representing $\geq 10\%$ of the total number of sequences) in
538 decaying oak sapwood. Ecological groups are defined as: p = parasite, wr = white-rot fungus, sr =
539 soft-rot fungus, sa = fungus with saprotrophic abilities, sp = combining both saprotrophic and
540 parasitic abilities, en = endophytic fungus. F = fresh, Y = young and O= old stumps, see Material
541 and Methods section for description of sampling methods.

Appendix C: Percentage of most abundant operational taxonomic units (OTUs) in samples collected from freshly cut oak stumps. (A): Ascomycota, (B): Basidiomycota. Light grey boxes represent OTUs that overlap with young and old oak stumps.

Appendix D: Percentage of most abundant operational taxonomic units (OTUs) in samples collected from young oak stumps. (A): Ascomycota, (B): Basidiomycota. Light grey boxes represent OTUs that overlap with fresh and old oak stumps; dark grey boxes represent OTUs that overlap with old oak stumps.

Appendix E: Percentage of most abundant operational taxonomic units (OTUs) in samples collected from old oak stumps. (A): Ascomycota, (B): Basidiomycota. Light grey boxes represent OTUs that overlap with fresh and young oak stumps; dark grey boxes represent OTUs that overlap with young oak stumps.

Appendix F: Co-occurrence analysis of sapwood-inhabiting fungal community data using C-score and checkerboard index analysis. * Only OTUs with 3 or more occurrences are included. † Estimated values from 5000 Monte Carlo randomizations. SES = Standardized effect size.

Appendix G: Number of sequence reads per phylum (A=Ascomycota, B = Basidiomycota, unk = unidentified sequence reads, G= Glomeromycota, Z= Zygomycota and C =Chytridiomycota), ratio between Basidiomycota and Ascomycota, H diversity, E evenness, the % Ascomycota and the % mass loss per sample. F=fresh, Y=young and O=old stumps.

Supplements

562 **Supplement 1:** Representative ITS sequences of operational taxonomic units (OTUs) obtained
563 from decaying oak stumps.

564

565

566

esa
preprint

Figure legends

Figure 1: Sampling design for a wood disc collected from an oak tree stump. The wedge-shaped piece (S) was cut out and separated into heartwood, sapwood and bark and for every separate piece the moisture content and density was determined. Drill holes are made to extract sawdust for further lab analyses (see material and methods).

Figure 2: Density of heartwood (red dots, red number indicates the average of the heartwood samples) and sapwood (green dots, green number indicates the average of the sapwood samples) from freshly cut samples, young samples (1.4 years since cutting) and old samples (5.3 years since cutting) of oak stumps.

Figure 3: Percentage of most abundant operational taxonomic units (OTUs) in samples collected from young oak stumps (1.4 years since cutting). (A): Ascomycota, (B): Basidiomycota. Numbers between brackets indicate OTU number (see Appendix B) of OTUs that could not be identified to the species level. On the horizontal axis the percentage of sapwood mass loss per sample is presented. Bars with white dots represent fungal OTUs that overlap between young and old samples.

Figure 4: Percentage of most abundant operational taxonomic units (OTUs) in samples collected from old oak stumps (5.3 years since cutting). (A): Ascomycota, (B): Basidiomycota. Numbers between brackets indicate OTU number (see Appendix B) of OTUs that could not be identified to the species level. On the horizontal axis the percentage of wood mass loss per sample is presented. Bars with white dots represent fungal OTUs that overlap between young and old samples. * indicates that nearly all sapwood was decayed, only about 0.5 g of sapwood could be sampled to perform the DNA extraction.

589 **Figure 5:** Detrended Correspondance Analysis (DCA) biplot showing the variation of the fungal
 590 community composition in sapwood samples taken from oak stumps. Black dots: samples
 591 collected from freshly cut trees, red dots: young samples (1.4 years since cutting), blue dots: old
 592 samples (5.3 years since cutting). Samples closer together have more similar fungal communities.

593

- bark
- sapwood
- heartwood
- drill hole
- S wedge-shaped piece