1
2
3 Title: **Do genetic modifications in crops affect soil fungi? ; a review**
4

5 S.E. Hannula¹, W. de Boer^{1,2} & J.A. van Veen^{1,3}
6

7 ¹ Netherlands Institute of Ecology (NIOO-KNAW), Department of Microbial Ecology,
8 Wageningen, The Netherlands
9

10 ²Wageningen University, Department of Soil Quality, Wageningen, The Netherlands
11

12
13 ³ Insitute of Biology, Leiden University, Leiden, The Netherlands
14
15
16
17
18
19

20 Corresponding author: Emilia Hannula (e.hannula@nioo.knaw.nl); Tel. +31317473507,
21 Fax. +31 317 47 36 75
22 Netherlands Institute of Ecology (NIOO-KNAW),
23 P.O. Box 50, 6708 PB Wageningen, The Netherlands
24
25
26
27
28
29

Abstract

The use of genetically modified (GM) plants in agriculture has been a topic in public debate for over a decade. Despite their potential to increase yields, there may be unintended negative side-effects of GM-plants on soil micro-organisms that are essential for functioning of agro-ecosystems. Fungi are important soil organisms and can have beneficial or harmful effects on plants. Their benefits to agro-ecosystems come from their activities as free-living saprobes breaking down soil organic matter thereby releasing nutrients to the crops, as well as from mutualistic interactions. On the other hand soil-borne plant pathogenic fungi can cause severe damage in crops. Understanding of the impact of GM plants on the dynamics and functioning of soil fungi is essential to evaluate the possible risks of introduction of GM plants for ecosystem functioning. In recent years, over 50 studies have addressed the effects of various GM-traits in crops on soil fungal community structure and function. These studies showed that GM-crops can have positive, negative or neutral effects on both free-living and plant-associated soil fungi. The observed discrepancy in results of these studies is discussed. This is done by highlighting a number of case studies. New methods developed in recent years have enabled microbial ecologists to get a better picture on the functioning and assembly of soil fungal communities. This review presents and discusses two of the most promising methods which are also readily usable in risk assessment of GM-plants on soil fungi and that could help answer remaining key questions in the field.

Keywords: Genetically modified (GM) plants – soil fungi – risk assessment – ‘normal operating range’

Introduction

Although conventional breeding has been successful in developing plants with desired traits, transgenic techniques have extended these possibilities by enabling the introduction of interesting genes from other organisms (Jones 2011). The introduction of new genes into crop-species may increase consistent food production for the growing world population as crop losses due to pests are reduced and optimal crop yields can be obtained. However, there are ethical concerns about the use of transgenic crops as have been discussed in many public forums and have spurred numerous discussions regarding their safety (Jones 2011). Despite these concerns, the number of fields allocated to transgenic crop production has increased each year worldwide (James 2012). However there are strong differences between continents; while in Europe companies are pulling out of the market due to the negative public opinion towards genetically modified (GM)-crops and more strict EU-legislation, growing of GM crops, such as herbicide resistant soybean, is a common practice in USA. The main concerns regarding the use of GM crops in agro-ecosystems are related to the possibility of unintended transgene flow to indigenous plants, development of super weeds, and the effects of transgenic plants on non-target organisms, including soil microbial communities (Wolfenbarger and Phifer 2000).

Effects on composition and activity of soil biota could occur via changes in the chemical composition and quantity of crop residues and rhizodeposits (compounds released by roots) as a result of the modification of the crop. However, growing of different non-GM crop species in crop rotations is nearly always coinciding with changes in the soil microbial communities making the interpretation of the implication of differences in microbial community composition between GM-crops and parental crops complicated (Bruinsma et al. 2003; Liu et al. 2005). The majority of the studies on GM-crop effects on soil microbes have focused on bacteria investigating numbers, activities and community composition whereas only relatively few studies have addressed the impacts on fungi in similar detail despite the importance of fungi for the functioning of soil ecosystems (Carlile et al. 2001).

In the 2003 review by Bruinsma et al (2003) , it was thought that the remaining gaps regarding the evaluation of impact of GM-crops on soil microbes were (1) incomplete knowledge of the functioning of soil microbial communities, (2) poor understanding of the range of responses of the microbial community to “normal” variation in soil systems (such as due to changes in season, weather, and

agricultural management practices including fertilizer use, crop rotation, pesticide use, etc.), and (3) inability to convert complex laboratory procedures to practical assays that are easy to perform and interpret. These knowledge gaps, in particular in the area of effects on fungi, have not yet been filled.

Here, we summarize the information on the effect of GM-plants on activity, biomass and community composition of non-target soil fungi and discuss possibilities to fill the remaining knowledge gaps. We pay special attention to recently developed methods such as next-generation sequencing and stable isotope probing which have the potential, both in their own way, to facilitate the evaluation of the response of soil fungi (and other microbes) to GM-crops.

Fungal diversity and functioning in agro-ecosystems

Before considering the potential effects of genetically modified plants on soil fungi, it is necessary to consider the general effects of agriculture on fungi. Besides the major effects that soil physical and chemical factors such as pH, moisture, soil texture, vegetation have on the composition and functioning of soil microbial communities, many studies have reported a reduction of fungal biomass under agriculture compared to more natural systems, and bacteria are thought to dominate in agricultural ecosystems (Kennedy 1999; Berg and Smalla 2009). This has been attributed both to the constant removal of crop plants, thereby reducing the input of litter (Berg and Smalla 2009) and to mechanical actions such as plowing which can potentially break extensive hyphal networks (Wang et al. 2010). Different agronomic practices such as tillage and fertilization also have profound impacts on the fungal communities (Oehl et al. 2010). Studies have found that organic farming had a significant positive effect on soil AMF richness compared to conventional farming (Verbruggen et al. 2010) and that type of fertilizer applied significantly affects fungal biomass (Heinze et al. 2010). Contradicting the idea of agro-ecosystems being bacterial dominated, recent evidence gained with stable isotope methods show that fungi might be more important organisms in the rhizosphere of crop-species than earlier thought (Gschwendtner et al. 2011; Hannula et al. 2012a). Further evidence suggests that the importance of fungi might be larger later in the season when the plant is more mature and fungi have had more time to establish their hyphal networks (Hannula et al. 2010) while bacteria are more abundant in the rhizosphere of early stages of plant growth and in the bulk soils (Inceoglu et al. 2010).

Fungi perform a wide range of ecosystem functions in agricultural soils and their importance in agro-ecosystems come from their activities as saprobes breaking down soil organic matter thereby releasing nutrients to the crop species as well as from their mutualistic and pathogenic interactions with the plant (Raaijmakers et al. 2009). Thus, a reduced fungal biomass in a field can potentially cause changes in both carbon and nutrient cycling and therefore change the functioning of the system (de Vries et al. 2011). Furthermore, as fungi are in the base of the soil foodweb, a change in either biomass or community composition of fungi caused by a GM-plant can have unforeseen cumulative effects on the higher trophic levels (de Vries et al. 2013).

According to the insurance hypothesis, the loss of biodiversity would cause a reduction in the ecosystem stability due to diminished probability to find species best able adapt to changing conditions (Loreau et al. 2002). The positive effect of increasing fungal biodiversity on stability of the soil ecosystems have been shown for simple communities (Setälä and McLean 2004) but the effect of biodiversity is less evident in natural systems (Nielsen et al. 2011). In this review the effects of GM-crops both on fungal community structure and diversity (and via that to resilience and resistance (Griffiths and Philippot 2013)), and on biomass and function are discussed.

Mechanisms by which GM-plants can affect soil fungi

GM crops can influence soil ecosystems positively, negatively or neutrally, (Oger et al. 1997). Birch et al. (2007) pointed out that the potential impacts of GM crops on soil ecosystem can be (1) direct (e.g. toxicity of an expressed new protein on key non-target species), (2) indirect (e.g. effects via trophic interactions), (3) caused by unintended changes in the metabolism of the plant and thus altering rhizodeposition and/or (4) caused by changes in the management regime used to cultivate GM crops. The effects of GM-trait can potentially be on the fungal biomass, community function or community composition. Harmful effects on fungal community function may cause a decrease in fertility and nutrient cycling in the soil which subsequently affects the following crops (de Vries et al. 2011), and changes in biomass and community structure via elimination of beneficial fungi such as AMF may affect plant growth as well and may result in the increased sensitivity of the plant to pathogens (van der Heijden et al. 2008). GM-caused shifts in

135 fungal community and biomass may also affect higher soil organisms through a cascade of effects in the
136 soil food web in which fungi are a major channel of energy and nutrients.

137 The possible effects discussed in this review are related to GM-induced changes in the chemical
138 composition of living and dead stages of crops that may affect non-target fungi.

141 *Root exudation and soil fungal communities*

142 Rhizodeposition has been identified as an important factor for the development of rhizosphere microbial
143 communities (Lynch and Whipps 1990; Berg and Smalla 2009). A substantial amount of photosynthetically
144 fixed carbon is released into the rhizosphere by roots and the composition and quantity of these exudates
145 differs among plant species and plant growth conditions (Berg and Smalla 2009). Therefore, the first
146 mechanism by which GM-crops can affect soil fungal communities is via intentional or unintentional
147 changes in rhizodeposition quantity and quality. The latter does not only include changes in composition of
148 well-known root-exudates (sugars, organic acids and amino acids) but also the presence of toxins,
149 introduced into the soil from the root. It has been shown that the presence of novel compounds in root
150 derived materials of a transgenic plant may confer a selective advantage to a specific group of soil bacteria
151 which are able to utilize this compound (Savka and Farrand 1997). However such a specific process has not
152 yet been demonstrated for fungi. The effects of toxin releases from roots of Bt-crops and its persistence in
153 the soils has been discussed in detail in an earlier review (Icoz and Stotzky 2008).

154 Several studies have compared GM-crops and parental isoline and other varieties of the same crop
155 species in field trials and greenhouse experiments. Most of these studies have shown that GM-crops do not
156 affect the composition of the free-living soil fungi nor the fungal biomass differently than their parental
157 isolines. Only in five studies significant differences in the soil fungal biomass or community structure
158 between the GM-variety and its parental isoline were observed (Fig. 1). The reason why these studies, and
159 not others have found differences between GM- and their parental isolines remains unclear as there is little
160 in common in these studies; different GM-traits were introduced and different methods were used. Two of
161 the studies were carried out with GM potatoes (increased resistance against nematode and pathogenic
162 bacteria) (Cowgill et al. 2002; Götz et al. 2006) , one with both GM maize and GM potato (Bt and viral

resistance) (Xue et al. 2005), one with GM soybean and GM maize (herbicide tolerance) (Kremer and Means 2009) and the last one with viral resistant GM papaya (Wei et al. 2006). Remarkably, other studies carried out with the same crops and same modifications showed no effect on fungi (Table 1). For example, the study by Kremer and Means (2009) found that frequency of potential plant pathogenic fungi (*Fusaria*) colonizing glyphosate resistant maize roots was higher than in the roots of the parental cultivar, whereas in the same year Hart et al. (2009) reported no differences in abundance or community structure of rhizosphere fungi between the same parental and GM-varieties. This lack of coherence in results is probably due to the large variety in interactions between crops and biotic and abiotic factors such as soil type, plant growth stage studied, climate and interaction with other soil organisms. All these factors are known to strongly influence dynamics of soil fungi. Furthermore, in seven studies differences in fungal biomass or community structure were found between GM- and parental varieties, but due to the large variation in time and space, these effects were deemed transient (Fig 1.). These ‘transient’ effects and lack of coherence in results are discussed later in this review.

Despite the importance of arbuscular mycorrhizal fungi (AMF) in plant-soil systems, only few studies evaluated the non-target effects of GM-crops on AMF colonization and community structure. (Liu 2010) (Fig 1). As plants vary naturally in their AMF-hosting ability, the GM trait in plants might, in some cases, alter their relationship with AMF. Because AMF are obligate symbionts and thus require the plant host for nutrition and reproduction, they may be more sensitive to changes in the physiology of the host plant than free-living soil fungi (Liu 2010; Cheeke et al. 2011). Earlier, it was shown that AMF are sensitive to different agronomic practices such as tillage and fertilization (Oehl et al. 2010). AMF are thought to be especially important in low input agro-ecosystems and are, therefore, an important component of sustainable agriculture (Verbruggen and Kiers 2010) and it is therefore crucial to understand the impacts of GM-traits on functioning and diversity of AMF. In two studies a reduction in arbuscular mycorrhizal colonization of the roots of a Bt-cultivar of maize was reported (Castaldini et al. 2005; Cheeke et al. 2012). Another study found no effect on colonization of the Bt-maize roots but an effect on the AM-fungal community structure assessed by DGGE (Tan et al. 2011). Transient effects of GM-crops on soil AMF community structure or AMF colonization of roots have been reported in 4 other studies on Bt-maize, starch modified potatoes and herbicide tolerant soybean (Turrini et al. 2004; Powell et al. 2007; Cheeke et

al. 2011; Hannula et al. 2012a). In the study by Powell et al. (2007) different levels of rhizobial and mycorrhizal colonization were observed between conventional and GM- soybeans. However, these differences could be attributed to variation found between the three different non-modified and six different modified cultivars and not the GM status of the plant. Other studies did not find effects of the GM-modification on any aspect of AM biology studied. For instance, four different modifications introducing insect resistance or herbicide tolerance in cotton had no effect on AMF colonization (Knox et al. 2008). In addition, de Vauflleury *et al.* (2007) did not find any significant effect of Cry1Ab (Bt-) modification of maize on AMF colonization. However, the total number of studies about effects of GM-crops on AMF is rather low and certain traits such as the herbicide tolerance and resistance to pathogens have been only studied in two studies (Table 1) which makes it difficult to come to a definite conclusion about the effects of GM-crops on the AMF community.

Effect of GM plants on residue decomposition and decomposer fungi

Decomposition of litter is a key function in the cycling of elements and, consequently, in mineral nutrient supply to plants thus any change in plant litter composition may potentially significantly affect soil functioning (Deacon et al. 2006; Berg and McClaugherty 2008; van der Wal et al. 2013). In general, fungi are more significant as litter-decaying agents than bacteria (Deacon et al. 2006). However, the large bulk of the relevant studies have addressed litter decomposition as a functional response to GM-traits without referring explicitly to the fungal communities involved.

The Bt-varieties of corn, cotton and rice have been the most studied modifications in litter decomposition due to the observed unintended effect of Cry1Ab on the lignin content of the plants (Saxena and Stotzky 2001b). Slower decomposition resulting from this altered lignin concentration has been reported in few studies (Castaldini et al. 2005; Flores et al. 2005) while a greater number of studies did not find a difference in decomposition between Bt and non-Bt corn (Jung and Sheaffer 2004; Fang et al. 2007; Zwahlen et al. 2007; Daudu et al. 2009; Zurbrugg et al. 2010). An early study on Cry1A expression in cotton found more species of fungi based on colony counting in the soils incubated with transgenic leaves than in the soil incubated with leaves from the parental variety (Donegan et al. 1995). However, this study seems to be an exception as it is the only one in which significant differences between the GM crop and the

parental isolate could not be explained by other factors than the genetic modification. The majority of studies on fungi in decomposing plant material did not show any significant effect or only a transient effect of genetic modifications on certain aspects of the fungal community (Fig.1). One of these studies on Cry3Bb expressing Bt-corn in a field experiment revealed no difference in the decomposition rate of roots, stalks, cobs or leaves between the Bt- and its parental variety at different locations but did detect a significant difference in fungal community composition as determined by T-RLFP in one of the soils tested and in one year which points to the transient nature of the observed effect (Xue et al. 2011). Other studies detected effects at one or more time points during decomposition but not for the overall decomposer community dynamics or for the total amount of plant material that was decomposed (Wu et al. 2004; Castaldini et al. 2005; Lu et al. 2010a).

As for AMF, it should be noted that effects of genetic modifications on decomposer fungi have only been addressed for a limited number of modifications and majority of the studies have investigated the effects of Bt-modifications (Table 1). Since modifications of pathogen resistance and structural changes of plant parts would be the most obvious GM-traits to affect the non-target decomposer fungal communities, it is surprising that no decomposition studies have addressed pathogen-resistance related modifications and only three dealt with the effect of structural changes of GM plants (lignin synthesis in tobacco, chitinase in birch and starch in potatoes) (Henault et al. 2006; Seppänen et al. 2007; Hannula et al. 2013). In the case of plants with genetic modifications to structural parts such as lignin synthesis or starch quality, risk assessment studies taking into account the effects on soil microbes and processes are essential.

Normal variation versus GM-induced variation

A common issue in the debate and a possible explanation for the lack of coherence in the results obtained in the previous studies on possible, harmful, side-effects of GM-crops is the difficulty to discern the effects of the modification from all the other abiotic and biotic factors (Fig.2). Usually 'normal variation' is defined as the variation in the responses of the fungal/microbial community to the non-GM crops under the prevailing conditions of the common agricultural practices (Kowalchuk et al. 2003). Factors such as weather, agricultural management and plant developmental stage can affect the outcome of the experiments more than the genetic modification (Griffiths et al. 2000; Lukow et al. 2000; Dunfield and Germida 2001).

For example, Hannula et al. (2012b) investigated the impact of different potato cultivars, including a GM amylopectin-accumulating potato line, on rhizosphere fungal communities over a period of three years under field conditions using molecular microbiological methods; they revealed occasional differences between the transgenic line and its parental variety, indicating that differences, if realistic at all, were mainly transient in nature and could only be detected either in one soil, at one growth stage or over a one-year period. Furthermore, decomposition of plant material is also affected by the soil type and burying depth of the tested plant material (Holland and Coleman 1987; Burgess et al. 2002; Powell et al. 2009).

The first variables to consider are site related variables. In general, soil type and field conditions, including the history of the site are considered to be among the most influential factors governing soil fungal community structure and function directly and indirectly via better plant growth (Costa et al. 2006; Singh et al. 2007; Wang et al. 2009; Chaparro et al. 2012). In case of Bt-modifications it is known that the physicochemical and biological characteristics of soils may influence the persistence of Cry class proteins (Icoz and Stotzky 2008) in the environment thus influencing the outcome of the studies. Unfortunately, only relatively few studies addressing possible effects of GM crops on soil fungal communities have included more than one soil type (Fig. 2). For instance, Blackwood and Buyer (2004) investigated the effects of Bt-modified maize on soil fungi in three soils and found that the soil type, but not the modification, had a significant effect on the fungal biomass. Furthermore, the decomposition of litter is found to vary between sites. Indeed, studies on Bt maize and rice have shown that both the site and the burying depth are very important factors governing the decomposer processes and the structure of associated fungal communities (Cortet et al. 2006; Lu et al. 2010b; Xue et al. 2011).

Few studies have compared effects of agricultural management practices in combination with GM-crops on soil fungi (Fig. 2). Cheeke et al. (2011) inoculated AMF *Glomus mosseae* in Bt- maize and parental roots and found that there was a significant interaction effect of cultivar and fertilizer level. The effect of the GM-trait could only be seen in the low or no fertilizer treatments but not in the high fertilizer treatment. Yet, this is an important aspect to consider as AMF are thought to be more beneficial to the plant in low-input agro-ecosystems.

The growth stage of the plant is a second factor determining the activity and community structure of fungi in soil. Jones et al. (2004) indicated that the amount and composition of rhizodeposition changes

during plant development with important consequences for the microbial activity and community composition in the rhizosphere. Indeed, this seems to be valid for saprophytic fungi as well (although their role in the rhizosphere is still a matter of debate) as plant growth stage and sampling time were found to have the largest effect on activity and composition of both fungi in general and AMF in many experiments (Fig. 2). The effect of growth stage was not seen in the bulk soil (Milling et al. 2004) or in the AMF community under a tree (aspen) (Kaldorf et al. 2002) but was apparent in all other studies in which the stage was evaluated. For example studies on genetically modified potatoes (Donegan et al. 1995; Cowgill et al. 2002; Weinert et al. 2009; Gschwendtner et al. 2010; Hannula et al. 2012b) have shown that growth stage is the single most important factor affecting the fungal biomass and community structure in the rhizosphere. In field trials effects of growth stage can be affected by coinciding changes in temperature and water availability, which are both important determinants of microbial growth. However, greenhouse experiments have shown that in controlled conditions there is an effect, although smaller than in the field, of plant growth stage on soil fungal communities (Girlanda et al. 2008; Wu et al. 2009; Gschwendtner et al. 2011). Finally, there is emerging evidence that plant parts collected at different stages of growth, also decompose differently (Zurbrügg et al. 2010) and might, thus, also have different effects on fungal communities.

Annual variation, including climatic factors such as precipitation and temperature often explains large part of the variation observed in decomposer experiments (Fig. 2). From 11 studies dealing with the effects of GM-crops on soil fungi in which annual variation was accounted for, 9 observed differences in fungal community composition or abundance between years whereas in 2 studies no annual variation was apparent (Milling et al. 2004; Li et al. 2011). An elegant field study during 3 years revealed that ‘year’ was the strongest explaining factor for changes in decomposition rate and structure of the associated fungal communities and far more important than the Bt-trait of maize (Xue et al. 2011). In a four year field study of Bt-corn, ‘year’ was shown to be a highly significant explanatory factor while the Bt-and its parental variety differed only in one of the years and thus had a transient effect on numbers of culturable soil fungi (Icoz et al. 2008).

The last important factor to consider when evaluating the effects of GM-traits on soil fungi is the variation in traits that affect the soil microbial community among cultivars that exists due to their long

history of breeding. This may explain some of the transient effects observed in those studies that have compared multiple GM-varieties or multiple ‘normal’ varieties against the GM (Fig. 2). In most cases it was found that the normal variation among cultivars and thus the variable impact of conventionally bred varieties on environment was larger than the difference between GM-variety and its parental cultivar. Icoz et al. (2008) compared 4 Bt-varieties and their corresponding parental isolines and observed that the Bt-modification did not have an effect on numbers of fungi while crop variety had a significant albeit transient effect on the soil fungal community. Cheeke et al. (2012) investigated the colonization percentages of AMF on 16 maize lines (9 Bt lines and 7 parental varieties) and were the first to find a significant relationship between Bt-trait and the (lowered) colonization of AMF in a multiple cultivar study. However, this pattern seemed to hold only for greenhouse conditions as in the field study with a subset of these cultivars no effect on AMF colonization was found (Cheeke et al. 2013). This further highlights the importance of including environmental factors when evaluating GM effects on AMF.

New methods and new possibilities in GM-research

Traditionally most of the studies on effects of GM-crops on fungi have used cultivation based methods and root colonization counts of AMF (Table 1) to assess the effects of GM crops on the size and the diversity of the fungal community. However, the inherent limitations of the studying of culturable microbes have also here prevented a full scale assessment of the effects of GM crops on fungal diversity and functioning. In few cases DNA based fingerprinting methods such as denaturing gradient gel electrophoresis (DGGE) (Milling et al. 2004; Götz et al. 2006) or terminal restriction length polymorphism (T-RFLP) (Hart et al. 2009) were applied. However, in order to answer fundamental questions still open in this field such as on the totality of the effects of GM-crops on soil fungal diversity and in particular the rare and non-culturable fractions of the community as well as the functioning of the community through the impact of changes in rhizodeposition patterns, new methodology is ready to be adopted. In this section we will discuss methodologies that could help answering these key questions.

Monitoring differences in root exudation patterns with stable isotope probing (SIP)

Many studies have reported on the differences in the community composition of rhizosphere fungi between cultivars and have hypothesized that this would have been the result of changes in rhizodepositions. However, only few studies have actually measured rhizodepositions or monitored carbon flow from the plant into the rhizosphere, the subsequent utilization of the rhizodepositions by the microbial community, and how this influence both the structure and the functioning of the rhizosphere community. An elegant approach to do such measurements is stable isotope probing (SIP,) where the whole plant is (pulse) labeled with $^{13}\text{CO}_2$ and the incorporation of ^{13}C in microbes is followed in the endosphere and rhizosphere. Alternatively, plant residues containing ^{13}C can be used in a decomposition study to monitor the soil fungi involved in breakdown of the (GM)-plant material. The measurement of the isotope in the DNA/RNA or fatty acids (PLFAs) extracted from soil allows for the detection and identification of the microbes actively involved in the assimilation of the labeled compound i.e. ^{13}C from the plant roots or dead plant material (Radajewski et al. 2000). SIP can be used to follow the fate of carbon in any system and has been used to study for example effects of soil management (Rangel-Castro et al. 2005; Lu et al. 2007) and climate change (Drigo et al. 2010) on soil microbial community structure and is proven to be a robust technique which can provide a quantitative insight in the rhizodeposit metabolizing community. In combination with phospholipid fatty acid (PLFA-SIP) analyses it has been used to evaluate the effects of GM-plants on carbon partitioning to different groups of soil organisms (Wu et al. 2009; Gschwendtner et al. 2011; Hannula et al. 2012a). All of these studies have shown the great importance of both saprotrophic fungi and especially AMF in the rhizosphere assimilating the (^{13}C) from the plant. The first two studies did not find significant differences between the GM-and its parental cultivar, although Wu et al. (2009) found significant differences between the Bt and parental rice in the amount of ^{13}C distribution at the seedling, booting and heading stages. The last study (Hannula et al. 2012a) found that a starch modified GM-potato line affected soil fungal communities slightly differently than its parental isolate did, but these effects were deemed transient. A study done with DNA-SIP revealed cultivar dependent distinctions in ^{13}C -label flow to endophytic bacteria of potato (Rasche et al. 2009). However, in these studies the baseline of environmental variation was not investigated and thus it is not clear whether these differences are ecologically relevant. SIP methodology will, nevertheless, offer a great opportunity to study the effects of GM-varieties on active members of rhizosphere communities.

Possibilities of high-throughput sequencing to reveal fine scale differences between GM- and parental variety

It has been recognized that in addition to using broad scale keystone indicators such as fungal biomass and community composition, there is a need to improve the sensitivity of detection methods for detailed analyses of the impacts of GM-crops on soil microbial communities (Lilley et al. 2006). This should further target relevant species and functions for each combination of modification and species. Earlier, microbial biodiversity was thought to be a very sensitive parameter to perturbation and a good indicator for soil functioning (van der Heijden et al. 1998; Kennedy 1999; Garbeva et al. 2004) but this has recently been debated (Prosser 2012). In earlier studies in which differences in fungal community structure have been found, often these changes have been small without clear influence on the functioning of the system. However, it is not known how rare microbes affect the functioning of the system and how a change in community structure and diversity affects its function (Nielsen et al. 2011). Modern molecular methods such as 454-sequencing are useful in evaluating the effects of GM-crops on soil diversity replacing the DGGE and T-RFLP methods (Lindahl et al. 2013). A recent study on the effects of Bt maize varieties on AMF communities done using 454-sequencing combined with T-RFLP analyses revealed a significant correlation between the two methods even though some of the relative abundances of individual taxa differed (Verbruggen et al. 2012). Thus, there is no need to repeat existing evaluations of various GM-crops with this new methodology but the method itself is a powerful tool for future studies on the effects of GM-crops on soil fungal community structure and diversity. However, changes in diversity and community structure might not always lead to changes in ecosystem function as the relationship between diversity of soil micro-organisms and soil ecosystem functioning remain unclear (Nielsen et al. 2011). Thus, functional parameters or indicators of community functioning (i.e. production of extracellular enzymes) should be measured parallel to the diversity estimates. These functional measurements are discussed in an earlier review in more detail (Bruinsma et al. 2003). In future, function based sequencing and meta-transcriptomics can be used to when evaluating the effects of the GM-crops on soil fungi and will give a more detailed insight into the functional consequences of the modifications.

386 *Final remarks*

387 One issue, that has been addressed only marginally in GM crop studies, but in our opinion highly relevant
388 to evaluate the real risk associated with the use of GM crops, is the effects of GM crops after harvest and/or
389 in the following growing season(s). Few examples of such studies are a study by Castaldini *et al.* (2005),
390 who reported a lower AMF colonization of *Medicago sativa* (alfalfa) roots grown in pots that had
391 previously been cultivated with Bt maize and contained extra Bt plant residues. Contradictory, Cheeke *et al.* 2012 did not find any residual effect of Bt maize cultivation on AMF colonization of subsequently
392 planted *Glycine max* (soybean). Besides, other studies in field and greenhouse conditions could not detect
393 an effect of GM-crops on soil fungi after their harvest and/or in the following growing season (Oliveira *et al.* 2008; Powell *et al.* 2009; Cheeke *et al.* 2012; Hannula *et al.* 2012b). We strongly argue that such
394 measurements should be considered in future risk assessment studies on the use of GM crops especially in
395 cases when GM-crops are grown consecutively in the same soils for years as continuous growing of GM-
396 crops can cause additive effect may strengthen the (negative) effects of the transgene.

399 The results available on the impact of GM plants on natural and agricultural ecosystems show that
400 specific effects of single transformation events should be tested on a case-by-case basis in a natural setting
401 where the baseline factors are all taken into the consideration, including biochemical, physiological, and
402 molecular parameters. As fungi are important to soil functioning and plant growth, an evaluation of these
403 organisms should be performed when evaluating the effects on soil biota. Furthermore, there is a need for
404 statistical methods which can evaluate the effects of GM-trait in relation to the baseline 'noise' in the
405 system. The new techniques such as SIP-experiments and high throughput sequencing and
406 metatranscriptomics should be used in parallel with carefully designed field experiments considering all the
407 'baseline' factors including effects on the subsequent crop species.

408
409 **Acknowledgements**

410 This review was financed by ERGO grant number 838.06.052 of the Netherlands Organization for Scientific
411 Research. We thank anonymous reviewers for their insightful comments that greatly improved the manuscript
412 This is publication 5563 of the Netherlands Institute of Ecology (NIOO-KNAW).

Legends for figures:

Figure 1 Studies on the effect of GM-crops on general fungal communities arbuscular mycorrhizal fungal (AMF) communities and decomposer fungal communities. This classification was done for convenience and because of the large differences between the types of studies and organisms. The observed effect of GM crop was categorized as i) a (lasting) effect, ii) a transient effect or iii) no effect on fungi. In the mycorrhizal studies, only effects on AMF were included; the few studies addressing effects of lignin-modified trees on ectomycorrhiza were not included. Despite the presumed role of fungi in decomposition processes, only studies actually measuring fungal activity were included. Further, studies based on fatty acid analysis (FAME, PLFA) were not included in AMF studies. For more details, see text.

Figure 2. Published studies on GM that have included the effect of other parameters than GM-trait on fungal community composition. List of studies which investigated each parameter and either found an effect (darker color) or did not detect an effect (lighter color). In the first row the studies detecting a significant effect of GM are marked with black, no effect with light gray and ‘transient’ effect with dark gray. Next rows are the effects of plant growth stage, field site and soil related parameters, season and climate and cultivar. For details on these categories, see text. Darker color marks that this factor was a significant explanatory factor in the study while lighter color marks that factor was studied but no effect was found. The totals are total number of studies looking at the factor (and studies in which an effect of the factor in question was detected). Some studies are featured many times in the table as they have looked at multiple aspects.

References:

- Berg B, McClaugherty C (2008) Plant Litter: Decomposition, Humus Formation, Carbon Sequestration (second ed.). Springer, Berlin; New York
- Berg G, Smalla K (2009) Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. FEMS Microbiol Ecol 68 (1):1-13
- Birch ANE, Griffiths BS, Caul S, Thompson J, Heckmann LH, Krogh PH, Cortet J (2007) The role of laboratory, glasshouse and field scale experiments in understanding the interactions between genetically modified crops and soil ecosystems: A review of the ECOGEN project. Pedobiologia 51 (3):251-260
- Blackwood CB, Buyer JS (2004) Soil microbial communities associated with Bt and non-Bt corn in three soils. J Environ Qual 33:832-836
- Bruinsma M, Kowalchuk GA, van Veen JA (2003) Effects of genetically modified plants on microbial communities and processes in soil. Biol Fertility Soils 37 (6):329-337
- Burgess MS, Mehuys GR, Madramootoo CA (2002) Decomposition of grain-corn residues (*Zea mays* L.): A litterbag study under three tillage systems. Can J Soil Sci 82 (2):127-138
- Carlile MJ, Watkinson SC, Gooday GW (2001) The Fungi. 2nd edn. Academic Press, San Diego, CA, USA
- Castaldini M, Turrini A, Sbrana C, Benedetti A, Marchionni M, Mocali S, Fabiani A, Landi S, Santomassimo F, Pietrangeli B, Nuti MP, Miclaus N, Giovannetti M (2005) Impact of Bt corn on rhizospheric and soil eubacterial communities on beneficial mycorrhizal symbiosis in experimental microcosms. Appl Environ Microbiol 71 (11):6719-6729. doi:10.1128/aem.71.11.6719-6729.2005
- Chaparro JM, Sheflin AM, Manter DK, Vivanco JM (2012) Manipulating the soil microbiome to increase soil health and plant fertility. Biol Fertility Soils 48 (5):489-499. doi:10.1007/s00374-012-0691-4
- Cheeke TE, Pace BA, Rosenstiel TN, Cruzan MB (2011) The influence of fertilizer level and spore density on arbuscular mycorrhizal colonization of transgenic Bt 11 maize (*Zea mays*) in experimental microcosms. FEMS Microbiol Ecol 75 (2):304-312. doi:10.1111/j.1574-6941.2010.01013.x
- Cheeke TE, Rosenstiel TN, Cruzan MB (2012) Evidence of reduced arbuscular mycorrhizal fungal colonization in multiple lines of Bt maize. Am J Bot 99 (4):700-707. doi:10.3732/ajb.1100529
- Cheeke TE, Cruzan MB, Rosenstiel TN (2013) Field evaluation of arbuscular mycorrhizal fungal colonization in *Bacillus thuringiensis* toxin-expressing (Bt) and non-Bt maize. Appl Environ Microbiol 79 (13):4078-4086. doi:10.1128/aem.00702-13
- Chun Y, Kim H-J, Park K, Jeong S-C, Lee B, Back K, Kim H, Kim C-G (2012) Two-year field study shows little evidence that PPO-transgenic rice affects the structure of soil microbial communities. Biol Fertility Soils 48 (4):453-461. doi:10.1007/s00374-011-0626-5
- Cortet J, Andersen MN, Caul S, Griffiths B, Joffre R, Lacroix B, Sausse C, Thompson J, Krogh PH (2006) Decomposition processes under Bt (*Bacillus thuringiensis*) maize: Results of a multi-site experiment. Soil Biol Biochem 38 (1):195-199
- Costa R, Gotz M, Mrozek N, Lottmann J, Berg G, Smalla K (2006) Effects of site and plant species on rhizosphere community structure as revealed by molecular analysis of microbial guilds. FEMS Microbiol Ecol 56 (2):236-249
- Cowgill SE, Bardgett RD, Kiezebrink DT, Atkinson HJ (2002) The effect of transgenic nematode resistance on non-target organisms in the potato rhizosphere. J Appl Ecol 39 (6):915-923
- Daudu CK, Muchaonyerwa P, Mkeni PNS (2009) Litterbag decomposition of genetically modified maize residues and their constituent *Bacillus thuringiensis* protein (Cry1Ab) under field conditions in the central region of the Eastern Cape, South Africa. Agric Ecosyst Environ 134 (3-4):153-158. doi:10.1016/j.agee.2009.06.012
- de Vauflery A, Kramarz PE, Binet P, Cortet J, Caul S, Andersen MN, Plumey E, Coeurdassier M, Krogh PH (2007) Exposure and effects assessments of Bt-maize on non-target organisms (gastropods, microarthropods, mycorrhizal fungi) in microcosms. Pedobiologia 51 (3):185-194. doi:10.1016/j.pedobi.2007.04.005
- de Vries FT, van Groenigen JW, Hoffland E, Bloem J (2011) Nitrogen losses from two grassland soils with different fungal biomass. Soil Biol Biochem 43 (5):997-1005. doi:<http://dx.doi.org/10.1016/j.soilbio.2011.01.016>

- de Vries FT, Thébault E, Liiri M, Birkhofer K, Tsiafouli MA, Bjørnlund L, Bracht Jørgensen H, Brady MV, Christensen S, de Ruiter PC, d'Hertefeldt T, Frouz J, Hedlund K, Hemerik L, Hol WHG, Hotes S, Mortimer SR, Setälä H, Sgardelis SP, Uteseny K, van der Putten WH, Wolters V, Bardgett RD (2013) Soil food web properties explain ecosystem services across European land use systems. *Proc Natl Acad Sci USA* 110 (35):14296-14301. doi:10.1073/pnas.1305198110
- Deacon LJ, Pryce-Miller EJ, Frankland JC, Bainbridge BW, Moore PD, Robinson CH (2006) Diversity and function of decomposer fungi from a grassland soil. *Soil Biol Biochem* 38 (1):7-20
- Donegan KK, Palm CJ, Fieland VJ, Porteous LA, Ganio LM, Schaller DL, Bucalo LQ, Seidler RJ (1995) Changes in levels, species and DNA fingerprints of soil-microorganisms associated with cotton expressing the *Bacillus-Thuringiensis* var *Kurstaki* endotoxin. *Applied Soil Ecology* 2 (2):111-124
- Donegan KK, Schaller DL, Stone JK, Ganio LM, Reed G, Hamm PB, Seidler RJ (1996) Microbial populations, fungal species diversity and plant pathogen levels in field plots of potato plants expressing the *Bacillus thuringiensis* var *tenebrionis* endotoxin. *Transgenic Res* 5 (1):25-35
- Donegan KK, Seidler RJ, Doyle JD, Porteous LA, Digiovanni G, Widmer F, Watrud LS (1999) A field study with genetically engineered alfalfa inoculated with recombinant *Sinorhizobium meliloti*: effects on the soil ecosystem. *J Appl Ecol* 36 (6):920-936
- Drigo B, Pijl AS, Duyts H, Kielak A, Gamper HA, Houtekamer MJ, Boschker HTS, Bodelier PLE, Whiteley AS, van Veen JA, Kowalchuk GA (2010) Shifting carbon flow from roots into associated microbial communities in response to elevated atmospheric CO₂. *Proc Natl Acad Sci USA* 107 (24):10938-10942. doi:10.1073/pnas.0912421107
- Dunfield KE, Germida JJ (2001) Diversity of bacterial communities in the rhizosphere and root interior of field-grown genetically modified *Brassica napus*. *FEMS Microbiol Ecol* 38 (1):1-9
- Fang M, Motavalli PP, Kremer RJ, Nelson KA (2007) Assessing changes in soil microbial communities and carbon mineralization in Bt and non-Bt corn residue-amended soils. *Appl Soil Ecol* 37 (1-2):150-160. doi:10.1016/j.apsoil.2007.06.001
- Fließbach A, Messmer M, Nietlispach B, Infante V, Mader P (2012) Effects of conventionally bred and *Bacillus thuringiensis* (Bt) maize varieties on soil microbial biomass and activity. *Biol Fertility Soils* 48 (3):315-324. doi:10.1007/s00374-011-0625-6
- Flores S, Saxena D, Stotzky G (2005) Transgenic Bt plants decompose less in soil than non-Bt plants. *Soil Biol Biochem* 37 (6):1073-1082
- Garbeva P, van Veen JA, van Elsas JD (2004) Microbial diversity in soil: Selection of microbial populations by plant and soil type and implications for disease suppressiveness. *Annu Rev Phytopathol* 42:243-270. doi:10.1146/annurev.phyto.42.012604.135455
- Girlanda M, Bianciotto V, Cappellazzo GA, Casieri L, Bergero R, Martino E, Luppi AM, Perotto S (2008) Interactions between engineered tomato plants expressing antifungal enzymes and nontarget fungi in the rhizosphere and phyllosphere. *FEMS Microbiol Lett* 288 (1):9-18. doi:10.1111/j.1574-6968.2008.01306.x
- Götz M, Nirenberg H, Krause S, Wolters H, Draeger S, Buchner A, Lottmann J, Berg G, Smalla K (2006) Fungal endophytes in potato roots studied by traditional isolation and cultivation-independent DNA-based methods. *FEMS Microbiol Ecol* 58 (3):404-413
- Griffiths BS, Geoghegan IE, Robertson WM (2000) Testing genetically engineered potato, producing the lectins GNA and Con A, on non-target soil organisms and processes. *J Appl Ecol* 37:159-170
- Griffiths BS, Philippot L (2013) Insights into the resistance and resilience of the soil microbial community. *FEMS Microbiol Rev* 37 (2):112-129. doi:10.1111/j.1574-6976.2012.00343.x
- Gschwendtner S, Reichmann M, Muller M, Radl V, Munch JC, Schlöter M (2010) Effects of genetically modified amylopectin-accumulating potato plants on the abundance of beneficial and pathogenic microorganisms in the rhizosphere. *Plant Soil* 335 (1-2):413-422. doi:10.1007/s11104-010-0430-2
- Gschwendtner S, Esperschütz J, Buegger F, Reichmann M, Müller M, Munch JC, Schlöter M (2011) Effects of genetically modified starch metabolism in potato plants on photosynthate fluxes into the rhizosphere and on microbial degraders of root exudates. *FEMS Microbiol Ecol* 76 (3):564-575. doi:10.1111/j.1574-6941.2011.01073.x
- Hannula SE, de Boer W, van Veen JA (2010) In situ dynamics of soil fungal communities under different genotypes of potato, including a genetically modified cultivar. *Soil Biol Biochem* 42 (12):2211-2223. doi:DOI: 10.1016/j.soilbio.2010.08.020
- Hannula SE, Boschker HTS, de Boer W, van Veen JA (2012a) ¹³C pulse-labeling assessment of the community structure of active fungi in the rhizosphere of a genetically starch-modified potato

- (*Solanum tuberosum*) cultivar and its parental isoline. *New Phytol* 194 (3):784-799.
doi:10.1111/j.1469-8137.2012.04089.x
- Hannula SE, de Boer W, van Veen J (2012b) A 3-year study reveals that plant growth stage, season and field site affect soil fungal communities while cultivar and GM-trait have minor effects. *PloS ONE* 7 (4):e33819. doi:10.1371/journal.pone.0033819
- Hannula SE, de Boer W, Baldrian P, Van Veen JA (2013) Effects of genetically modified amylopectin-accumulating potato in decomposer processes and fungal diversity in litter and soil. *Soil Biol Biochem* 58:88-98
- Hart MM, Powell JR, Gulden RH, Dunfield KE, Pauls KP, Swanton CJ, Klironomos JN, Antunes PM, Koch AM, Trevors JT (2009) Separating the effect of crop from herbicide on soil microbial communities in glyphosate-resistant corn. *Pedobiologia* 52 (4):253-262.
doi:10.1016/j.pedobi.2008.10.005
- Heinze S, Raupp J, Joergensen RG (2010) Effects of fertilizer and spatial heterogeneity in soil pH on microbial biomass indices in a long-term field trial of organic agriculture. *Plant Soil* 328 (1-2):203-215. doi:10.1007/s11104-009-0102-2
- Henault C, English LC, Halpin C, Andreux F, Hopkins DW (2006) Microbial community structure in soils with decomposing residues from plants with genetic modifications to lignin biosynthesis. *FEMS Microbiol Lett* 263 (1):68-75. doi:10.1111/j.1574-6968.2006.00416.x
- Holland EA, Coleman DC (1987) Litter placement effects on microbial and organic-matter dynamics in an agroecosystem. *Ecology* 68 (2):425-433. doi:10.2307/1939274
- Icoz I, Saxena D, Andow DA, Zwahlen C, Stotzky G (2008) Microbial populations and enzyme activities in soil in situ under transgenic corn expressing Cry proteins from *Bacillus thuringiensis*. *J Environ Qual* 37:647-662
- Icoz I, Stotzky G (2008) Fate and effects of insect-resistant Bt crops in soil ecosystems. *Soil Biol Biochem* 40 (3):559-586
- Inceoglu O, Salles JF, van Overbeek L, van Elsas JD (2010) Effects of plant genotype and growth stage on the betaproteobacterial communities associated with different potato cultivars in two fields. *Appl Environ Microbiol* 76 (11):3675-3684. doi:10.1128/aem.00040-10
- James C (2012) Global Status of Commercialized Biotech/GM Crops: 2012. ISAAA Brief No. 44. . International Service for the Acquisition of Agri-Biotech Applications, Ithaca, NY
- Jones DL, Hodge A, Kuzyakov Y (2004) Plant and mycorrhizal regulation of rhizodeposition. *New Phytol* 163:459-480
- Jones JDG (2011) Why genetically modified crops? *Phil Trans R Soc A* 369 (1942):1807-1816.
doi:10.1098/rsta.2010.0345
- Jung HG, Sheaffer CC (2004) Influence of Bt transgenes on cell wall lignification and digestibility of maize stover for silage. *Crop Sci* 44 (5):1781-1789
- Kaldorf M, Fladung M, Muhs H-JM, Buscot F (2002) Mycorrhizal colonization of transgenic aspen in a field trial. *Planta* 214 (4):653-660. doi:10.1007/s004250100658
- Kennedy AC (1999) Bacterial diversity in agroecosystems. *Agric Ecosyst Environ* 74 (1-3):65-76
- Knox OGG, Nehl DB, Mor T, Roberts GN, Gupta VVSR (2008) Genetically modified cotton has no effect on arbuscular mycorrhizal colonisation of roots. *Field Crops Res* 109 (1-3):57-60.
doi:10.1016/j.fcr.2008.06.005
- Kowalchuk GA, Bruinsma M, van Veen JA (2003) Assessing responses of soil microorganisms to GM plants. *Trends Ecol Evol* 18 (8):403-410
- Kremer RJ, Means NE (2009) Glyphosate and glyphosate-resistant crop interactions with rhizosphere microorganisms. *European Journal of Agronomy* 31 (3):153-161. doi:10.1016/j.eja.2009.06.004
- Kuramae EE, Verbruggen E, Hillekens R, de Hollander M, Roling WFM, van der Heijden MGA, Kowalchuk GA (2013) Tracking Fungal Community Responses to Maize Plants by DNA- and RNA-Based Pyrosequencing. *PloS ONE* 8 (7). doi:10.1371/journal.pone.0069973
- Lawhorn CN, Neher DA, Dively GP (2009) Impact of coleopteran targeting toxin (Cry3Bb1) of Bt corn on microbially mediated decomposition. *Applied Soil Ecology* 41 (3):364-368.
doi:10.1016/j.apsoil.2008.12.003
- Lee S-H, Kim C-G, Kang H (2011) Temporal dynamics of bacterial and fungal communities in a genetically modified (GM) rice ecosystem. *Microb Ecol* 61 (3):646-659. doi:10.1007/s00248-010-9776-5

- Li X, Liu B, Cui J, Liu D, Ding S, Gilna B, Luo J, Fang Z, Cao W, Han Z (2011) No evidence of persistent effects of continuously planted transgenic insect-resistant cotton on soil microorganisms. *Plant Soil* 339 (1):247-257. doi:10.1007/s11104-010-0572-2
- Lilley AK, Bailey MJ, Cartwright C, Turner SL, Hirsch PR (2006) Life in earth: the impact of GM plants on soil ecology? *Trends Biotechnol* 24 (1):9-14
- Lindahl BD, Nilsson RH, Tedersoo L, Abarenkov K, Carlsen T, Kjølner R, Kõljalg U, Pennanen T, Rosendahl S, Stenlid J, Kauserud H (2013) Fungal community analysis by high-throughput sequencing of amplified markers – a user's guide. *New Phytol* 199 (1):288-299. doi:10.1111/nph.12243
- Liu B, Zeng Q, Yan FM, Xu HG, Xu CR (2005) Effects of transgenic plants on soil microorganisms. *Plant Soil* 271 (1-2):1-13
- Liu W, Hao Lu H, Wu W, Kun Wei Q, Xu Chen Y, Thies JE (2008) Transgenic Bt rice does not affect enzyme activities and microbial composition in the rhizosphere during crop development. *Soil Biol Biochem* 40 (2):475-486
- Liu WK (2010) Do genetically modified plants impact arbuscular mycorrhizal fungi? *Ecotoxicology* 19 (2):229-238. doi:10.1007/s10646-009-0423-1
- Loreau M, Downing A, Emmerson M, Gonzalez A, Hughes J, Inchausti P, Joshi J, Norberg J, Sala O (2002) A new look at the relationship between diversity and stability. In: Naeem S (ed) *Biodiversity and ecosystem functioning : synthesis and perspectives* Oxford University Press Oxford;New York,
- Lu H, Wu W, Chen Y, Wang H, Devare M, Thies JE (2010a) Soil microbial community responses to Bt transgenic rice residue decomposition in a paddy field. *J Soils Sed* 10 (8):1598-1605. doi:10.1007/s11368-010-0264-9
- Lu H, Wu W, Chen Y, Zhang X, Devare M, Thies JE (2010b) Decomposition of Bt transgenic rice residues and response of soil microbial community in rapeseed-rice cropping system. *Plant Soil* 336 (1-2):279-290. doi:10.1007/s11104-010-0476-1
- Lu YH, Abraham WR, Conrad R (2007) Spatial variation of active microbiota in the rice rhizosphere revealed by in situ stable isotope probing of phospholipid fatty acids. *Environ Microbiol* 9 (2):474-481
- Lukow T, Dunfield PF, Liesack W (2000) Use of the T-RFLP technique to assess spatial and temporal changes in the bacterial community structure within an agricultural soil planted with transgenic and non-transgenic potato plants. *FEMS Microbiol Ecol* 32 (3):241-247. doi:10.1111/j.1574-6941.2000.tb00717.x
- Lynch JM, Whipps JM (1990) Substrate flow in the rhizosphere. *Plant Soil* 129 (1):1-10
- Milling A, Smalla K, Maidl FX, Schlöter M, Munch JC (2004) Effects of transgenic potatoes with an altered starch composition on the diversity of soil and rhizosphere bacteria and fungi. *Plant Soil* 266 (1-2):23-39
- Naef A, Defago G (2006) Population structure of plant-pathogenic *Fusarium* species in overwintered stalk residues from Bt-transformed and non-transformed maize crops. *Eur J Plant Pathol* 116 (2):129-143
- Nielsen UN, Ayres E, Wall DH, Bardgett RD (2011) Soil biodiversity and carbon cycling: a review and synthesis of studies examining diversity–function relationships. *Eur J Soil Sci* 62 (1):105-116. doi:10.1111/j.1365-2389.2010.01314.x
- O'Callaghan M, Gerard EM, Bell NL, Waipara NW, Aalders LT, David BB, Conner AJ (2008) Microbial and nematode communities associated with potatoes genetically modified to express the antimicrobial peptide magainin and unmodified potato cultivars. *Soil Biol Biochem* 40:1446-1459. doi:10.1016/j.soilbio.2007.12.028
- Oehl F, Laczkó E, Bogenrieder A, Stahr K, Bösch R, van der Heijden M, Sieverding E (2010) Soil type and land use intensity determine the composition of arbuscular mycorrhizal fungal communities. *Soil Biol Biochem* 42 (5):724-738. doi:10.1016/j.soilbio.2010.01.006
- Oger P, Petit A, Dessaux Y (1997) Genetically engineered plants producing opines alter their biological environment. *Nat Biotechnol* 15 (4):369-372
- Oliveira AP, Pampulha ME, Bennett JP (2008) A two-year field study with transgenic *Bacillus thuringiensis* maize: Effects on soil microorganisms. *Sci Total Environ* 405 (1-3):351-357. doi:10.1016/j.scitotenv.2008.05.046

- Powell JR, Gulden RH, Hart MM, Campbell RG, Levy-Booth DJ, Dunfield KE, Pauls KP, Swanton CJ, Trevors JT, Klironomos JN (2007) Mycorrhizal and rhizobial colonization of genetically modified and conventional soybeans. *Appl Environ Microbiol* 73 (13):4365-4367
- Powell JR, Levy-Booth DJ, Robert HG, Wendy LA, Rachel GC, Kari ED, Allan SH, Miranda MH, Sylvain L, Robert EN, Pauls KP, Peter HS, Clarence JS, Jack TT, John NK (2009) Effects of genetically modified, herbicide-tolerant crops and their management on soil food web properties and crop litter decomposition. *J Appl Ecol* 46 (2):388-396
- Prosser JI (2012) Ecosystem processes and interactions in a morass of diversity. *FEMS Microbiol Ecol* 81 (3):507-519. doi:10.1111/j.1574-6941.2012.01435.x
- Raaijmakers J, Paulitz T, Steinberg C, Alabouvette C, Moënné-Loccoz Y (2009) The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant Soil* 321:341-361
- Radajewski S, Ineson P, Parekh NR, Murrell JC (2000) Stable-isotope probing as a tool in microbial ecology. *Nature* 403 (6770):646-649
- Rangel-Castro JI, Killham K, Ostle N, Nicol GW, Anderson IC, Scrimgeour CM, Ineson P, Meharg A, Prosser JI (2005) Stable isotope probing analysis of the influence of liming on root exudate utilization by soil microorganisms. *Environ Microbiol* 7 (6):828-838
- Rasche F, Lueders T, Schlöter M, Schaefer S, Buegger F, Gättinger A, Hood-Nowotny RC, Sessitsch A (2009) DNA-based stable isotope probing enables the identification of active bacterial endophytes in potatoes. *New Phytol* 181 (4):802-807. doi:10.1111/j.1469-8137.2008.02744.x
- Savka MA, Farrand SK (1997) Modification of rhizobacterial populations by engineering bacterium utilization of a novel plant-produced resource. *Nat Biotechnol* 15 (4):363-368
- Saxena D, Stotzky G (2001a) *Bacillus thuringiensis* (Bt) toxin released from root exudates and biomass of Bt corn has no apparent effect on earthworms, nematodes, protozoa, bacteria, and fungi in soil. *Soil Biol Biochem* 33 (9):1225-1230. doi:10.1016/s0038-0717(01)00027-x
- Saxena D, Stotzky G (2001b) Bt corn has a higher lignin content than non-Bt corn. *Am J Bot* 88 (9):1704-1706. doi:10.2307/3558416
- Seppänen SK, Pasonen HL, Vauramo S, Vahala J, Toikka M, Kilpeläinen I, Setälä H, Teeri TH, Timonen S, Pappinen A (2007) Decomposition of the leaf litter and mycorrhiza forming ability of silver birch with a genetically modified lignin biosynthesis pathway. *Appl Soil Ecol* 36 (2-3):100-106
- Setälä H, McLean MA (2004) Decomposition rate of organic substrates in relation to the species diversity of soil saprophytic fungi. *Oecologia* 139 (1):98-107
- Singh BK, Munro S, Potts JM, Millard P (2007) Influence of grass species and soil type on rhizosphere microbial community structure in grassland soils. *Appl Soil Ecol* 36 (2-3):147-155. doi:10.1016/j.apsoil.2007.01.004
- Tan F, Wang J, Chen Z, Feng Y, Chi G, Rehman SU (2011) Assessment of the arbuscular mycorrhizal fungal community in roots and rhizosphere soils of Bt corn and their non-Bt isolines. *Soil Biol Biochem* 43 (12):2473-2479. doi:10.1016/j.soilbio.2011.08.014
- Tan FX, Wang JW, Feng YJ, Chi GL, Kong HL, Qiu HF, Wei SL (2010) Bt corn plants and their straw have no apparent impact on soil microbial communities. *Plant Soil* 329 (1-2):349-364. doi:10.1007/s11104-009-0163-2
- Tilston EL, Halpin C, Hopkins DW (2013) Simultaneous down-regulation of enzymes in the phenylpropanoid pathway of plants has aggregated effects on rhizosphere microbial communities. *Biol Fertility Soils*:1-9. doi:10.1007/s00374-013-0862-y
- Turrini A, Sbrana C, Nuti MP, Pietrangeli BM, Giovannetti M (2004) Development of a model system to assess the impact of genetically modified corn and aubergine plants on arbuscular mycorrhizal fungi. *Plant Soil* 266 (1-2):69-75
- van der Heijden MGA, Klironomos JN, Ursic M, Moutoglou P, Streitwolf-Engel R, Boller T, Wiemken A, Sanders IR (1998) Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature* 396 (6706):69-72
- van der Heijden MGA, Bardgett RD, van Straalen NM (2008) The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol Lett* 11 (3):296-310. doi:10.1111/j.1461-0248.2007.01139.x
- van der Wal A, Geydan TD, Kuyper TW, de Boer W (2013) A thready affair: linking fungal diversity and community dynamics to terrestrial decomposition processes. *FEMS Microbiol Rev* 37 (4):477-494

- Verbruggen E, Kiers TE (2010) Evolutionary ecology of mycorrhizal functional diversity in agricultural systems. *Evolutionary Applications* 3 (5-6):547-560. doi:10.1111/j.1752-4571.2010.00145.x
- Verbruggen E, Rölting WFM, Gamper HA, Kowalchuk GA, Verhoef HA, van der Heijden MGA (2010) Positive effects of organic farming on below-ground mutualists: large-scale comparison of mycorrhizal fungal communities in agricultural soils. *New Phytol* 186 (4):968-979
- Verbruggen E, Kuramae EE, Hillekens R, de Hollander M, Kiers ET, Rölting WFM, Kowalchuk GA, van der Heijden MGA (2012) Testing potential effects of maize expressing the *Bacillus thuringiensis* Cry1Ab Endotoxin (Bt Maize) on mycorrhizal fungal communities via DNA- and RNA-based pyrosequencing and molecular fingerprinting. *Appl Environ Microbiol* 78 (20):7384-7392. doi:10.1128/aem.01372-12
- Wang GH, Xu YX, Jin J, Liu JD, Zhang QY, Liu XB (2009) Effect of soil type and soybean genotype on fungal community in soybean rhizosphere during reproductive growth stages. *Plant Soil* 317 (1-2):135-144. doi:10.1007/s11104-008-9794-y
- Wang Y, Xu J, Shen J, Luo Y, Scheu S, Ke X (2010) Tillage, residue burning and crop rotation alter soil fungal community and water-stable aggregation in arable fields. *Soil Till Res* 107 (2):71-79. doi:10.1016/j.still.2010.02.008
- Weaver MA, Krutz LJ, Zablotowicz RM, Reddy KN (2007) Effects of glyphosate on soil microbial communities and its mineralization in a Mississippi soil. *Pest Manage Sci* 63 (4):388-393. doi:10.1002/ps.1351
- Wei XD, Zou HL, Chu LM, Liao B, Ye CM, Lan CY (2006) Field released transgenic papaya affects microbial communities and enzyme activities in soil. *Plant Soil* 285 (1-2):347-358
- Weinert N, Meincke R, Gottwald C, Heuer H, Gomes NCM, Schlöter M, Berg G, Smalla K (2009) Rhizosphere communities of genetically modified zeaxanthin-accumulating potato plants and their parent cultivar differ less than those of different potato cultivars. *Appl Environ Microbiol* 75 (12):3859-3865. doi:10.1128/aem.00414-09
- Wolfenbarger LL, Phifer PR (2000) Biotechnology and ecology - The ecological risks and benefits of genetically engineered plants. *Science* 290 (5499):2088-2093
- Wrobel-Kwiatkowska M, Turnau K, Goralska K, Anielska T, Szopa J (2012) Effects of genetic modifications to flax (*Linum usitatissimum*) on arbuscular mycorrhiza and plant performance. *Mycorrhiza* 22 (7):493-499. doi:10.1007/s00572-011-0427-6
- Wu WX, Ye QF, Min H (2004) Effect of straws from Bt-transgenic rice on selected biological activities in water-flooded soil. *Eur J Soil Biol* 40 (1):15-22
- Wu WX, Liu W, Lu HH, Chen YX, Devare M, Thies J (2009) Use of C-13 labeling to assess carbon partitioning in transgenic and nontransgenic (parental) rice and their rhizosphere soil microbial communities. *FEMS Microbiol Ecol* 67 (1):93-102. doi:10.1111/j.1574-6941.2008.00599.x
- Xue K, Luo HF, Qi HY, Zhang HX (2005) Changes in soil microbial community structure associated with two types of genetically engineered plants analyzing by PLFA. *J Environ Sci (China)* 17 (1):130-134
- Xue K, Serohijos RC, Devare M, Thies JE (2011) Decomposition rates and residue-colonizing microbial communities of *Bacillus thuringiensis* insecticidal protein Cry3Bb-expressing (Bt) and non-Bt corn hybrids in the field. *Appl Environ Microbiol* 77 (3):839-846. doi:10.1128/aem.01954-10
- Zurbrugg C, Hoenemann L, Meissle M, Romeis J, Nentwig W (2010) Decomposition dynamics and structural plant components of genetically modified Bt maize leaves do not differ from leaves of conventional hybrids. *Transgenic Res* 19 (2):257-267. doi:10.1007/s11248-009-9304-x
- Zwahlen C, Hilbeck A, Nentwig W (2007) Field decomposition of transgenic Bt maize residue and the impact on non-target soil invertebrates. *Plant Soil* 300 (1-2):245-257

776
777**Table 1.** Studies on the effects of GM-crops on soil fungi, AMF and decomposer fungi. The primary crop species and modification studied, the method used, the set-up of the experiment and the outcome are listed in the columns.

Organisms	Research	Crop	Modification	Method	Environment studied	Outcome
Fungi	Donegan et al. 1996	Potato	Bt	CFU	Field study	Minor effect of GM trait in phyllosphere
	Donegan et al. 1999	Alfalfa	Lignin peroxidase and alpha amylase	CFU	Field study	No effect on fungi
	Saxena & Stoltzky 2001	Maize	Btk (Cry1Ab)	CFU	Greenhouse study	No effect on fungi
	Cowgill et al. 2002	Potato	Nematode resistance	FAME	Field study	Decrease in fungi in GM treatment
	Dunfield & Germida 2003	Canola	Glyphosate resistant	FAME	Field study	Transient effect
	Blackwood & Buyer 2004	Maize	Bt	PLFA	Greenhouse study	No effect on fungi
	Milling et al. 2004	Potato	granule bound starch synthase gene (gbs)	DGGE	Field study	No effect on fungi
	Xue et al. 2005	Maize and Potato	Bt and PVY (potato virus Y)	FAME	Field study	Decrease in fungi in GM treatment
	Gotz et al. 2006	Potato	T4 lysozyme	Microscopic analyses and DGGE	Field study	Differences in relative abundances between treatments
	Wei et al. 2006	Papaya	mutant gene of papaya rin spot virus	CFU	Greenhouse study	Increase in fungi in the soils grown with GM-papaya
	Weaver et al. 2007	Soybean	Glyphosate resistant (RoundUP)	FAME	Field study	No effect
	Girlanda et al. 2008	Tomato	Expression of tobacco b-1,3-glucanase and chitinase	CFU and AMF colonization	Greenhouse study	No effect on AMF or fungi
	Icoz et al. 2008	Corn	Bt	CFU	Field study	Transient effect
	Liu et al. 2008	Rice	Bt	TRFLP	Field study	No effect on fungi
	O'Callaghan et al. 2008	Potato	The antimicrobial peptide magainin II (against bacteria)	CFU	Field study	More fungi in roots of GM- potato
	Oliveira et al. 2008	Corn	Bt	CFU	Field study	Transient effect
	Hart et al. 2009	Corn	Glyphosate resistant (RoundUP)	qPCR - TRFLP	Field study	No effect on fungi
	Kremer & Means 2009	Soybean & corn	Glyphosate resistant	CFU of endophytes	Field study	Increase in <i>Fusaria</i> colonizing the GM-roots
	Weinert et al. 2009	Potato	carotenoid zeaxanthin accumulation	DGGE	Field study	No effect on fungi
	Wu et al. 2009	Rice	Bt	FAME	Greenhouse study	No effect on fungi
	Gschwendtner et al. 2010	Potato	granule bound starch synthase gene (gbs)	qPCR	Field and greenhouse studies	No effect on fungi
	Hannula et al. 2010	Potato	granule bound starch synthase gene (gbs)	TRFLP - enzymatic measurements, fungal biomass	Field study	No effect on AMF or fungi
	Tan et al. 2010	Corn	Bt	DGGE	Greenhouse study	No effect on fungi
	Gschwendtner et al. 2011	Potato	granule bound starch synthase gene (gbs)	FAME	Greenhouse study	No effect on fungi
	Lee et al. 2011	Rice	Fusion of trehalose-6-phosphate synthase	TRFLP and qPCR	Field study	No effect on fungi
	Li et al. 2011	Cotton	Bt	CFU- diversity	Field study	No effect on fungi
	Hannula et al. 2012a	Potato	granule bound starch synthase gene (gbs)	SIP- TRFLP	Greenhouse study	Transient effect on fungi and AMF
	Hannula et al. 2012b	Potato	granule bound starch synthase gene (gbs)	TRFLP, enzymatic measurements and fungal biomass	Field study	No effect on AMF or fungi
	Chun et al. 2012	Rice	Herbicide resistant	TRFLP	Field study	No effect on fungi
	Tilston et al. 2013	Tobacco	Phenylpropanoid metabolism	PLFA	Greenhouse study	Decrease in fungi in GM treatment
	Kuramae et al. 2013	Corn	Bt	454-Sequencing of DNA and RNA	Greenhouse study	No effect on fungi
AMF	Kaldorf et al. 2001	Aspen	Phytohormone balance	Colonization counts	Field study	No effect on AMF
	Turrini et al. 2004	Maize & Aubergine	Bt / antifungal	Colonization counts	Artificial greenhouse system (microcosm)	Transient effect
	Castaldini et al. 2005	Maize	Bt	Colonization counts	Artificial greenhouse system (microcosm)	Decreased colonization in one of the non-GM varieties
	de Vaulleury et al. 2007	Maize	Bt	Infectivity and colonization	Artificial greenhouse system (microcosm)	No effect on AMF
	Powell et al. 2007	Soybean	Herbicide resistant	Colonization counts	Greenhouse study	Transient effect
	Knox et al. 2008	Cotton	Bt and Glyphosate resistant	Colonization counts	Field study	No effect on AMF
	Hannula et al. 2010	Potato	granule bound starch synthase gene (gbs)	Colonization counts, TRFLP	Field study	No effect on AMF
	Cheeke et al. 2011	Maize	Bt	Colonization counts	Artificial greenhouse system (microcosm)	Transient effect
	Tan et al. 2011	Maize	Bt	Colonization counts and DGGE	Greenhouse study	Effect on the community structure but not on colonization
	Verbruggen et al. 2012	Maize	Bt	Microscopy, 454-seq RNA and DNA / T-RFLP	Greenhouse study	No effect
	Cheeke et al. 2012	Maize	Bt	Colonization counts	Greenhouse study	Decreased colonization in one of the GM varieties
	Wrobel-Kwiatkowska et al. 2012	Flax	Structural change	Colonization counts	Greenhouse study	No effect on AMF
	Hiebbach et al. 2012	Maize	Bt	Colonization counts	Greenhouse study	Transient effect
	Cheeke et al. 2013	Maize	Bt	Colonization counts and spore diversity estimate	Field study	No effect on AMF
Decomposer Fungi	Donegan et al. 1995	Cotton	Bt	CFU	Greenhouse study	Transient effect
	Wu et al. 2004	Rice	Bt	CFU	Greenhouse study	Transient effect
	Castaldini et al. 2005	Maize	Bt	Colonization counts	Greenhouse study	Transient effect
	Flores et al. 2005	Maize, canola, potatoes, cotton, rice, tobacco	Bt	CFU	Greenhouse study	No effect
	Henault et al. 2006	Tobacco	Lignin-synthesis	FAMES	Greenhouse study	Twice as much fungal marker in the GM soil
	Naef & DeFago 2006	Corn	Bt	Multiplex PCR - micro assay	Field study	Transient effect
	Seppinen et al. 2007	Birch	Lignin-synthesis	Ergosterol	Field study	No effect
	Lawhorn et al. 2009	Corn	Bt	Extracellular enzymes	Field study	No effect
	Powell et al. 2009	Soybean - maize	Glyphosate-tolerant	DFS and hyphal length	Greenhouse study	No effect
	Lu et al. 2010a	Rice	Bt	TRFLP	Field study	Transient effect
	Lu et al. 2010b	Rice	Bt	TRFLP	Field study	No effect
	Tan et al. 2010	Corn	Bt	DGGE	Greenhouse study	Transient effect
	Xue et al. 2011	Corn	Bt	TRFLP	Field study	Transient effect
	Hannula et al. 2013	Potato	granule bound starch synthase gene (gbs)	TRFLP, qPCR	Greenhouse study	Transient effect

778
779