Herbicides in the environment alter infection dynamics in a microbial host-parasite system

Silke Van den Wyngaert ^{1,2*}, Alena S. Gsell ³, Piet Spaak ^{1,2} and Bas W. Ibelings ^{1,2,3,4}

¹Aquatic Ecology, Eawag, Ueberlandstrasse 133, P.O. Box 611, 8600 Duebendorf, Switzerland

²Institute of Integrative Biology, ETH Zürich, CH-8092 Zürich, Switzerland

³Netherlands Institute of Ecology, Department of Aquatic Ecology, Droevendaalsesteeg 10, 6708 PB Wageningen, The Netherlands

⁴Institut F.-A. Forel, Université de Genève, 10 Route de Suisse, Versoix, Switzerland

***Present Address & Corresponding author:**

Silke Van den Wyngaert

Laboratory of Aquatic Biology, Department of Biology, K.U.Leuven Kulak

E. Sabbelaan 53, 8500 Kortrijk, Belgium

Telephone: +32 56 24 60 59

Fax: +32 56 24 69 99

silke.vandenwyngaert@kuleuven-kulak.be

Running title: Infection dynamics and environmental pollution

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/j.1462-2920.2012.02874.x

23 *Summary*

24 Parasites play an important role in the regulation of host population growth. How these
25 ubiquitous stressors interact with anthropogenic stressors is less often studied. In a full
26 factorial experiment we explored the independent and combined effects of the widely used
27 herbicide diuron and a chytrid parasite on the fitness of genetically different monoclonal
28 diatom populations. Furthermore we evaluated how herbicide exposure influenced infection
29 dynamics, parasite fitness and the impact of infectious disease on host populations. We found
30 no evidence of host genetic variation for diuron sensitivity and parasite resistance. Instead,
31 host population phenotype was a decisive factor in controlling parasite growth. Although
32 herbicide exposure initially posed a constraint on disease transmission, it enhanced the spread
33 of disease over time. Consequently the nature of the parasite-toxicant stressor interaction
34 shifted from antagonistic (on exponential host growth) towards additive (on final uninfected
35 host density). We conclude that herbicide exposure can modify infection dynamics and
36 impact of disease on host populations through the complex interplay between host and
37 parasite growth dynamics and host population phenotype.

38 *Introduction*

39 In nature, organisms are exposed simultaneously to biotic and abiotic stressors. Depending on
40 the stressor combination, joint effects can be additive, synergistic or antagonistic. Although
41 an increasing number of studies consider the impact and interaction of multiple stressors, it
42 remains difficult to make general predictions about the outcome of their combined effects in
43 nature. This lack of knowledge has become a major concern in ecological risk assessment
44 (Crain et al. 2008; Holmstrup et al. 2010)

45 Parasitism represents an important biotic stress factor that influences host populations
46 in the field. Considering the high abundance and diversity of parasites we can fairly say that
47 every species is affected by parasites (Windsor 1998). They can play a significant role in
48 shaping population dynamics by altering the genetic structure and diversity of their host
49 populations (Decaestecker et al. 2007; Duncan and Little 2007; Jokela et al. 2003). Effects of
50 parasitism and anthropogenic environmental stressors such as chemical pollutants interact in
51 intricate ways. Host organisms exposed to chemical stressors may lack the energy for
52 mounting an efficient defence against parasite attack and therefore become more susceptible
53 to infection (Holmes 1996). Studies on vertebrate host parasite model systems provide
54 evidence for this line of reasoning: e.g. an amphibian-trematode system (Koprivnikar et al.
55 2007; Rohr et al. 2008a; Rohr et al. 2008b) or fish-pathogen systems (Kreutz et al. 2010).
56 Studies on the Eastern Oyster, an aquatic invertebrate, also reported increased vulnerability to
57 a protozoan pathogen when hosts were pre-exposed to pollutants (Chu and Hale 1994; Fisher
58 et al. 1999). Similarly, negative effects of pesticide exposure on disease resistance in the
59 invertebrate water flea *Daphnia magna* were found (Coors and De Meester 2008; Coors et al.
60 2008). Also for higher plant parasite systems there is evidence that pesticides can lower plant
61 defence causing increased disease severity (Johal and Huber 2009). However, interactions
62 between stress and disease are much more complex (Lafferty and Holt 2003). Ecosystem

63 stressors may reduce host density and/or host quality and therefore have negative effects on
64 parasite transmission and/or parasite reproduction (Bittner et al. 2002; Seppälä et al. 2008).
65 Furthermore, host genotypes may harbour large genetic variation for traits involved in
66 infection and pollution resistance. Whether these traits are positively or negatively correlated
67 will affect selection processes and consequently the impact of multiple stressors on host
68 population dynamics.

69 Phytoplankton, i.e. cyanobacteria and eukaryotic microalgae, as primary producers provide
70 many important ecosystem services, form the basis of entire aquatic food webs and play a key
71 role in the global carbon cycle (e.g. Falkowski et al. (2000)). Therefore, understanding
72 phytoplankton responses to combined natural and anthropogenic stressors is important
73 because of knock-on effects at higher trophic levels and whole ecosystem functioning.
74 Although still largely understudied, many phytoplankton species have obligate, host specific
75 parasites that have the potential to alter phytoplankton population dynamics and succession
76 (Canter 1950; Canter and Lund 1948; Ibelings et al. 2011). In addition, phytoplankton species
77 are relevant non-target organisms for a multitude of pesticides, especially for herbicides,
78 which are applied in agricultural weed-control and reach stream- and lake ecosystems via run-
79 off from the land (Cedergreen and Streibig 2005). Peak concentrations of herbicides are
80 commonly found in spring after field application and rainfall (Rabiet et al. 2010). This is also
81 the time when phytoplankton spring blooms develop and chytrid epidemics occur (Ibelings et
82 al. 2004). We therefore believe that this stressor interaction may significantly affect
83 phytoplankton population dynamics and disease in the field. To our knowledge, no study so
84 far has explored the interaction between chemical pollution and disease in phytoplankton.

85 In this study we address three main questions using a well studied microbial host-parasite
86 system, consisting of the diatom *Asterionella formosa* Hassall host, and the chytrid fungus
87 *Zygorhizidium planctonicum* Canter as parasite, exposed to the herbicide diuron: 1) Do host

88 genotypes show variation for parasite resistance and herbicide resistance traits? If so, are
89 these two traits correlated? 2) How does herbicide exposure interact with parasite stress: is the
90 nature of the interaction additive, synergistic or antagonistic? 3) Does herbicide exposure alter
91 infection dynamics and impact of parasitism on the algal host population? We demonstrate
92 experimentally that environmental pollution has the potential to alter infection dynamics and
93 the severity of disease in this diatom-chytrid host-parasite system.

94

95 *Results*

96 **Single and combined effects of diuron and parasite on host fitness**

97 For average host population growth rate the AIC-selected model discarded all random
98 interaction terms but included the interaction terms between the fixed factors diuron x parasite
99 and between both fixed factors and the covariate host population cell volume. Variation
100 explained by the random factor host genotype compared to the residual variance was very low
101 (almost 0%). The interaction between mean host population cell volume and stressors was
102 significant for the parasite treatment and marginally significant for the diuron treatment
103 (Table 1). The estimates of the interaction terms were low, however, and the interaction was
104 mainly driven by the host population with the smallest mean cell volume. The combined
105 stress treatment decreased host fitness compared to the control treatment but the decrease was
106 less (with the exception of Strain 26) than expected by the reference model of independent
107 action (IA) (Fig. 1a). This indicates an antagonistic interaction which was substantiated by a
108 significant two way interaction between diuron and parasite (Table 1).

109 For final uninfected host density, the AIC-selected model also discarded all random
110 interaction terms but included the three way interaction term between the fixed factors diuron
111 and parasite and the covariate mean host cell volume (Table 2). The significant three way
112 interaction indicates that the interaction between diuron and parasite depends on host

population cell size (Table 2). The interaction between diuron and parasite was therefore analyzed separately for each of the five host populations (corresponding to each level of the covariate host population cell volume). The results of this analysis show that the combined stress treatment also decreased the final uninfected host density in each population separately. However, for the host population with the smallest cell size (strain 37) this decrease was less than expected by IA (antagonistic interaction; $F = 11.83$, $P = 0.0063$, Fig 1b). The interaction became additive for populations with larger cell size. This was indicated by non-significant interactions between diuron and parasite and by the expected IA value lying within the confidence interval of the observed uninfected host density (Fig.1b). The host fitness reduction by the two stressors followed an opposite trend with host populations of smaller mean cell volume being more sensitive to diuron, but suffering less from parasite stress, and vice versa for populations with larger mean cell volume (Figs. 2a&b). This trade off pattern, mediated by host cell volume, was expressed strongest in the response variable percentage reduction in final uninfected host density (Fig. 2b).

Parasite fitness

Parasite population growth rate increased significantly in all host populations exposed to diuron (Table 3, Fig. 3) and was positively related to mean host population cell volume, independent of herbicide treatment (no diuron; linear regression, $r^2 = 0.245$, $P = 0.012$, $n = 25$, diuron; $r^2 = 0.276$, $P = 0.007$, $n = 25$, Fig. 3). Tukey HSD posthoc analysis revealed a significant difference between parasite growth rate infecting host populations with the smallest and largest mean cell volume ($P < 0.011$).

Impact of diuron on parasite infection dynamics

The infection dynamics differed significantly between the two treatments (time $\times$ treatment interaction; $F = 10.49$, $P < 0.001$). Since the three way interaction between the covariate (mean host population cell volume), time and treatment was not significant ($F = 0.58$, $P = 0.67$) the data from all host populations were pooled together in order to better visualize the infection dynamics of both treatments. Infection started immediately in the non diuron treatment whereas an infection lag phase occurred in the presence of diuron. After day 4 the slope of infection increase was steeper with, than without diuron (Fig. 4a). Multiple infections were rare so that the increase in number of infections was equivalent to the increase in number of sporangia. Therefore, parasite fitness (based upon the exponential increase of number of sporangia) was also higher in the presence of diuron (see results above). Total host density in the parasite exposed treatment increased over time but remained overall lower in the presence of diuron (Fig. 4b). This resulted in a final infection prevalence that was almost twice as high in the diuron treatment as compared to the non diuron treatment (Fig. 4c).

Discussion

In this study we investigated how abiotic and biotic stressors interact, resulting in environment by environment ($E \times E$) interactions. In particular we studied the interaction between a herbicide that inhibits algal photosynthesis and a virulent chytrid parasite in their effects on a common planktonic diatom. In addition we included a third interaction term “host genotype” (G) in order to identify whether genotypic heterogeneity in response to single and multiple stressors does occur (i.e. host $G \times E$ and $G \times E \times E$).

Host population phenotype and response to stress

Contrary to many other studies we found no evidence for genetic variation in resistance traits and host $G \times E$ interactions (Blanford et al. 2003; Mitchell et al. 2005). It was previously shown that different *A. formosa* genotypes differed in their susceptibility to two different *Z. planktonicum* strains and that parasite strains differed in their ability to infect particular host genotypes (De Bruin et al. 2004). However, from the 17 different host genotypes tested in that paper the majority (9 genotypes) were severely infected by both parasite strains. Therefore the number of host genotypes used in this experiment - which were genetically differentiated based on AFLP marker analysis - was presumably too low to detect differences in susceptibility to parasitism. The study of De Bruin et al. (2004), however, did not consider effects of host population phenotype. Our study shows that host population phenotype, i.e. mean cell volume, plays a role in the response to single and multiple stressors and is positively correlated with parasite growth. In diatoms, asexual reproduction results in a progressive reduction in cell size. Differences in cell size between genotypes are most likely explained by genotypes being at different stages of cell-size reduction. The first question of our study (i.e. do host genotypes show variation for parasite resistance and herbicide resistance traits?) can therefore not be answered since we are unable to disentangle phenotype from genotype.

The host population phenotype $\times$ environment interactions suggest that populations with smaller mean cell size are more sensitive to the herbicide diuron but suffer less from parasite stress, while the opposite is true for populations with larger mean cell size (Table 1, Fig. 2). We are aware that slope estimates for the interaction terms were low and the linear relationship between fitness reduction and cell volume was only significant for percent inhibition of final uninfected host density with diuron. Our experiment was initially not designed to investigate phenotypic responses and only the host population with smallest mean

cell volume significantly differed from all other host populations. Therefore we lack power to properly estimate the effect of host cell volume on stress response, but a trend was visible. Several studies found a negative relationship between cell volume and toxicant sensitivity in phytoplankton (Lockert et al. 2006; Tang et al. 1998). A similar relationship for invertebrate crustaceans was found by (Vesela and Vijverberg 2007) with smaller sized daphnids being more sensitive to heavy metal toxicity. In contrast, a positive relationship between body size and infection has been demonstrated in several other host-parasite systems with free-living parasite transmission stages: e.g. *Daphnia*-yeast system (Duffy et al. 2011; Hall et al. 2007) or a snail-trematode system (Seppälä et al. 2011). The proposed mechanisms that lead to this relationship are the higher feeding rate of larger hosts which subsequently lead to higher parasite clearance rates and the increased production of parasite transmission stages on larger hosts (Duffy et al. 2011). Since *A. formosa* is not a filter feeding organism, the higher feeding rate mechanism can obviously be excluded. However, increased parasite production on larger host strains would be a possible mechanism. The correlation between host population cell size and parasite population growth which we observed, could well be explained by increased parasite reproduction - i.e. increased zoospore production - on larger host cells as more nutrients can be extracted and exploited by the parasite. Moreover, a positive correlation between parasite sporangium size and host cell size has also been observed in another diatom-chytrid association (Holfeld 2000). This ecological dependence of parasitism on host phenotype likely rivals and/or interacts with genetic mechanisms that often get more attention in the literature (Hall et al. 2009).

Interaction between stressors

Our second objective was to identify the direction of the interaction (synergistic versus antagonistic) between herbicide and parasite exposure. There is increasing consensus that

parasitism acting concertedly with other stressors has stronger detrimental effects on host organisms than either stressor acting alone presumably due to reduced host immune defences (Marcogliese et al. 2009).

Our results show that the direction of stressor interactions depends on the focal host fitness parameter (reflecting different stages of host population growth – see paragraph on infection dynamics) and on the host population phenotype. The reduction in host growth rate in the combined stress treatment was less than the sum of both single stressors and resulted in an antagonistic interaction between parasite and diuron. However, when the final uninfected host cell density is considered, a three way interaction between mean population cell size, diuron and parasite indicates that the interaction between the two stressors is dependent on host population phenotype, ranging from antagonistic for the host population with smallest mean cell size to additive for populations with larger mean cell size. However, this effect was mainly driven by the smallest host population where the single effect of diuron was very strong and the negative effect of parasite remained weak.

Host defence strategies against parasites range from avoidance or structural resistance to immune responses associated with excretion of defence molecules (Rigby et al. 2002) or recognition of pathogen associated molecular patterns (Chisholm et al. 2006) leading to programmed cell death (Lam et al. 2001). For some clonal strains of *A. formosa* a hypersensitive death reaction to infection by *Rhizophyidium planktonicum* has been observed resulting in growth arrestment and finally death of the parasite (Canter and Jaworski 1979). However, the exact mechanisms that lead to successful inhibition of infection in the *A. formosa* - chytrid interaction are not known and we did not observe any signs of hypersensitive reaction during our experiment. Our results on the combined effect of parasite and diuron may be not so much linked to host immune defence but more to the complex

interplay between host population phenotype and host and parasite population growth dynamics.

Infection dynamics

Infection dynamics differed in the presence or absence of the environmental stressor diuron (Fig. 4a). Diuron exposure resulted in higher parasite growth (Fig. 3) and a reduction in uninfected host density (Fig 1b). However, initially, the infection started off slower when the herbicide was present, which explains the antagonistic interaction between diuron and parasite on exponential host population growth rate.

The reduction of host growth rate and thereby host cell density through diuron exposure could directly constrain parasite transmission efficiency. However, the initial host density in our experiment was relatively high compared to threshold host densities allowing epidemics in the field (Ibelings et al. 2011). Alternatively, low initial infection success may be explained by the fact that many parasites with free living stages use chemical cues to track their host (Gerardo et al. 2006; Kuhn 1997). Previous experiments have shown that zoospores are unable to infect *A. formosa* in darkness or at very low irradiance (Bruning 1991b; Canter and Jaworski 1981). The release of extracellular organic carbon from living algal cells increases with photosynthetic production and generally correlates with increased irradiance (Espeland and Wetzel 2001). Therefore it is supposed that zoospores are attracted by specific extracellular products from *A. formosa* that are excreted during active photosynthesis (Bruning 1991b). The concentration of diuron used in this experiment provokes ca. 50% photosynthesis inhibition (see supporting information, Fig. S1A), so that diuron probably interfered with zoospore behaviour and negatively affected host finding mechanisms. Nevertheless infection in the diuron treatment caught up with infections in the controls, and from day four onwards parasite infections even increased more rapidly in presence than in

262 absence of diuron. How to explain this? Despite the inhibiting effect of diuron on
263 photosynthesis host cell density continuously increased during the experiment, also in
264 presence of the herbicide. Higher host densities may have increased chance contacts between
265 host cells and zoospores so that the herbicide imposed constraints on chemotaxis are less
266 relevant for infections in the late stages of the experiment, when host cell density had
267 sufficiently increased.

268 Although parasite transmission was initially negatively affected by diuron, the overall
269 parasite growth rate increased in the presence of the herbicide. As already mentioned above,
270 the effect of increasing host density may have annulled the possible transmission constraints
271 imposed by lower initial host density. However, this does not explain the steeper infection
272 increase with diuron after the second parasite generation. The effect of diuron on parasite
273 growth can be direct or indirect. For example, some studies have shown a direct growth
274 stimulating effect of herbicides on pathogenic soil fungi (Altman and Campbell 1977; Davis
275 et al. 1976) and the potential of several asco-, basidio-, and zygomycetes to metabolize
276 phenylurea herbicides (Gondim-Tomaz et al. 2005; Ronhede et al. 2005). On the other hand,
277 continuous exposure to herbicides could increase host susceptibility. Diuron affects
278 photosynthetic electron transport and thereby causes the inhibition of sugar production in the
279 Calvin cycle through the depletion of ATP and NADPH supplied by the light reaction.
280 Inhibition of photosystem II electron transport also generates reactive oxygen species (ROS)
281 that have the potential to cause membrane protein damage (Cobb and Reade 2010). Longer
282 exposure to the herbicide may therefore disrupt physiological processes which could increase
283 host susceptibility to parasites in the longer term. The possible alteration of host physiology
284 through diuron exposure may also have affected parasite development time. Phosphorous
285 limitation of *A. formosa* for example was found to reduce sporangia development time
286 (Bruning and Ringelberg 1987). However, it also reduced the number of zoospores per

sporangium. Hence the overall effect of phosphorous limitation on parasite growth was negative (Bruning and Ringelberg 1987). Faster sporangia development time could be an underlying mechanism for higher infection rates if zoospore production is not reduced. Finally parasite growth rate integrates different life history traits, each of which may interact differently with changes in the batch culture environment as the experiment progresses (Bruning 1991a; b; Vale and Little 2009). It is therefore difficult to pinpoint the underlying mechanisms of the observed differences in infection dynamics and the outcome between the treatments (Pulkkinen and Ebert 2004).

Does stress increase the impact of disease on host populations?

Lafferty and Holt (2003) differentiate between the spread of disease (R_0) which describes the parasite population growth and the impact of disease on host population growth. In our system, the spread of disease was initially negatively related to the stressor diuron. This resulted in an antagonistic interaction between diuron and parasite with a weaker negative effect of the parasite on the exponential host population growth rate in the presence of diuron. The presence of this herbicide did therefore not implicate an increased impact of disease on exponential host population growth. However, this interaction between stress and spread of disease shifted from antagonistic (early infection phase) towards additive which is reflected by the increased infection rate after day four in presence of diuron. Together with this shift towards an increased R_0 , also the impact of disease on host populations increased (lower density of uninfected host cells, Fig.4b, and higher prevalence of infection, Fig.4c). Although we do not have a mechanistic explanation yet for this shift this would indicate a risk of more severe epidemics when host populations are exposed to the photosynthesis inhibiting herbicide diuron. However, care must be taken with this general statement since our results suggest that mechanisms of density dependent efficiency of parasite transmission are in

operation. Therefore the outcome of our laboratory infection experiment is probably highly dependent on initial host and parasite densities (Ben-Ami et al. 2008). Moreover, the initial host density in our experiment was relatively high compared to natural host densities in the field (Ibelings et al. 2011). Therefore, the negative effect of diuron exposure on parasite transmission could be even more pronounced under natural conditions, resulting in the reduction of epidemics. Furthermore, when our hypothesis is true that reduced photosynthesis of the algal hosts negatively affects parasite transmission (through chemotaxis interference), one could generalize that any herbicide (or any other stress factor) affecting photosynthesis will result in lower chytrid infection rates. This hypothesis requires further investigation. The strength and impact of the interaction between parasite and herbicide probably also depends on the concentration and duration of herbicide exposure. Our experimental setup was limited to one concentration of diuron and one exposure scenario (continuous exposure) causing continuous inhibition of photosynthesis (Fig. S1). However, exposure to herbicides in the aquatic environment often occurs as repeated pulses with possible recovery of photosynthesis and algal growth between exposures (Vallotton et al. 2008). Further studies are needed to elucidate how such a pulsed exposure scenario would affect chytrid infection.

Conclusion

We present the first experimental study on the impact of disease in phytoplankton populations exposed to an anthropogenic chemical stressor. Simultaneous exposure of host populations to a parasite and a toxicant allowed the observation of how initial disease transmission and later disease development are affected at the population level. An important finding of this study is that environmental stress modifies the infection dynamics, thereby changing the nature of the stressor interaction from antagonistic towards additive. We therefore gained valuable insights into the dynamic nature of interactions between environmental pollution and disease. Further,

our results showed that host population phenotype can be a decisive factor in controlling parasite growth. We conclude that anthropogenic stress can modify infection dynamics and the impact of disease on host populations through the complex interplay between stressor, host and parasite growth dynamics and host population phenotype.

Experimental Procedures

Host-Parasite study system

The host *Asterionella formosa* is a cosmopolitan freshwater diatom which forms stellate colonies. Reproduction in this species is predominantly asexual and characterized by a “shrinking division” mode which is specific for diatoms. This process leads to a progressive reduction in cell size. So far, no sexual reproduction has been observed for *A. formosa*. However, rare sexual events are assumed since these may explain the high genetic diversity in a Dutch lake population (De Bruin et al. 2008) and regular size rejuvenation in the Lake Zurich population (Nipkow 1927).

The parasite *Zygorhizidium planktonicum* is an aquatic fungus belonging to the Chytridiomycetes (James et al. 2006). It is an obligate, host-specific parasite of *A. formosa* (Van Donk and Ringelberg 1983). Hence, the parasite cannot survive and/or reproduce in absence of its host. *Z. planktonicum* is an extremely virulent parasite i.e. every infection inhibits host reproduction and leads to host cell death. The life cycle of this chytrid parasite begins with the attachment of a motile zoospore to the surface of a host cell. After zoospore encystment a germ tube is formed which enters the host cell through the girdle zone. Via the germ tube nutrients are extracted from the host cell and used for the development of the sporangium. New zoospores are formed either asexually or sexually and are released from the sporangium by dehiscence (Doggett and Porter 1996; Ibelings et al. 2004).

362 Experimental design

363 All host and parasite strains were isolated during the spring bloom of 2008 from Lake
364 Maarsseveen, (52.142828 N, 5.085711 E, The Netherlands). We conducted a full factorial
365 experiment with five different host genotypes (based upon AFLP fingerprinting, Gsell et al.
366 *in press*), one parasite genotype and four treatments: 1) stress-free (positive control), 2)
367 diuron, 3) parasite, 4) diuron $\times$ parasite. All treatment combinations were tested in batch
368 cultures, incubated in water baths at 18 °C and 100 $\mu\text{E m}^{-2} \text{s}^{-1}$ irradiance. The total volume of
369 a batch was 65 mL (in 100 mL Erlenmeyer flasks) and each treatment was replicated five
370 times resulting in 100 experimental units. Each flask was manually shaken once per day. The
371 experiment was started with an initial host density of 10 000 cells mL^{-1} . Approximately 0.5
372 mL of an old infected *A. formosa* culture (100% prevalence of infection) was added to obtain
373 an initial parasite sporangia concentration of 500 sporangia mL^{-1} corresponding to a starting
374 prevalence of infection of 5%. Based on the results of a previous dose response experiment
375 with diuron (DCMU (3-(3,4-dichlorophenyl)-1,1-dimethylurea; CAS 330 54 1, PESTANAL®
376 analytical standard, Sigma-Aldrich) and three different *A. formosa* strains from Lake
377 Maarsseveen, a sublethal concentration of 8 $\mu\text{g L}^{-1}$ diuron was added. This concentration
378 results in ca. 25% growth inhibition and 50% photosynthesis inhibition (see supporting
379 information). Diuron, is a herbicide that inhibits photosynthesis by binding to the D-1 protein
380 of the Photosystem 2 reaction centre, thereby blocking photosynthetic electron transport
381 (Duke 1990). It is one of the most frequently and permanently observed agrochemicals in
382 surface waters (Blanchoud et al. 2004; Chèvre et al. 2006) and is highly persistent in the
383 environment (Giacomazzi and Cochet 2004). Peak concentrations of this herbicide of up to 28
384 $\mu\text{g L}^{-1}$ are found in the environment (Field et al. 2003), therefore a concentration of 8 $\mu\text{g L}^{-1}$
385 can be considered ecologically plausible and relevant. The experiment was conducted for 8
386 days (if experiments last longer (multiple) infections gets so widespread that infection is

impossible to quantify accurately) and every two days 3 mL subsamples were fixed with a mixture of paraformaldehyde (0.01% final concentration) and glutaraldehyde (0.1% final concentration). Before the experiment was started, approximately 60 cells (from 60 different colonies) of each host population were measured and their cell biovolume was calculated assuming the shape of a rectangular box ($V = \text{length} \times (\text{width})^2$). We counted total abundance of host cells (N), abundance of infected (N_i) and uninfected (N_u) host cells, and abundance of fungal sporangia and attached zoospores using an inverted fluorescence microscope (Fluovolt FS, Leitz) according to the Uthermoehl method (see Van Donk and Ringelberg 1983). Average host population growth rate and parasite population growth rate were calculated from the exponential increase of total host cells mL^{-1} and sporangia mL^{-1} respectively (Guillard 1973): $K' = \ln(N_2 / N_1) / (t_2 - t_1)$.

Where: N_1 and N_2 is abundance at t_1 and t_2 respectively.

A second host fitness parameter was determined; which we termed “final uninfected host density”. It was calculated as the abundance of live, uninfected host cells at the last day of the experiment minus the initial host cell abundance. To assess host genotype sensitivity for diuron and parasites the percentage of host growth rate and final host density inhibition compared to the control treatment was calculated.

Statistical analysis

Statistical analyses were performed using the free software R, version 2.8.1 (R Development Core Team 2008). The effects of the single and combined stress treatments on two fitness parameters of the five *A. formosa* genotypes were analyzed by a linear mixed effect model (lmer function of the lme4 package) with average growth rate and final uninfected host density as dependent variables, diuron exposure (yes/no) and parasite exposure (yes/no) as fixed factors and host genotype as random factor. To account for cell size differences between

the host genotype populations mean host cell volume was included as covariate in the model. Models were fitted using the maximum likelihood method in R and model selection was conducted by standard Akaike information criterion (AIC) methods, i.e. starting from the model including all higher order interactions and progressively deleting the least significant interaction term. In a similar analysis the effect of diuron on parasite fitness was tested with parasite population growth rate as dependent variable, diuron exposure (yes/no) as fixed factor, host mean cell biovolume as covariate and host genotype as random factor. The significance of the fixed factors in the linear mixed effects models was estimated by Markov Chain Monte Carlo (MCMC) simulations with the function pvals.fnc from language R package (Baayen et al. 2008). To analyze the infection dynamics a repeated measure analysis was performed using the lme function of the nlme package. Sampling day was the repeated factor and number of infections the dependent variable. Diuron exposure (yes/no) was fixed and the mean host population cell volume was included as covariate. The best fitting covariance structure was determined by comparing AIC values. To meet the assumptions of normality (tested with q-q plots) and homogeneity of variance (Bartlett test) the data for number of infections was log transformed.

Prediction of joint effects

We assume that parasitism and the photosynthesis inhibiting herbicide diuron act independently from each other. Therefore the Bliss model of independent action (Bliss 1939) was used to calculate the reference value for additive effects of combined stressors, which is obtained by taking the sum of each single stressor minus their product (Andersen et al. 2009). Deviations from this predicted additive effect are thus an indication of antagonistic (less than predicted) or synergistic (greater than predicted) effects.

437 *Acknowledgements*

438 We thank Christoph Tellenbach for statistical advice and helpful comments on the
439 manuscript. Further we thank Ellen van Donk for providing the lab facilities to conduct the
440 experiment. This research was funded by the ETH Board (CCES GEDIHAP).

442 *References*

- 443 Altman J, Campbell CL (1977) Effect of herbicides on plant diseases. *Annu. Rev.*
444 *Phytopathol.* 15:361-385
- 445 Andersen IS, Voie OA, Fonnum F, Mariussen E (2009) Effects of Methyl Mercury in
446 Combination with Polychlorinated Biphenyls and Brominated Flame Retardants on
447 the Uptake of Glutamate in Rat Brain Synaptosomes: A Mathematical Approach for
448 the Study of Mixtures. *Toxicol. Sci.* 112:175-184
- 449 Baayen RH, Davidson DJ, Bates DM (2008) Mixed-effects modeling with crossed random
450 effects for subjects and items. *J. Mem. Lang.* 59:390-412
- 451 Ben-Ami F, Regoes RR, Ebert D (2008) A quantitative test of the relationship between
452 parasite dose and infection probability across different host-parasite combinations.
453 *Proc.R. Soc. B* 275:853-859
- 454 Bittner K, Rothhaupt K-O, Ebert D (2002) Ecological interactions of the microparasite
455 *Caullerya mesnili* and its host *Daphnia galeata*. *Limnol. Oceanogr.* 47:300-305
- 456 Blanchoud H, Farrugia F, Mouchel JM (2004) Pesticide uses and transfers in urbanised
457 catchments. *Chemosphere* 55:905-913
- 458 Blanford S, Thomas MB, Pugh C, Pell JK (2003) Temperature checks the Red Queen?
459 Resistance and virulence in a fluctuating environment. *Ecol. Lett.* 6:2-5
- 460 Bliss CI (1939) The toxicity of poisons applied jointly. *Ann. Appl. Biol.* 26:585-615

- 461 Bruning K (1991a) Effects of phosphorus limitation on the epidemiology of a chytrid
462 phytoplankton parasite. *Freshw. Biol.* 25:409-417
- 463 Bruning K (1991b) Infection of the diatom *Asterionella* by a chytrid.1. Effects of light on
464 reproduction and infectivity of the parasite. *J. Plankton Res.* 13:103-117
- 465 Bruning K, Ringelberg J (1987) The Influence of Phosphorus Limitation of the Diatom
466 *Asterionella-Formosa* on the Zoospore Production of Its Fungal Parasite
467 *Rhizophydium-Planktonicum*. *Hydrobiol. Bull.* 21:49-54
- 468 Canter HM (1950) Fungal parasites of the phytoplankton. 1. Studies on British chytrids X.
469 *Ann. Bot.* 14:263-&
- 470 Canter HM, Jaworski GHM (1979) Occurrence of a hypersensitive reaction in the planktonic
471 diatom *Asterionella formosa* Hassall parasitized by the chytrid *Rhizophydium*
472 *planktonicum* Canter Emend., in culture. *New Phytol.* 82:187-&
- 473 Canter HM, Jaworski GHM (1981) The effect of light and darkness upon infection of
474 *Asterionella-formosa* Hassall by the chytrid *Rhizophydium-planktonicum* Canter
475 Emend. *Ann. Bot.* 47:13-30
- 476 Canter HM, Lund JWG (1948) Studies on plankton parasites. 1. Fluctuations in the numbers
477 of *Asterionella-formosa* Hassall in relation to fungal epidemics. *New Phytol.* 47:238-
478 261
- 479 Cedergreen N, Streibig JC (2005) The toxicity of herbicides to non-target aquatic plants and
480 algae: assessment of predictive factors and hazard. *Pest Manag. Sci.* 61:1152-1160
- 481 Chèvre N, Loeppé C, Singer H, Stamm C, Fenner K, Escher BI (2006) Including mixtures in
482 the determination of water quality criteria for herbicides in surface water. *Environ.*
483 *Sci. Technol.* 40:426-435
- 484 Chisholm ST, Coaker G, Day B, Staskawicz BJ (2006) Host-microbe interactions: Shaping
485 the evolution of the plant immune response. *Cell* 124:803-814

- 486 Chu FLE, Hale RC (1994) Relationship between pollution and susceptibility to infectious
487 disease in the eastern oyster, *Crassostrea virginica*. Mar. Environ. Res. 38:243-256
- 488 Cobb AH, Reade JPH (2010) Herbicides That Inhibit Photosynthesis, in Herbicides and Plant
489 Physiology, Second edition, Wiley-Blackwell, Oxford, UK
- 490 Coors A, De Meester L (2008) Synergistic, antagonistic and additive effects of multiple
491 stressors: predation threat, parasitism and pesticide exposure in *Daphnia magna*. J.
492 Appl. Ecol. 45:1820-1828
- 493 Coors A, Decaestecker E, Jansen M, De Meester L (2008) Pesticide exposure strongly
494 enhances parasite virulence in an invertebrate host model. Oikos 117:1840-1846
- 495 Crain CM, Kroeker K, Halpern BS (2008) Interactive and cumulative effects of multiple
496 human stressors in marine systems. Ecol. Lett. 11:1304-1315
- 497 Davis DE, Pillai CGP, Truelove B (1976) Effects of Prometryn, Diuron, Fluometuron, and
498 MSMA on Chlorella and two fungi. Weed Sci. 24:587-593
- 499 De Bruin A, Ibelings BW, Kagami M, Mooij WM, Van Donk E (2008) Adaptation of the
500 fungal parasite *Zygorhizidium planktonicum* during 200 generations of growth on
501 homogeneous and heterogeneous populations of its host, the diatom *Asterionella*
502 *formosa*. J. Eukaryot. Microbiol. 55:69-74
- 503 De Bruin A, Ibelings BW, Rijkeboer M, Brehm M, van Donk E (2004) Genetic variation in
504 *Asterionella formosa* (Bacillariophyceae): Is it linked to frequent epidemics of host-
505 specific parasitic fungi? J. Phycol. 40:823-830
- 506 Decaestecker E et al. (2007) Host-parasite 'Red Queen' dynamics archived in pond sediment.
507 Nature 450:870-873
- 508 Doggett MS, Porter D (1996) Sexual reproduction in the fungal parasite, *Zygorhizidium*
509 *planktonicum*. Mycologia 88:720-732

- 510 Duffy MA, Housley JM, Penczykowski RM, Caceres CE, Hall SR (2011) Unhealthy herds:
511 indirect effects of predators enhance two drivers of disease spread. *Funct. Ecol.*
- 512 Duke SO (1990) Overview of herbicide mechanisms of action. *Environ. Health Perspect.*
513 87:263-271
- 514 Duncan AB, Little TJ (2007) Parasite-driven genetic change in a natural population of
515 *Daphnia*. *Evolution* 61:796-803
- 516 Espeland EM, Wetzel RG (2001) Effects of photosynthesis on bacterial phosphatase
517 production in biofilms. *Microb. Ecol.* 42:328-337
- 518 Falkowski P et al. (2000) The global carbon cycle: A test of our knowledge of earth as a
519 system. *Science* 290:291-296
- 520 Field JA, Reed RL, Sawyer TE, Griffith SM, Wigington PJ (2003) Diuron occurrence and
521 distribution in soil and surface and ground water associated with grass seed
522 production. *J. Environ. Qual.* 32:171-179
- 523 Fisher WS, Oliver LM, Walker WW, Manning CS, Lytle TF (1999) Decreased resistance of
524 eastern oysters (*Crassostrea virginica*) to a protozoan pathogen (*Perkinsus marinus*)
525 after sublethal exposure to tributyltin oxide. *Mar. Environ. Res.* 47:185-201
- 526 Gerardo NM, Jacobs SR, Currie CR, Mueller UG (2006) Ancient host-pathogen associations
527 maintained by specificity of chemotaxis and antibiosis. *PLoS Biol.* 4:1358-1363
- 528 Giacomazzi S, Cochet N (2004) Environmental impact of diuron transformation: a review.
529 *Chemosphere* 56:1021-1032
- 530 Gondim-Tomaz RMA, Franco TT, Durrant LR (2005) Biodegradation of diuron and
531 pyruthiobac-sodium by white-rot and soil fungi. In: Calabrese EJ, Kostecki PT,
532 Dragun J (eds) *Contaminated Soils, Sediments and Water: Science in the Real World*,
533 Vol 9, vol 9. Springer, New York, pp 21-32

- 534 Guillard RRL (1973) Division rates. In: Press CU (ed) Handbook of Phycological Methods -
 535 Culture Methods and Growth Measurements. Cambridge Univ. Press, Cambridge, pp
 536 289-311
- 537 Hall SR, Knight CJ, Becker CR, Duffy MA, Tessier AJ, Caceres CE (2009) Quality matters:
 538 resource quality for hosts and the timing of epidemics. Ecol. Lett. 12:118-128
- 539 Hall SR, Sivers-Becker L, Becker C, Duffy MA, Tessier AJ, Caceres CE (2007) Eating
 540 yourself sick: transmission of disease as a function of foraging ecology. Ecol. Lett.
 541 10:207-218
- 542 Holfeld H (2000) Infection of the single-celled diatom *Stephanodiscus alpinus* by the chytrid
 543 *Zygorhizidium*: Parasite distribution within host population, changes in host cell size,
 544 and host-parasite size relationship. Limnol. Oceanogr. 45:1440-1444
- 545 Holmes JC (1996) Parasites as threats to biodiversity in shrinking ecosystems. Biodivers.
 546 Conserv. 5:975-983
- 547 Holmstrup M et al. (2010) Interactions between effects of environmental chemicals and
 548 natural stressors: A review. Sci. Total Environ. 408:3746-3762
- 549 Ibelings BW, De Bruin A, Kagami M, Rijkeboer M, Brehm M, van Donk E (2004) Host
 550 parasite interactions between freshwater phytoplankton and chytrid fungi
 551 (Chytridiomycota). J. Phycol. 40:437-453
- 552 Ibelings BW, Gsell AS, Mooij WM, van Donk E, van den Wyngaert S, Domis LND (2011)
 553 Chytrid infections and diatom spring blooms: paradoxical effects of climate warming
 554 on fungal epidemics in lakes. Freshw. Biol. 56:754-766
- 555 James TY et al. (2006) A molecular phylogeny of the flagellated fungi (Chytridiomycota) and
 556 description of a new phylum (Blastocladiomycota). Mycologia 98:860-871
- 557 Johal GS, Huber DM (2009) Glyphosate effects on diseases of plants. Eur. J. Agron. 31:144-
 558 152

- 559 Jokela J, Lively CM, Dybdahl MF, Fox JA (2003) Genetic variation in sexual and clonal
560 lineages of a freshwater snail. *Biol. J. Linn. Soc.* 79:165-181
- 561 Koprivnikar J, Forbes MR, Baker RL (2007) Contaminant effects on host-parasite
562 interactions: Atrazine, frogs, and trematodes. *Environ. Toxicol. Chem.* 26:2166-2170
- 563 Kreutz LC, Barcellos LJG, Marteninghe A, dos Santos ED, Zanatta R (2010) Exposure to
564 sublethal concentration of glyphosate or atrazine-based herbicides alters the
565 phagocytic function and increases the susceptibility of silver catfish fingerlings
566 (*Rhamdia quelen*) to *Aeromonas hydrophila* challenge. *Fish Shellfish Immunol.*
567 29:694-697
- 568 Kuhn SF (1997) Infection of *Coscinodiscus* spp. by the parasitoid nanoflagellate *Pirsonia*
569 *diadema*: I. Behavioural studies on the infection process. *J. Plankton Res.* 19:791-804
- 570 Lafferty KD, Holt RD (2003) How should environmental stress affect the population
571 dynamics of disease? *Ecol. Lett.* 6:654-664
- 572 Lam E, Kato N, Lawton M (2001) Programmed cell death, mitochondria and the plant
573 hypersensitive response. *Nature* 411:848-853
- 574 Lockert CK, Hoagland KD, Siegfried BD (2006) Comparative sensitivity of freshwater algae
575 to atrazine. *Bull. Environ. Contam. Toxicol.* 76:73-79
- 576 Marcogliese DJ et al. (2009) Combined effects of agricultural activity and parasites on
577 biomarkers in the bullfrog, *Rana catesbeiana*. *Aquat. Toxicol.* 91:126-134
- 578 Mitchell SE, Rogers ES, Little TJ, Read AF (2005) Host-parasite and genotype-by-
579 environment interactions: Temperature modifies potential for selection by a sterilizing
580 pathogen. *Evolution* 59:70-80
- 581 Nipkow HF (1927) Ueber das Verhalten der Skelette planktischer Kieselalgen im
582 geschichteten Tiefenschlamm des Zürich- und Baldeggersees (Dissertation). In. ETH
583 Zürich

- 584 Pulkkinen K, Ebert D (2004) Host starvation decreases parasite load and mean host size in
585 experimental populations. *Ecology* 85:823-833
- 586 Rabiet M, Margoum C, Gouy V, Carluier N, Coquery M (2010) Assessing pesticide
587 concentrations and fluxes in the stream of a small vineyard catchment - Effect of
588 sampling frequency. *Environ. Pollut.* 158:737-748
- 589 Rigby MC, Hechinger RF, Stevens L (2002) Why should parasite resistance be costly? *Trends*
590 *Parasitol.* 18:116-120
- 591 Rohr JR, Raffel TR, Sessions SK, Hudson PJ (2008a) Understanding the net effects of
592 pesticides on amphibian trematode infections. *Ecol. Appl.* 18:1743-1753
- 593 Rohr JR et al. (2008b) Agrochemicals increase trematode infections in a declining amphibian
594 species. *Nature* 455:1235-U1250
- 595 Ronhede S, Jensen B, Rosendahl S, Kragelund BB, Juhler RK, Aamand J (2005)
596 Hydroxylation of the herbicide isoproturon by fungi isolated from agricultural soil.
597 *Appl. Environ. Microbiol.* 71:7927-7932
- 598 Seppälä O, Karvonen A, Haataja M, Kuosa M, Jokela J (2011) Food makes you a target:
599 Disentangling genetic, physiological, and behavioral effects determining susceptibility
600 to infection. *Evolution* 65:1367-1375
- 601 Seppälä O, Liljeroos K, Karvonen A, Jokela J (2008) Host condition as a constraint for
602 parasite reproduction. *Oikos* 117:749-753
- 603 Tang J, Hoagland KD, Siegfried BD (1998) Uptake and bioconcentration of atrazine by
604 selected freshwater algae. *Environ. Toxicol. Chem.* 17:1085-1090
- 605 Vale PF, Little TJ (2009) Measuring parasite fitness under genetic and thermal variation.
606 *Heredity* 103:102-109

- 607 Vallotton N, Eggen RIL, Escher BI, Krayenbühl J, Chèvre N (2008) Effect of pulse herbicidal
608 exposure on *Scenedesmus vacuolatus*: a comparison of two photosystem II inhibitors.
609 Environ. Toxicol. Chem. 27:1399-1407
- 610 Van Donk E, Ringelberg J (1983) The effect of fungal parasitism on the succession of
611 diatoms in Lake Maarsseveen I (The Netherlands). Freshw. Biol. 13:241-252
- 612 Vesela S, Vijverberg J (2007) Effect of body size on toxicity of zinc in neonates of four
613 differently sized *Daphnia* species. Aquat. Ecol. 41:67-73
- 614 Windsor DA (1998) Most of the species on Earth are parasites. Int. J. Parasitol. 28:1939-1941

615 **Table 1**

616 Mixed-effects model to test for the effects of the single and combined stress treatments on
 617 average host population growth rate

Host population growth rate	Estimate	Std. Error	t value	pMCMC
Intercept	0.4204	0.0593	7.093	0.0001
diuron (D)	-0.2062	0.0569	-3.621	0.0014
parasite (P)	0.0524	0.0621	0.843	0.4260
cell volume (CV)	0.0003	0.0002	1.309	0.2686
D×P	0.0645	0.0173	3.731	0.0008
D×CV	0.0005	0.0002	2.010	0.0600
P×CV	-0.0006	0.0003	-2.414	0.0246

618 **Table 2**

619 Mixed-effects model to test for the effects of the single and combined stress treatments on
 620 final uninfected host density

Final uninfected host density:	Estimate	Std. Error	t value	pMCMC
Intercept	461844.0	121025.5	3.816	0.0004
diuron (D)	-392947.7	104548.2	-3.759	0.0002
parasite (P)	-36263.3	87471.3	-0.415	0.6589
cell volume (CV)	-897.2	495.8	-1.810	0.0087
D×P	295654.3	123703.0	2.390	0.0130
D×CV	1225.7	428.3	2.862	0.0032
P×CV	-118.2	358.3	-0.330	0.7253
D×P×CV	-1066.9	506.8	-2.105	0.0278

Table 3

Mixed-effects model to test for the effect of diuron exposure on parasite population growth rate.

Parasite population growth rate:	Estimate	Std. Error	t value	pMCMC
Intercept	-0.8175	0.7452	-1.097	0.342
diuron (D)	0.5656	0.0969	5.835	0.001
cell volume (CV)	0.0124	0.0030	4.086	0.004

Figure legends

Fig. 1 Average host growth rate (a) and final uninfected host density (b) of the five *A. formosa* strains exposed to no stressors (squares), single stressors; parasite (triangles up), diuron (triangles down) and combined stressors (circles). Filled symbols denote observed responses (means and 95% confidence intervals) and empty symbols represent additive effects predicted by the model of independent action. The mean cell volume of each host strain is given in parentheses below each host strain name.

Fig. 2 Relationship between host population cell volume (mean $\pm$ SE) and host fitness inhibition; a) % growth reduction, b) % reduction in final uninfected host density. The broken line shows linear regression analysis for the stressor diuron (black circles) and the unbroken line shows linear regression analysis for the stressor parasite (white circles).

Fig. 3 Relationship between parasite population growth rate (mean $\pm$ SE) and host population cell volume (mean $\pm$ SE). The broken line shows linear regression analysis for the parasite exposed treatment with diuron (black circles) and the unbroken line shows linear regression analysis for the parasite exposed treatment without diuron (white circles).

Fig. 4 Comparison of a) the increase of infections (mean $\pm$ SE), b) host density (mean $\pm$ SE) and c) prevalence of infection (mean $\pm$ SE) between the parasite exposed treatments in the absence of diuron (white circles) and presence of diuron (black circles). Data pooled together for all five host populations.

Figure 1

Figure 2

Figure 3

Figure 4