



Royal Netherlands Academy of Arts and Sciences (KNAW) KONINKLIJKE NEDERLANDSE AKADEMIE VAN WETENSCHAPPEN

Nitrogen fixation and respiratory electron transport in the cyanobacterium *Cyanothece* under different light:dark cycles

Rabouille, S.A.M.; Van de Waal, D.B.; Matthijs, H.C.P.; Huisman, J.

published in

FEMS Microbiology Ecology

2014

DOI (link to publisher)

[10.1111/1574-6941.12251](https://doi.org/10.1111/1574-6941.12251)

document version

Peer reviewed version

[Link to publication in KNAW Research Portal](#)

citation for published version (APA)

Rabouille, S. A. M., Van de Waal, D. B., Matthijs, H. C. P., & Huisman, J. (2014). Nitrogen fixation and respiratory electron transport in the cyanobacterium *Cyanothece* under different light:dark cycles. *FEMS Microbiology Ecology*, 87(3), 630-638. <https://doi.org/10.1111/1574-6941.12251>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the KNAW public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the KNAW public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

pure@knaw.nl

1 **NITROGEN FIXATION AND RESPIRATORY ELECTRON**
2 **TRANSPORT IN THE CYANOBACTERIUM *CYANOTHECE* UNDER**
3 **DIFFERENT LIGHT:DARK CYCLES**

4

5 **Sophie Rabouille^{1,2,3*}, Dedmer B. Van de Waal^{1,4}, Hans C.P. Matthijs¹ and Jef Huisman¹**

6

7

8 ¹ Aquatic Microbiology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, P.O.
9 Box 94248, 1090 GE Amsterdam, The Netherlands.

10 ² UPMC Univ Paris 06, UMR 7093, LOV, Observatoire océanologique, F-06234, Villefranche/mer, France

11 ³ CNRS, UMR 7093, LOV, Observatoire océanologique, F-06234, Villefranche/mer, France

12 ⁴ Department of Aquatic Ecology, Netherlands Institute of Ecology (NIOO-KNAW),
13 P.O. Box 50, 6700 AB Wageningen, The Netherlands.

14

15 Running Title: Timing of N₂ fixation in *Cyanothece*

16

17

* Author for correspondence: email srabouille@obs-vlfr.fr; Tel +33 (0)493763832; Fax +33 (0)493763834

1 Abstract

2 Incompatibility of nitrogen fixation and oxygen production compel unicellular diazotrophic
3 cyanobacteria to perform photosynthesis during daytime and restrict nitrogen fixation to
4 nighttime. The marine diazotroph *Cyanothece* BG 043511 was grown in continuous culture
5 under three light:dark regimes (16L:8D, 12L:12D and 8L:16D hours); we monitored nitrogen
6 fixation and potential photosynthetic efficiency simultaneously online to reveal how their
7 temporal separation is affected by different LD regimes. An increase in nitrogen fixation rate
8 at night coincided with a rise in pulse-amplitude modulated fluorescence, indicating that the
9 enhanced respiratory electron transport to fuel diazotrophy affects the oxidation state of the
10 plastoquinone pool. This may offer an alternative approach to assess instantaneous nitrogen
11 fixation activity. Regardless of photoperiod, the maximum rate of nitrogen fixation was
12 conserved at about 20 hours after the onset of the light. Consequently, nitrogen fixation rates
13 peaked at different moments in the dark: relatively early in the 16L:8D cycle, at midnight in
14 12L:12D, and relatively late in 8L:16D. Under 16L:8D, nitrogen fixation extended into the
15 light, demonstrating the functional plasticity of nitrogen fixation in *Cyanothece*. Highest daily
16 amounts of nitrogen fixed were obtained in 12L:12D, which is consistent with the natural
17 light:dark cycle of subtropical latitudes in which *Cyanothece* thrives.

18

19

20 Keywords: cyanobacteria, light:dark cycle, nitrogen fixation, PAM fluorescence, *Cyanothece*

21 Abbreviations: PSII, photosystem II; PQ pool, plastoquinone pool

22

1

2 **Introduction**

3 Fixation of dinitrogen (N₂) by marine cyanobacteria provides an important source of
4 nitrogen in the tropical and subtropical oceans (Karl *et al.*, 1997). Fixed nitrogen supports 20
5 to 40 % of the total marine primary production (Lee *et al.*, 2002). Predominant open-ocean N₂
6 fixers include the filamentous *Trichodesmium* (Carpenter & Romans, 1991; Letelier & Karl,
7 1996; Capone *et al.*, 1997) and the endosymbiont *Richelia* (Mague *et al.*, 1974; Venrick,
8 1974). Some unicellular cyanobacteria can also fix nitrogen (Wyatt & Silvey, 1969; Mitsui *et*
9 *al.*, 1986), although they were, for many years, not recognized as main actors in the open
10 ocean biogeochemistry. The discovery that unicellular diazotrophic cyanobacteria actually
11 thrive in open oceans, and contribute substantially to the oceanic nitrogen budget (Zehr *et al.*,
12 2001; Moisaner *et al.*, 2010), prompted a reconsideration of the role of different N₂ fixers in
13 the oceanic nitrogen cycle (Zehr *et al.*, 2000; Karl *et al.*, 2002; Zehr, 2011).

14 Autotrophic, unicellular N₂-fixing cyanobacteria are largely confined to the tropics and
15 subtropics (Zehr, 2011). This biogeographical distribution has been attributed to the high
16 temperature requirements of unicellular diazotrophs (e.g., Stal, 2009; Brauer *et al.*, in press).
17 However, it is conceivable that the day-night regime in the (sub)tropics, with an
18 approximately similar duration of day and night, might also play a role. N₂-fixing
19 cyanobacteria handle two processes, mutually exclusive at a first glance: on the one hand,
20 they gather energy and reducing power from oxygenic photosynthesis, while on the other
21 hand the nitrogenase enzyme complex, responsible for N₂ fixation, is inactivated by oxygen
22 (Fay, 1992; Gallon, 1992). To accomplish ‘the impossible’, temporal separation is the
23 common strategy developed by unicellular cyanobacteria, i.e. restraining N₂ fixation to
24 darkness at night when oxygen evolution of light-dependent photosynthesis is absent (Rippka
25 *et al.*, 1971; Mitsui *et al.*, 1986; Bergman *et al.*, 1997). The principal timing of these
26 processes is via the light on-off rhythm, facilitated by a circadian clock (Sherman *et al.*, 1998;

1 Johnson & Golden, 1999; Cervený & Nedbal, 2009; for detail see Cervený *et al.*, 2013). Such
2 a strategy implies that nitrogenase activity cannot be fuelled by photosynthesis directly, but
3 relies instead on carbon reserves accumulated during the light period (Gallon *et al.*, 1988;
4 Schneegurt *et al.*, 1994; Rabouille *et al.*, 2006; Mohr *et al.*, 2010; Dron *et al.*, 2012; Brauer *et*
5 *al.*, 2013). We thus expect that the relative duration of the light and dark period will be of
6 ecological significance for the diazotrophic growth of unicellular cyanobacteria.

7 In this study, we therefore investigate how the diel pattern of N₂ fixation and
8 photosynthesis in the marine, unicellular cyanobacterium *Cyanothece* sp. strain BG 043511
9 (hereafter *Cyanothece*) responds to different light:dark cycles. We monitored N₂ fixation rates
10 by an online nitrogenase activity assay (Staal *et al.*, 2001) and photosynthetic activity by
11 online fluorescence measurements. Interestingly, the results do not only support the view that
12 the photoperiod is an important determinant of the ecological niche of *Cyanothece*, but also
13 reveal that nighttime fluorescence measurements provide a measure of the respiratory electron
14 transport associated with nitrogen fixation.

15

16 **Materials and Methods**

17 *Cyanothece* strain BG 043511, originally isolated as *Synechococcus* BG 043511 near
18 the Bahama Islands in the Atlantic Ocean (León *et al.*, 1986), was kindly provided by the
19 Hawaii Culture Collection (University of Hawaii, Honolulu).

20 *Cyanothece* was grown in continuous culture under three different light:dark (LD)
21 regimes. The light regime at tropical latitudes was represented by a 12L:12D cycle, while
22 16L:8D and 8L:16D cycles reflected summer and winter regimes of the temperate zone. The
23 continuous cultures were specifically designed to study phytoplankton growth (Huisman *et*
24 *al.*, 2002; Agawin *et al.*, 2007), using a flat culture vessel with an optical path length of 5 cm
25 and an effective working volume of 1.6 L. A water jacket placed between the light source and
26 the culture vessel maintained the temperature of the culture at 26°C. The vessels were heat

1 sterilized for 1 h at 121°C prior to inoculation. Cultures were grown in modified YBC II
2 medium devoid of any combined nitrogen source (Chen *et al.*, 1996). The mineral medium
3 was pumped into the culture vessel by a peristaltic pump (Watson Marlow 101U/R MkII) at a
4 dilution rate of $D = 0.12 \text{ d}^{-1}$. A constant culture volume was maintained by a horizontally
5 placed outflow mounted at the surface level of the culture. Homogeneous mixing and
6 sufficient supply of CO₂ were ensured by aerating the cultures with sterile and moistened air
7 enriched with 2.4 % CO₂, at a flow rate of 70 L h⁻¹ regulated by mass flow controllers
8 (Brooks Instruments, Hatfield, PA, USA).

9 Light was supplied by white fluorescent tubes (Philips PLL 24W/840/4P; Philips
10 Lighting, Eindhoven, The Netherlands) that were placed on the front side of the culture vessel
11 (see Huisman *et al.*, 2002). Light intensities (PAR from 400 to 700 nm) were measured with a
12 Licor LI-189 quantum sensor attached to a LI-250 light meter (LI-COR, Lincoln, Nebraska,
13 USA). The incident light intensity (I_{in}) was set at 50 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ by neutral density
14 filters. Light transmitted through the culture vessel (I_{out}) was measured at the back surface of
15 the culture vessel. To account for spatial variation, I_{in} and I_{out} were measured at 14 regularly
16 spaced positions at the front and back surface of the vessel, respectively.

17 At start up, two continuous cultures were exposed to a 12L:12D regime, and stabilized
18 at a steady state at which cell growth equals the dilution rate. Once the population densities
19 approached equilibrium values, one of the continuous cultures was shifted to 16L:8D and the
20 second one to a 8L:16D regime.

21 Cell abundances were measured daily in all cultures, by fixing samples of 2 mL taken in
22 triplicate with 20 μl of a solution of 1% formaldehyde and 10% glutaraldehyde in water.
23 Samples were then frozen with liquid nitrogen and stored at -80°C until cell count analyses
24 with a Coulter Epics Elite flow cytometer (Beckman Coulter Nederland BV, Woerden, The
25 Netherlands). N₂ fixation activity was monitored online using custom designed incubators
26 (Staal *et al.*, 2001). Once or twice per day during four to six consecutive days, samples were

1 taken from the continuous cultures and filtered on pre-combusted (4h at 400°C) and pre-
2 weighted Watson GF/F glass fiber filters. The filters with cellular material were subsequently
3 placed in the custom designed, gas-tight cell incubator which was placed adjacent to the
4 culture vessel and exposed to the same light regime as the continuous culture. The cell
5 incubator was connected to a Shimadzu Gas Chromatograph GC-14B, controlled by
6 Shimadzu Gas Chromatograph software CLASS-VP (Shimadzu, Kyoto, Japan). Samples were
7 incubated during 12 to 24 hours and nitrogenase activity was monitored online every 10 min
8 using the acetylene reduction assay method (Staal *et al.*, 2001). The N₂ fixation rate was
9 calculated according to Stal (1988), with a conversion factor of 4 to calculate N₂ fixation from
10 acetylene reduction.

11 Using the same samples, fluorescence emission was measured every 30 min online by
12 pulse amplitude modulated (PAM) fluorescence using a Walz MINI-PAM (Walz GmbH,
13 Effeltrich, Germany) equipped with white excitation light. Records during the light period
14 were taken without pre-adaptation of the cells to dark conditions. Hence, measurements
15 during the light period gave the in-situ fluorescence (F_s), and a saturating light flash gave the
16 maximum fluorescence of light-adapted cells (F'_m). During the dark period, the cells were
17 obviously dark adapted and the same measurements thus gave what is commonly known as
18 the minimum fluorescence (F_0) and maximum fluorescence (F_m). The quantum yield of
19 photosystem II was calculated as: $\Phi = (F'_m - F_s) / F'_m$ in the light and $(F_m - F_0) / F_m$ in the dark
20 (Genty *et al.*, 1989).

21 Similarities in the fluctuations of the nitrogen fixation rate and quantum yield were
22 investigated with cross-correlation analysis, which calculates the Pearson correlation
23 coefficient between the two time series at different time lags. For this purpose, we improved
24 the stationarity of the time series by rescaling both time series between 0 and 1 on a daily
25 basis. Statistical significance of the cross-correlations was assessed by bootstrapping. Each
26 individual bootstrap sample was represented by the cross-correlation between two artificial

1 time series randomly generated using the same sample size, mean, variance and lag-1
2 autocorrelation of the residuals as the measured time series of the nitrogen fixation rate and
3 quantum yield. A two-sided 95% confidence interval of the cross-correlation coefficient was
4 estimated based on the 2.5 and 97.5 percentile of the bootstrap distribution obtained from a
5 total of 1,000 bootstrap samples.

6

7 **Results**

8 *Population dynamics*

9 Two continuous cultures (C1 and C2) were inoculated at concentrations of $\sim 8 \times 10^5$ cells
10 mL^{-1} , and exposed to a 12L:12D regime. Cell concentrations increased during the transient
11 phase, leading to enhanced self-shading as revealed by a gradual decline in light transmission
12 (I_{out}) (data not shown). At cell densities of $7.33 \pm 1.67 \times 10^6$ cells per mL^{-1} ($n = 9$) in C1 and
13 $7.77 \pm 2.61 \times 10^6$ cells per mL^{-1} ($n = 3$) in C2, and corresponding stabilized I_{out} values of $2.95 \pm$
14 0.12 ($n = 6$) $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ and 2.97 ± 0.23 ($n = 5$) $\mu\text{mol photons m}^{-2} \text{s}^{-1}$, the light
15 regimes were changed. C1 was shifted to 16L:8D and C2 to 8L:16D. Equilibrium was reached
16 within a few days in both cultures, for which cell abundances were $13.2 \pm 1.07 \times 10^6$ cells mL^{-1}
17 ($n = 13$) in C1 (16L:8D) and $9.97 \pm 0.56 \times 10^6$ cells mL^{-1} ($n = 3$) in C2 (8L:16D).
18 Corresponding light transmission values were then $I_{\text{out}} = 1.86 \pm 0.10$ ($n = 6$) $\mu\text{mol photons m}^{-2}$
19 s^{-1} in C1 and $I_{\text{out}} = 2.97 \pm 0.23$ ($n = 5$) $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ in C2.

20

21 *N₂ fixation*

22 At equilibrium, N_2 fixation showed a periodicity of 24 hours under all LD regimes with
23 consistent repeatability. A comparison is made between the 12L:12D regime (control), in
24 which the nitrogenase enzyme was active during the entire dark period, and the 16L:8D and
25 8L:16D regimes (Fig. 1). Under 12L:12D, cells began to fix N_2 at or very shortly before the

1 onset of darkness, nitrogenase activity first increased, peaked between 6.5 h and 9 h into the
2 dark period, and then decreased to a negligible activity at the end of the dark period (Fig. 1a).
3 The average maximum N₂ fixation rate for 12L:12D was 0.87 ± 0.2 pmol N₂ cell⁻¹ h⁻¹ (n = 4)
4 and the total amount of N₂ fixed during an entire day (i.e., 24 hours) averaged at 6.24 ± 1.25
5 pmol N₂ cell⁻¹, from which more than 97 % was acquired in the dark (Table 1). Under
6 16L:8D, nitrogenase activity was not exclusively restricted to the 8 hours of darkness but
7 started already in the light, shortly before the onset of the dark period, and peaked between
8 1.5 and 4 h into the dark period (Fig. 1b). Moreover, after this short dark period, nitrogenase
9 activity continued in the light for about 6 hours, albeit at a lower rate (Fig. 1b). Almost 14%
10 of the total daily N₂ fixation was obtained during the light period (Table 1). Nevertheless,
11 both the total amount of N₂ fixed in the dark and the total daily N₂ fixation were substantially
12 lower in the 16L:8D culture than in the 12L:12D culture (Table 1). In the 8L:16D culture,
13 nitrogenase activity initiated after about 8 h in the dark and lasted for about 8 h (Fig. 1c). The
14 total amount of N₂ fixed during the dark period was again substantially lower than in the
15 12L:12D culture, while we did not observe N₂ fixation activity in the light (Table 1). These
16 results demonstrate that day-night regimes different from 12L:12D reduce the rate and daily
17 amount of N₂ fixation.

18

19 *Fluorescence*

20 To monitor the potential efficiency of photosystem II during the dark period and its
21 actual efficiency during the light period, pulse-amplitude modulated fluorescence was
22 applied, using the 16L:8D and 8L:16D light regimes. The photosynthetic performance of cell
23 samples was studied in the same chamber as the N₂ fixation assays. Fluorescence showed
24 distinct patterns during the light and dark periods. In the dark, maximum fluorescence F_m first
25 increased and then decreased, in a very similar way as the nitrogenase activity. Both
26 maximum fluorescence and quantum yield peaked during periods of highest N₂ fixation

1 activity (Fig. 1b,c). Hence, it appears that the temporal dynamics of PSII fluorescence and
2 nitrogen fixation are closely connected. We noticed that minimum fluorescence during the
3 dark period (F_0) showed constant values throughout each night: values fluctuated by only 3%
4 in the 16L:8D regime and by only 5% in the 8L:16D regime (data not shown). In the 16L:8D
5 culture, where the nitrogenase activity started in the late light period shortly before the onset
6 of the dark, both the maximum fluorescence and quantum yield of PSII also began to increase
7 already in the late light period (Fig. 1b). After the nitrogen fixation rate had reached
8 maximum values in the first half of the short night, the maximum fluorescence and quantum
9 yield started to decrease again. Under the 8L:16D regime (Fig. 1c), the maximum
10 fluorescence and quantum yield both decreased during the entire light period, continued
11 decreasing after the onset of the dark period as long as nitrogenase activity remained
12 undetected, and only started to increase later in the dark, together with nitrogenase activity.

13 Cross-correlation analysis confirmed the similarity in the diel fluctuations of the
14 quantum yield of PSII and the nitrogen fixation rate, both for the 16L:8D regime (Fig. 2a) and
15 the 8L:16D regime (Fig. 2b). For both LD regimes, the cross-correlation was significant. In
16 the 16L:8D regime, the highest value of the cross-correlation coefficient was $r \approx 0.6$ at a time
17 lag of 30 min. In the 8L:16D regime, the cross-correlation coefficient approached $r \approx 0.8$ at
18 time lags of 0 and 30 min. In other words, the temporal variation in quantum yield was
19 closely tracked by the nitrogen fixation rate with a small delay of 0 - 30 min.

20

21 **Discussion**

22 *N₂ fixation dynamics*

23 Our results show that the day-night regime affects the efficiency of N_2 fixation in
24 unicellular cyanobacteria, in agreement with earlier studies (e.g., Ikemoto & Mitsui, 1994;
25 Schneegurt *et al.*, 1994; Sherman *et al.*, 1998; Gallon, 2001; Taniuchi & Ohki, 2007; Toepel
26 *et al.*, 2008; Mohr *et al.*, 2010). In the 12L:12D control, *Cyanothece* showed a typical cyclic

1 nitrogenase activity in which N₂ fixation was restricted to the dark period and peaked around
2 the mid-dark phase (Fig. 1a). Data of the 16L:8D culture show that light periods longer than
3 12 hours did not promote an increase in the gross rate and cumulative yield of nitrogen
4 acquisition. Interestingly, the apparent reduction in dark N₂ fixation capacity as a result of a
5 shorter nighttime was partly compensated by continuation of N₂ fixation in the first 4 to 8 h
6 after the onset of the light period (Fig. 1b). Daytime N₂ fixation added a substantial fraction
7 (14%) to the total amount of N₂ fixed over a 24 h time span. Hence, the present experiment
8 confirms the ability of *Cyanothece* to fix nitrogen in the light (see also, e.g., Červený *et al.*,
9 2013). Despite this continued activity in the light, the total amount of N₂ fixed per cell under
10 16L:8D was about a third lower than in the 12L:12D culture. In the 8L:16D culture, N₂ was
11 fixed only during the second half of the dark period, leading to a daily amount of N₂ fixation
12 that was only half of the 12L:12D control.

13 Our results thus indicate that a 12:12 LD regime provided optimal conditions for N₂
14 fixation by *Cyanothece* BG 043511, as both day lengths longer and shorter than 12 h rendered
15 less product formation by the nitrogenase enzyme. This is most likely caused by an imbalance
16 between the carbon and nitrogen metabolism of the cells. Carbon reserves are needed for the
17 redox and energy requirements of N₂ fixation, but also for covalent bonding of the ammonium
18 ions formed during nitrogen fixation via the GS-GOGAT reaction (Muro-Pastor *et al.*, 2001;
19 von Wobeser *et al.*, 2011; Krasikov *et al.*, 2012). Hence, we hypothesize that a shortage of
20 carbon may limit nitrogen fixation if the light period becomes too short, while there is
21 insufficient time for nitrogen fixation if the nights become too short. N₂ fixation therefore
22 appears to be regulated not only by the prevalence of active nitrogenase enzyme, but also by
23 the accumulation of carbon reserves fueling the nitrogenase activity and providing carbon
24 skeletons for the GS-GOGAT controlled N metabolism.

25 Irrespective of the length of the light period, nitrogenase activity in our cultures was
26 first detected at 12 to 16 hours and reached maximum values at 18 to 21 hours after the onset

1 of the light period. Similar results have been reported for the unicellular diazotroph
2 *Gloeotheca* sp. 68DGA, where nitrogenase activity consistently peaked 16 hours after the
3 onset of the light period, even when the 12L:12D cycle was modified by the addition or
4 subtraction of a single 6 h light period or dark period (Taniuchi & Ohki, 2007). This indicates
5 that the timing of the onset of N₂ fixation is apparently well conserved and obeys a tight
6 connection to the 24 h biological clock. Energy transducing systems, such as photosynthesis,
7 glycolysis, respiration and nitrogen fixation generate cycles of redox and phosphorylation
8 potentials that both modulate metabolic activities and affect gene transcription, via
9 Transcriptional/Translational Control Loops (Stock *et al.*, 2000; Albrechtova *et al.*, 2006).
10 The assembly of the machinery for nitrogen fixation also responds to this control (Chen *et al.*,
11 1998) and this may be the reason why, in the 8L:16D culture, N₂ fixation was postponed to
12 the second half of the night. Moreover, the biochemical processes initiated by these control
13 loops will be temperature dependent, which may explain the recent observation that the
14 timing of the onset of N₂ fixation is delayed at low temperature (Brauer *et al.*, 2013).

15

16 *N₂ fixation and respiratory electron transport*

17 We used pulse-amplitude modulated (PAM) fluorescence to monitor changes in
18 photosynthetic electron transport. During daytime, this technique reports on the quantum
19 yield of photosystem II (Φ), which is commonly used as a measure of the efficiency of
20 photosynthetic electron transport (Schreiber *et al.*, 1986; Genty *et al.*, 1989). Use of the PAM
21 technique during nighttime seems counterintuitive, as there is no photosynthetic electron
22 transport during the night. However, nighttime nitrogen fixation in cyanobacteria makes use
23 of thylakoid membrane bound electron transfer components; in fact, photosynthetic and dark
24 respiratory electron transport both take place in the thylakoid membranes (Hirano *et al.*, 1980;
25 Matthijs *et al.*, 1984a,b; Scherer *et al.*, 1988). These two processes share several components
26 of the electron transport chain including the plastoquinone (PQ) pool and the cytochrome b₆/f

1 complex (Jones & Myers, 1963; Hirano *et al.*, 1980; Scherer *et al.*, 1988; Vermaas, 2001).
2 Thus, electron transfer during photosynthesis and respiration affects the actual redox state of
3 the PQ pool in the thylakoid membranes, which in turn affects the PAM fluorescence signal.

4 During the dark period, measurement of the potential quantum yield of photosystem II
5 informs on the PQ redox state. Typically, the quantum yield of photosystem II remains
6 constant during the dark period in phototrophic eukaryotes such as green algae (Flameling &
7 Kromkamp, 1997). In contrast, the quantum yield of *Cyanothece* showed a bell-shaped
8 pattern in close synchrony with nitrogenase activity (Fig 1b,c). Our explanation is that the
9 increased quantum yield during N₂ fixation at night is caused by enhanced respiratory
10 electron transfer through the thylakoid membranes, intended for ATP generation to support
11 the energy requirements for N₂ fixation. Cross-correlation analysis confirmed the close
12 relationship between the fluctuations in quantum yield and nitrogen fixation rate, and pointed
13 at a small time delay of 0 - 30 min (Fig. 2). This minor time delay may roughly indicate the
14 time between the respiratory electron transport in the thylakoid membrane and the actual
15 fixation of N₂ by the nitrogenase enzyme complex.

16 At the molecular level, these results indicate that Q_A, the first stable acceptor of
17 electrons from PSII, transfers its electrons more efficiently to the plastoquinone pool, due to
18 electron transport associated with the N₂ fixation process (Fig. 3). In particular, enhanced
19 respiration of carbohydrate reserves at night produces NADPH and succinate, which transfer
20 electrons via NADPH dehydrogenase (NDH) and succinate dehydrogenase (SDH; Cooley &
21 Vermaas, 2001) into the PQ pool and onwards to the terminal electron acceptor oxygen via
22 cytochrome c and cytochrome oxidase (Fig. 3). This respiratory electron transport sets up a
23 proton gradient across the thylakoid membrane, which drives ATP production. N₂ fixation in
24 unicellular diazotrophs is probably the most energy-consuming process in the dark and thus
25 constitutes a drain of both ATP and electrons (Fig. 3). High activity of the electron transport
26 chain, to sustain the ATP production required for N₂ fixation, results in relative oxidation of

1 the PQ pool. An oxidized state of the PQ pool is measured as a high “quantum yield” (Genty
2 *et al.*, 1989), and hence changes in nitrogenase activity are accompanied by changes in the
3 apparent quantum yield of PSII measured by PAM fluorescence. This finding has two
4 important implications: (i) during daytime, changes in PAM fluorescence of N₂-fixing
5 cyanobacteria do not necessarily reflect changes in their photosynthetic activity, because N₂
6 fixation also affects the fluorescence signal, and (ii) at night, PAM fluorescence
7 measurements of unicellular, N₂-fixing cyanobacteria such as *Cyanothece* can provide an
8 experimentally easy method to monitor their actual N₂ fixation activity.

9 In conclusion, our results revealed a tight link between respiratory electron transport
10 and nocturnal nitrogenase activity, and support the hypothesis that the day-night regime is an
11 important determinant of the ecological niche and biogeographical distribution of unicellular
12 diazotrophic cyanobacteria. Nitrogen acquisition in *Cyanothece* BG 043511 was maximized
13 when the day-night regime provided balanced accumulation of both carbon and nitrogen. It
14 would be interesting to investigate whether the same 12:12 LD cycle would also be optimal
15 for other *Cyanothece* strains with a different timing of their nitrogenase activity peak
16 (Bandyopadhyay *et al.*, 2013). While the biogeographical distribution of unicellular
17 diazotrophs is currently being revealed (Moisander *et al.*, 2010), the finding that a 12:12
18 light:dark cycle appears to yield the highest daily nitrogen fixation rate might offer an elegant
19 explanation why unicellular diazotrophs seem particularly successful at (sub)tropical
20 latitudes.

21

22

23 **Acknowledgements**

24 The authors are grateful to the anonymous reviewers for their helpful comments on the
25 manuscript, to Yannick Huot and Hugh MacIntyre for thoughtful discussions on the
26 chlorophyll fluorescence data and to Anouk Blauw for her advice on the cross-correlation

1 analysis. The research of S. Rabouille was supported by the European Commission (contract
2 MEIF-CT-2003-500516) and the CYBER-LEFE project CROCOCYCLE. The research of
3 D.B. Van de Waal and J. Huisman was supported by the Earth and Life Sciences Foundation
4 (ALW), which is subsidized by the Netherlands Organization for Scientific Research (NWO).

5
6

7 **References**

8

- 9 Agawin NSR, Rabouille S, Veldhuis MJW, Servatius L, Hol S, van Overzee HMJ & Huisman
10 J (2007) Competition and facilitation between unicellular nitrogen-fixing cyanobacteria
11 and non-nitrogen-fixing phytoplankton species. *Limnology and Oceanography* **52**: 2233-
12 2248.
- 13 Albrechtová JTP, Vervliet-Scheebaum M, Normann J, Veit J & Wagner E (2006) Metabolic
14 control of transcriptional-translational control loops (TTCL) by circadian oscillations in
15 the redox- and phosphorylation state of cells. *Biological Rhythm Research* **37**: 381-389.
- 16 Bandyopadhyay A, Elvitigala T, Liberton M & Pakrasi HB (2013) Variations in the rhythms
17 of respiration and nitrogen fixation in members of the unicellular diazotrophic
18 cyanobacterial genus *Cyanothece*. *Plant Physiology* **161**: 1334-1346.
- 19 Bergman B, Gallon JR, Rai AN & Stal LJ (1997) N₂ fixation by non-heterocystous
20 cyanobacteria. *FEMS Microbiology Reviews* **19**: 139-185.
- 21 Brauer VS, Stomp M, Rosso C, van Beusekom SAM, Emmerich B, Stal LJ & Huisman J
22 (2013) Low temperature delays timing and enhances the cost of nitrogen fixation in the
23 unicellular cyanobacterium *Cyanothece*. *ISME Journal* (advance online publication). doi:
24 10.1038/ismej.2013.103
- 25 Capone DG, Zehr JP, Paerl HW, Bergman B & Carpenter EJ (1997) *Trichodesmium*, a
26 globally significant marine cyanobacterium. *Science* **276**: 1221-1229.

- 1 Carpenter EJ & Romans K (1991) Major role of the cyanobacterium *Trichodesmium* in
2 nutrient cycling in the North Atlantic Ocean. *Science* **254**: 1356-1358.
- 3 Červený J & Nedbal L (2009) Metabolic rhythms of the cyanobacterium *Cyanothece* sp.
4 ATCC 51142 correlate with modeled dynamics of circadian clock. *Journal of Biological*
5 *Rhythms* **24**: 295-303.
- 6 Červený J, Sinetova MA, Valledor L, Sherman LA & Nedbal L (2013) Ultradian metabolic
7 rhythm in the diazotrophic cyanobacterium *Cyanothece* sp. ATCC 51142. *Proceedings of*
8 *the National Academy of Sciences USA* (advance online publication). doi:
9 10.1073/pnas.1301171110.
- 10 Chen YB, Zehr JP & Mellon M (1996) Growth and nitrogen fixation of the diazotrophic
11 filamentous nonheterocystous cyanobacterium *Trichodesmium* sp. IMS 101 in defined
12 media: Evidence for a circadian rhythm. *Journal of Phycology* **32**: 916-923.
- 13 Chen YB, Dominic B, Mellon MT & Zehr JP (1998) Circadian rhythm of nitrogenase gene
14 expression in the diazotrophic filamentous nonheterocystous cyanobacterium
15 *Trichodesmium* sp. strain IMS 101. *Journal of Bacteriology* **180**: 3598-3605.
- 16 Cólón-López MS, Sherman DM & Sherman LA (1997) Transcriptional and translational
17 regulation of nitrogenase in light- dark- and continuous-light-grown cultures of the
18 unicellular cyanobacterium *Cyanothece* sp. strain ATCC 51142. *Journal of Bacteriology*
19 **179**: 4319-4327.
- 20 Cólón-López MS & Sherman LA (1998) Transcriptional and translational regulation of
21 photosystem I and II genes in light-dark- and continuous-light-grown cultures of the
22 unicellular cyanobacterium *Cyanothece* sp. strain ATCC 51142. *Journal of Bacteriology*
23 **180**: 519-526.
- 24 Cooley JW & Vermaas WFJ (2001) Succinate dehydrogenase and other respiratory pathways
25 in thylakoid membranes of *Synechocystis* sp strain PCC 6803: Capacity comparisons and
26 physiological function. *Journal of Bacteriology* **183**: 4251-4258.

- 1 Dron A, Rabouille S, Claquin P, Le Roy B, Talec A & Sciandra A (2012) Light-dark (12:12)
2 cycle of carbon and nitrogen metabolism in *Crocospaera watsonii* WH8501: relation to
3 the cell cycle. *Environmental Microbiology* **14**: 967-981.
- 4 Fay, P. (1992) Oxygen relations of nitrogen fixation in cyanobacteria. *Microbiological*
5 *Reviews* **56**: 340-373.
- 6 Flameling IA & Kromkamp J (1997) Photoacclimation of *Scenedesmus protuberans*
7 (Chlorophyceae) to fluctuating irradiances simulating vertical mixing. *Journal of Plankton*
8 *Research* **19**: 1011-1024.
- 9 Gallon JR (1992) Reconciling the incompatible: N₂ fixation and O₂. *New Phytologist* **122**:
10 571-609.
- 11 Gallon JR (2001) N₂ fixation in phototrophs: adaptation to a specialized way of life. *Plant and*
12 *Soil* **230**: 39-48.
- 13 Gallon JR, Perry SM, Rajab TMA, Flayeh KAM, Yunes JS & Chaplin AE (1988) Metabolic
14 changes associated with the diurnal pattern of N₂ fixation in *Gloeotheca*. *Journal of*
15 *General Microbiology* **134**: 3079-3087.
- 16 Genty B, Briantais JM & Baker NR (1989) The relationship between the quantum yield of
17 photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochimica*
18 *et Biophysica Acta* **990**: 87-92.
- 19 Hirano M, Satoh K & Katoh S (1980) Plastoquinone as a common link between
20 photosynthesis and respiration in a blue-green alga. *Photosynthesis Research* **1**: 149-162.
- 21 Huisman J, Matthijs HCP, Visser PM *et al.* (2002) Principles of the light-limited chemostat:
22 theory and ecological applications. *Antonie van Leeuwenhoek* **81**: 117-133.
- 23 Ikemoto H & Mitsui A (1994) Diazotrophic synchronous growth of a marine unicellular
24 cyanobacterium, *Synechococcus* sp. strain Miami BG 043511, under aerobic and
25 microaerobic/anaerobic conditions. *Microbiology-UK* **140**: 2153-2158.
- 26 Johnson CH & Golden SS (1999) Circadian programs in cyanobacteria: Adaptiveness and
27 mechanism. *Annual Review of Microbiology* **53**: 389-409.

- 1 Jones LW & Myers J (1963) A common link between photosynthesis and respiration in a
2 blue-green alga. *Nature* **199**: 670-672.
- 3 Karl D, Letelier R, Tupas L, Dore J, Christian J & Hebel D (1997) The role of nitrogen
4 fixation in biogeochemical cycling in the subtropical North Pacific Ocean. *Nature* **388**:
5 533-538.
- 6 Karl D, Michaels A, Bergman B *et al.* (2002) Dinitrogen fixation in the world's oceans.
7 *Biogeochemistry* **57/58**: 47-98.
- 8 Krasikov V, von Wobeser EA, Dekker HL, Huismann J & Matthijs HCP (2012) Time-series
9 resolution of gradual nitrogen starvation and its impact on photosynthesis in the
10 cyanobacterium *Synechocystis* PCC 6803. *Physiologia Plantarum* **145**: 426-439.
- 11 Lee K, Karl DM, Wanninkhof R & Zhang JZ (2002) Global estimates of net carbon
12 production in the nitrate-depleted tropical and subtropical oceans. *Geophysical Research*
13 *Letters* **29**: ARTN 1907.
- 14 León C, Kumazawa S & Mitsui A (1986) Cyclic appearance of aerobic nitrogenase activity
15 during synchronous growth of unicellular cyanobacteria. *Current Microbiology* **13**: 149-
16 153.
- 17 Letelier RM & Karl DM (1996) Role of *Trichodesmium* spp. in the productivity of the
18 subtropical North Pacific Ocean. *Marine Ecology-Progress Series* **133**: 263-273.
- 19 Mague TH, Weare NM & Holm-Hansen O (1974) Nitrogen fixation in the North Pacific
20 Ocean. *Marine Biology* **24**: 109-119.
- 21 Matthijs HCP, Luderus EME, Scholts MJC & Kraayenhof R (1984a) Energy-metabolism in
22 the cyanobacterium *Plectonema boryanum*: oxidative phosphorylation and respiratory
23 pathways. *Biochimica et Biophysica Acta* **766**: 38-44.
- 24 Matthijs HCP, Luderus EME, Loffler HJM, Scholts MJC & Kraayenhof R (1984b) Energy-
25 metabolism in the cyanobacterium *Plectonema boryanum*: participation of the thylakoid
26 photosynthetic electron-transfer chain in the dark respiration of NADPH and NADH.
27 *Biochimica et Biophysica Acta* **766**: 29-37.

- 1 Mitsui A, Cao S, Takahashi A & Arai T (1987) Growth synchrony and cellular parameters of
2 the unicellular nitrogen-fixing marine cyanobacterium, *Synechococcus* sp. strain Miami
3 BG 043511 under continuous illumination. *Physiologia Plantarum* **69**: 1-8.
- 4 Mitsui A, Kumazawa S, Takahashi A, Ikemoto H, Cao S & Arai T (1986) Strategy by which
5 nitrogen-fixing unicellular cyanobacteria grow photoautotrophically. *Nature* **323**: 720-
6 722.
- 7 Mohr W, Intermaggio MP & LaRoche J (2010) Diel rhythm of nitrogen and carbon
8 metabolism in the unicellular, diazotrophic cyanobacterium *Crocospaera watsonii*
9 WH8501. *Environmental Microbiology* **12**: 412-421.
- 10 Moisander PH, Beinart RA, Hewson I *et al.* (2010) Unicellular cyanobacterial distributions
11 broaden the oceanic N₂ fixation domain. *Science* **327**: 1512-1514.
- 12 Muro-Pastor MI, Reyes JC & Florencio FJ (2001) Ammonium assimilation in cyanobacteria.
13 *Photosynthesis Research* **83**: 135-150.
- 14 Rabouille S, Staal M, Stal LJ & Soetaert K (2006) Modeling the dynamic regulation of
15 nitrogen fixation in the cyanobacterium *Trichodesmium* sp. *Applied and Environmental*
16 *Microbiology* **72**: 3217-3227.
- 17 Rippka R, Neilson A, Kunisawa R & Cohen-Bazire G (1971) Nitrogen fixation by unicellular
18 blue-green algae. *Archiv für Mikrobiologie* **76**: 341-348.
- 19 Scherer S, Almon H & Böger P (1988) Interaction of photosynthesis, respiration and nitrogen
20 fixation in cyanobacteria. *Photosynthesis Research* **15**: 95-114.
- 21 Schneegurt MA, Sherman DM, Nayar S & Sherman LA (1994) Oscillating behavior of
22 carbohydrate granule formation and dinitrogen fixation in the cyanobacterium *Cyanothece*
23 sp. strain ATCC 51142. *Journal of Bacteriology* **176**: 1586-1597.
- 24 Schreiber U, Schliwa U & Bilger W (1986) Continuous recording of photochemical and non-
25 photochemical chlorophyll fluorescence quenching with a new type of modulation
26 fluorometer. *Photosynthesis Research* **10**: 51-62.

- 1 Sherman LA, Meunier P & Colon-Lopez MS (1998) Diurnal rhythms in metabolism: A day in
2 the life of a unicellular, diazotrophic cyanobacterium. *Photosynthesis Research* **58**: 25-42.
- 3 Staal M, Lintel-Hekkert ST, Harren F & Stal LJ (2001) Nitrogenase activity in cyanobacteria
4 measured by the acetylene reduction assay: a comparison between batch incubation and
5 on-line monitoring. *Environmental Microbiology* **3**: 343-351.
- 6 Stal LJ (1988) Nitrogen fixation in cyanobacterial mats. *Methods in Enzymology*,
7 *Cyanobacteria, Vol. 167* (Packer L & Glazer AN, eds), pp. 474-484. Academic Press, San
8 Diego.
- 9 Stal LJ (2009) Is the distribution of nitrogen-fixing cyanobacteria in the oceans related to
10 temperature? *Environmental Microbiology* **11**: 1632-1645.
- 11 Stock AM, Robinson VL & Goudreau PN (2000) Two-component signal transduction.
12 *Annual Review of Biochemistry* **69**: 183-215.
- 13 Taniuchi Y & Ohki K (2007) Relation between nitrogenase synthesis and activity in a marine
14 unicellular diazotrophic strain, *Gloeotheca* sp 68DGA (Cyanophyte), grown under
15 different light/dark regimens. *Phycological Research* **55**: 249-256.
- 16 Toepel J, Welsh E, Summerfield TC, Pakrasi HB & Sherman LA (2008) Differential
17 transcriptional analysis of the cyanobacterium *Cyanothece* sp. strain ATCC 51142 during
18 light-dark and continuous-light growth. *Journal of Bacteriology* **190**: 3904-3913.
- 19 Venrick EL (1974) The distribution and significance of *Richelia intracellularis* Schmidt in the
20 North Pacific Central Gyre. *Limnology and Oceanography* **19**: 437-445.
- 21 Vermaas WFJ (2001) Photosynthesis and respiration in cyanobacteria. In: Encyclopedia of
22 Life Sciences. John Wiley & Sons (eds), New York.
- 23 von Wobeser EA, Ibelings BW, Bok J, Krasikov V, Huisman J & Matthijs HCP (2011)
24 Concerted changes in gene expression and cell physiology of the cyanobacterium
25 *Synechocystis* sp strain PCC 6803 during transitions between nitrogen and light-limited
26 growth. *Plant Physiology* **155**: 1445-1457.
- 27 Wyatt JT & Silvey JKG (1969) Nitrogen fixation by *Gloeocapsa*. *Science* **165**: 908-909.

1 Zehr JP (2011) Nitrogen fixation by marine cyanobacteria. *Trends in Microbiology* **19**: 162-
2 173.

3 Zehr JP, Carpenter EJ & Villareal TA (2000) New perspectives on nitrogen-fixing
4 microorganisms in tropical and subtropical oceans. *Trends in Microbiology* **8**: 68-73.

5 Zehr JP, Waterbury JB, Turner PJ *et al.* (2001) Unicellular cyanobacteria fix N₂ in the
6 subtropical North Pacific Ocean. *Nature* **412**: 635-638.

7
8
9

1 **Figure Legends**

2

3 **Figure 1:** Dynamics of N_2 fixation (closed circles), maximum fluorescence F'_m in the light
4 and F_m in the dark (open circles), and the quantum yield of photosystem II (crosses) in
5 continuous cultures exposed to different light:dark regimes. a) 12L:12D regime (fluorescence
6 data not available), b) 16L:8D regime, and c) 8L:16D regime. Shaded areas represent dark
7 periods.

8

9 **Figure 2:** The estimated cross-correlation function between time series of the quantum yield
10 of PSII and the nitrogen fixation rate. a) 16L:8D regime and b) 8L:16D regime. Shaded areas
11 represent the 95% confidence interval based on a bootstrapping approach using the same
12 sample size, mean, variance and lag-1 autocorrelation as in the measured time series. The
13 vertical, dashed line locates the highest cross correlation and the corresponding lag time.

14

15 **Figure 3:** Schematic representation of electron flow within the thylakoid membrane,
16 illustrating the role of the plastoquinone (PQ) pool. a) In the light, electron fluxes are
17 essentially related to the light reactions of photosynthesis: a linear transport of electrons into
18 the PQ pool originates from PSII, and are transported to the first stable terminal acceptor
19 ferredoxin (Fd) by photosystem I (PSI). b) In the dark, light reactions of photosynthesis are
20 inactive; electron fluxes are related to respiration only and reduction of the PQ pool occurs
21 through oxidation of NADPH and succinate during the respiration of carbon reserves. $h\nu$:
22 light energy; $C_{b/f}$: cytochrome b_6/f ; NDH: NADPH dehydrogenase; NR: NADP reductase; PQ:
23 oxidized plastoquinone pool; PQH₂: reduced plastoquinone pool; PSI: photosystem I; PSII:
24 photosystem II; SDH: succinate dehydrogenase.

25

1

2 **Table 1.** Characteristics of the N₂ fixation rate (\pm SD) under three different light:dark regimes.

LD regime (h:h)	Maximum N₂ fixation rate (pmol N₂ cell⁻¹ h⁻¹)	Total N₂ fixed in the dark (pmol N₂ cell⁻¹)	Total N₂ fixed in 24 hours (pmol N₂ cell⁻¹)	Percentage of N₂ fixed in the light (%)
12:12	0.87 \pm 0.20	6.09 \pm 1.23	6.24 \pm 1.25	2.4
16:8	0.81 \pm 0.18	3.78 \pm 0.50	4.39 \pm 0.38	13.9
8:16	0.74 \pm 0.04	3.14 \pm 0.29	3.14 \pm 0.29	0

3

4

5