

# *Acidobacteria* strains from subdivision 1 act as plant growth-promoting bacteria

Anna M. Kielak<sup>1</sup> · Matheus A. P. Cipriano<sup>1</sup> · Eiko E. Kuramae<sup>1</sup>

Received: 21 March 2016 / Revised: 9 June 2016 / Accepted: 15 June 2016  
© The Author(s) 2016. This article is published with open access at Springerlink.com

**Abstract** *Acidobacteria* is one of the most abundant phyla in soils and has been detected in rhizosphere mainly based on cultivation-independent approaches such as 16S rRNA gene survey. Although putative interaction of *Acidobacteria* with plants was suggested, so far no plant–bacterial interactions were shown. Therefore, we performed several in vitro tests to evaluate *Acidobacteria*–plant interactions and the possible mechanisms involved in such interaction. We observed that *Arabidopsis thaliana* inoculated with three strains belonging to *Acidobacteria* subdivision 1 showed increase in biomass of roots and shoots as well as morphological changes in root system. Our results indicate that the plant hormone indole-3-acetic acid production and iron acquisition are plausibly involved in the plant and *Acidobacteria* interactions. Here, we confirm for the first time that *Acidobacteria* can actively interact with plants and act as plant growth-promoting bacteria. In addition, we show that *Acidobacteria* strains produce exopolysaccharide which supports the adhesion of bacteria to the root surfaces.

**Keywords** *Acidobacteria* · PGPB · IAA · Root biomass · *Arabidopsis thaliana*

## Introduction

*Acidobacteria* is a very diverse and ubiquitous bacterial phylum. Furthermore, those bacteria seem to be especially well adapted to soil environment, often representing one the most abundant bacterial phylum (Janssen 2006; Lee et al. 2008). Although there are inconsistencies in reports regarding the preference of *Acidobacteria* in inhabiting bulk versus rhizosphere soils (Fierer et al. 2007; Singh et al. 2007; Kielak et al. 2008), there are clear evidences for the association of some *Acidobacteria* with plants (da Rocha et al. 2010, 2013). The enormous phylogenetic diversity within the phylum also suggests that *Acidobacteria* are genetically and, most likely, metabolically dissimilar; thus, the results of single studies cannot be generalized and easily extrapolated to the whole phylum.

Due to the still low number of sequenced genomes and difficulties associated with cultivation, the ecological role of this phylum remains rather unknown (Kielak et al. 2016). Nevertheless, a number of studies have compared distribution and diversity of *Acidobacteria* in relation to plant root proximity (Chow et al. 2002; Filion et al. 2004; da Rocha et al. 2010; Chaparro et al. 2014) and/or plant exudates (Shi et al. 2011; Mao et al. 2014). For example, acidobacterial strains have been obtained from internal plant tissues hinting to an endophytic lifestyle (Idris et al. 2004; Nissinen et al. 2012; Poosakkannu et al. 2015). Mendes et al. (2014) using culture-independent approach technique have shown that *Acidobacteria* are overrepresented in soybean rhizosphere, and da Rocha et al. (2010) have reported by means of qPCR, the *Holophagae* (*Acidobacteria* subdivision 8) being more abundant in leek rhizosphere. However, in the second case bacterial cell number was lower in spheres very proximate to roots or on the root surface itself.

Communicated by Erko Stackebrandt.

✉ Eiko E. Kuramae  
e.kuramae@nioo.knaw.nl

<sup>1</sup> Department of Microbial Ecology, Netherlands Institute of Ecology (NIOO-KNAW), P.O. Box 50, Droevendaalsesteeg 10, 6708 PB Wageningen, The Netherlands

*Acidobacteria* were also shown to be dominant in the rhizosphere of *Arabidopsis thaliana*, and moreover, change in terms of phylum composition and abundance during plant development, possibly due to changes in plant exudation (Chaparro et al. 2014). However, most of the studies were based on culture-independent method based on 16S ribosomal gene marker sequencing due to difficulty to culture *Acidobacteria* and perform experiments under laboratory conditions. Thus, those type of studies do not investigate the nature of plant–bacteria interactions. Concerning few studies on *Acidobacteria* physiology, non-traditional sources of carbon such as complex polysaccharides were suggested to improve cultivability of *Acidobacteria* (Koch et al. 2008; Pankratov and Dedysh 2010; Eichorst et al. 2011). Also some of the characterized strains were shown to be able to utilize plant-derived polymers (Pankratov et al. 2008, 2012; Eichorst et al. 2011) further suggesting close relation between plants and specific *Acidobacteria* subdivisions. Nevertheless, available strains can be used for attempts of studying interactions with plants under experimental conditions. In order to test the hypothesis that *Acidobacteria* strains effect plant growth, we assayed the interactions of three class *Acidobacteria* strains with *A. thaliana* ecotype Columbia 0 (Col 0) under in vitro conditions.

## Materials and methods

### Bacterial strains

Three *Acidobacteria* strains belonging to the class *Acidobacteria* from the NIOO-KNAW microorganisms' collection were used in this study. Two of the strains are affiliated with genus *Granulicella*, namely *Granulicella* sp. WH15 (Valášková et al. 2009) and 5B5 (KM979383), and one is a type strain of the genus *Acidicapsa*, *A. ligni* WH120T (Valášková et al. 2009; Kulichevskaya et al. 2012). *Pseudomonas putida* IAC-RBa14 (KJ590499) and *Escherichia coli* WA321 (DSM no. 4509) strains were used as positive and negative controls of plant growth-promoting bacteria, respectively.

### Plant–bacteria interaction experiment

*Arabidopsis thaliana* ecotype Columbia 0 seeds were surface sterilized by washing in 70 % ethanol for 5 min followed by submerging for 10 min in 50 % bleach and rinsing four times with sterile distilled water. Sterile seeds were placed on half-strength Murashige and Skoog (MS) medium pH 5.7 (Murashige and Skoog 1962) supplemented with 12 g L<sup>-1</sup> plant agar (Duchefa Biochemie bv) and 5 g L<sup>-1</sup> sucrose. Six plants were grown per plate. Seedlings were incubated at 21 °C with the light cycle of

photoperiod 16 h/8 h day/night. The root tips of 5-day-old seedlings were inoculated with 2.5 µL of bacterial suspension (phosphate saline buffer, pH 5.5) of OD600 = 1 corresponding to 1.7 × 10<sup>6</sup>, 1.5 × 10<sup>7</sup> and 1.3 × 10<sup>7</sup> CFU for *A. ligni* WH120T, *Granulicella* sp. 5B5 and WH15, respectively, or by direct transfer from the solid medium (0.1 × TSA, pH 5.0 see below) as an alternative method. The effect of bacteria on plant growth was evaluated 3 weeks post-inoculation.

### Indole acetic acid (IAA) production


IAA production was determined based on the method described by Bric et al. (1991). *Acidobacteria* strains and *P. putida* IAC-RBa14 (positive control) were inoculated on 0.1 × tryptone soy agar (TSA) plates (pH 5.0 and 5.7) supplemented with 5 mM L<sup>-1</sup> tryptophan and covered with a cellulose nitrate filter (0.45 µm pore size, Sartorius). TSA contained 1 g L<sup>-1</sup> NaCl, 3.0 g L<sup>-1</sup> TSB (Oxoid), 1.95 g L<sup>-1</sup> MES, 20 g L<sup>-1</sup> agar (Boom, Netherlands). Plates were incubated at 20 °C until colonies reached approximately 4–5 mm diameter (5 days), and then the membranes were washed in the Salkowski reagent (1.2 % FeCl<sub>2</sub> in 37 % sulfuric acid). The reaction was allowed to proceed for 30 min at RT until purple color appeared. All strains were tested in triplicates on separate plates.

### Phosphate solubilization assay

*Acidobacteria* strains were tested for their ability to solubilize a mineral form of phosphate. *P. putida* IAC-RBa14 was used as a positive control. Tests were performed on the National Botanical Research Institute's phosphate growth medium (NBRI PM) containing per liter 15 g agar–agar ultrapure (Merck KGaA), 10 g glucose, 5 g Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>, 5 g MgCl<sub>2</sub>·6H<sub>2</sub>O, 0.25 g MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.2 g KCl and 0.1 g (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (Nautiyal 1999). All strains were inoculated by transfer from the 0.1 × TBA pH 5.0 media using inoculation loop and incubated for 6 weeks at 20 °C. The clearing zones around the colonies indicated phosphate solubilization by the isolates. The experiment was carried out in triplicate on separate plates.

### Siderophore production

Detection of siderophore production was carried out in chrome azurol S (CAS) agar plates. Removal of iron from the CAS dye by iron-chelating compounds results in a color change from blue to yellow/orange. The CAS medium was prepared according to the method described by Schwyn and Neilands (1987). Bacteria were collected from 0.1 × TSA plates, resuspended and washed twice with phosphate-buffered saline pH 6.5. An aliquot of 10 µL



**Fig. 1** Effects of inoculation with *Acidobacteria* strains on *Arabidopsis thaliana* seedlings. Root tips of 5-day-old seedlings were inoculated by direct transfer with a loop of bacteria grown on  $0.1 \times$  TSA medium, pH 5.0. **A** Changes in morphology. Image was

taken 3 weeks post-inoculation. **B** Changes in root length and fresh biomass. Different letters (*a*, *b*) indicate statistically significant differences ( $P < 0.05$ ) between inoculated and control plants according to *t* test. Error bars represent SD ( $n = 6$  plates each with six plants)


of bacterial suspension was spotted on CAS agar plates. Plates were checked daily for color change around each colony. *E. coli* WA321 was used as a positive control.

### *nifH* targeting PCR

The *nifH* gene targeting PCR was performed according to the modified protocol by Brankatschk et al. (2012). PCR amplification was performed in a 25- $\mu$ L reaction mixture including DNA template, 0.6  $\mu$ M of primers (*nifHF/nifHR*), 200  $\mu$ M dNTPs, 1 $\times$  of Taq buffer and 0.04 U FastStart High Fidelity Taq Enzyme Blend (Roche). The PCR were performed under the following conditions: initial denaturation step 5 min at 95  $^{\circ}$ C, followed by touch-down cycles of denaturation for 15 s at 95  $^{\circ}$ C, annealing starting at 63  $^{\circ}$ C with temperature decreases of 2  $^{\circ}$ C per two cycles and elongation at 72  $^{\circ}$ C for 45 s followed by 30 cycles with annealing at 53  $^{\circ}$ C. The final extension was extended to 10 min at 72  $^{\circ}$ C.

## Results and discussion

In this study, we tested three acidobacterial strains for possible interactions with *A. thaliana* (Col 0) roots. The growth of the plantlets was clearly positively affected by the presence of bacteria (Fig. 1). The presented results are shown with bacteria transferred directly from the solid media since this method of inoculation resulted in a stronger plant response. Root length, lateral root formation and root hair number were increased in plants exposed to *Acidobacteria* strains used in this study (Fig. 1A). Moreover, the root biomass increased significantly for plantlets inoculated with all three strains (Fig. 1B). The improved root architecture,




**Fig. 2** Root colonization and biofilm formation around *Arabidopsis thaliana* roots by strain 5B5

more lateral branches and/or higher number of root hairs assist in more efficient water and nutrients uptake (Herder et al. 2010). Increased shoot biomass was also observed; however, the differences were not significant. We hypothesize that the stronger effect observed on plant growth with bacteria from the growth media in comparison with the bacterial suspension is not only related to higher bacterial biomass in the inoculum but also to the stress and longer adaptation time experienced by bacteria under the unfavorable culture conditions.

Bacterial adhesion, biofilm formation and growth along the root surfaces were observed for all three strains (Fig. 2). *Acidobacteria* strains *Granulicella paludicola*, *G. pectinivorans*, *G. aggregans* and *G. rosea* (Pankratov and Dedysch 2010), *Acidicapsa borealis* and *A. ligni* (Kulichevskaya




**Fig. 3** Isolates were tested for indole acetic acid (IAA) production by bacteria immobilization on a nitrocellulose membrane followed by washing the membrane with Salkowski reagent. *P. putida* IAC-RBa4 was used as a positive control. **A** *P. putida* IAC-RBa4, **B** strain WH120T, **C** strain 5B5 and **D** strain WH15



et al. 2012) and *Terriglobus tenax* (Whang et al. 2014) were proven to produce extracellular polysaccharide. By genome mining, Ward et al. (2009) have suggested *Acidobacteria* being involved in soil matrix formation, water and nutrition trapping, or bacterial adhesion that lead to soil aggregate formation. However, here, for the first time we show that *Acidobacteria* strains produce exopolysaccharide (EPS) in the adhesion of bacteria to the root surfaces.


Bacteria can have positive effect on plant growth indirectly by acting as a biocontrol agent or directly by modulating plant hormone levels (Hayat et al. 2010) or/and by facilitating resource acquisition (mostly nitrogen, phosphorus and iron). Among the plant hormones, the auxin indole-3-acetic acid (IAA) has received most of the attention. As auxin production could best explain the observed changes in the plant phenotypes, we tested our strains for production of this phytohormone. A color change due to IAA production was observed for all three strains indicating all of them as positive for indolic substances production (Fig. 3).

Moreover, we tested our strains for nutrient acquisition abilities. Phosphorus (P) is an important macronutrient for plant growth and development. However, in general, the concentration of soluble P in soil is quite low. It is postulated that bacteria can enhance the P acquisition of



**Fig. 4** Isolates were tested for phosphate solubilization on agar plate using National Botanical Research Institute's phosphate medium. *P. putida* IAC-RBa4 was used as a positive control

plants (Richardson and Simpson 2011). All three strains were proven to be not able to solubilize mineral phosphate (Fig. 4). Nevertheless, the enhanced P uptake by plants can be also achieved via hormonal stimulation of root growth,



**Fig. 5** Isolates were tested for siderophore production on CAS medium. *E. coli* WA321 was used as a positive control. **A** CAS plates prepared with the MM9, pH 6.8. **B** CAS plates prepared with

the MM9, pH 6.0 and supplemented with casamino acid. *a E. coli* WA321, *b* strain WH120T, *c* strain 5B5 and *d* strain WH15

branching or root hair development mediated by IAA among others (Richardson and Simpson 2011).

On the other hand, the assay for siderophore production showed that all three strains react with the media (Fig. 5). Detection of siderophore production was carried out on chrome azurol S (CAS) agar plates. Removal of iron from the CAS dye by iron-chelating compounds results in a color change. The discoloration was observed for all strains with stronger discoloration zones observed around 5B5 and WH120T colonies than around colonies of strain WH15. Plants have high iron requirement but similarly to P, most of the iron in soil is in ferric form, which is unavailable for plant uptake (Hayat et al. 2010). The strategies of iron uptake by plants are similar to those from bacteria. Those include acidification of the rhizosphere resulting in reduction of  $\text{Fe}^3$ – $\text{Fe}^2$  or synthesis of  $\text{Fe}^3$  chelators (Morrissey and Gueriot 2009; Saha et al. 2013). Additional advantage of bacterial siderophore production is competition with pathogens by removing iron from the environment (Saha et al. 2013).

In order to test the ability of the strains to fix  $\text{N}_2$ , we have carried out PCR targeting nitrogenase (*nifH*) gene. However, the results showed no evidence of such ability of tested strains. The absence of *nifH* is in agreement with acidobacterial genome mining studies (Ward et al. 2009). Up to now, there is no experimental evidence for the ability of *Acidobacteria* type strains to fix nitrogen.

## Conclusion

Based on our findings, we provide for the first time a direct evidence of active *Acidobacteria*–plant interaction and data indicating growth-promoting effects by *Acidobacteria*. We verified that a possible auxin production is involved in plant growth promotion. Although commonly used, the test conducted to unravel possible mechanisms of this phenomenon, for the first time was applied for *Acidobacteria*. Further studies are needed to better understanding the beneficial *Acidobacteria*–plant interaction as well as the mechanisms involved in such interaction. In addition, we conclude that EPS production during root colonization by *Acidobacteria* might be helpful in root adhering to soil particles and in root protection. Taking into account the dominance in abundance of this phylum in soil environment, the overall impact of *Acidobacteria* on plant growth may be significant and the results shown here indicate for the first time that *Acidobacteria* can act as plant growth-promoting bacteria.

**Acknowledgments** We thank Manuela Di Lorenzo and Tanja R. Scheublin for technical advices. This work was financially supported by the Dutch Ministry of Economic Affairs, Agriculture and Innovation (729.004.016) and the BE-Basic partner organization ([www.be-basic.org](http://www.be-basic.org)). Publication 6101 of the Netherlands Institute of Ecology (NIOO-KNAW).

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Brankatschk R, Bodenhausen N, Zeyer J, Bürgmann H (2012) Simple absolute quantification method correcting for quantitative PCR efficiency variations for microbial community samples. *Appl Environ Microbiol* 78(12):4481–4489
- Bric JM, Bostock RM, Silverstone SE (1991) Rapid in situ assay for indoleacetic acid production by bacteria immobilized on a nitrocellulose membrane. *Appl Environ Microbiol* 57(2):535–538
- Chaparro JM, Badri DV, Vivanco JM (2014) Rhizosphere microbiome assemblage is affected by plant development. *ISME J* 8(4):790–803
- Chow ML, Radomski CC, McDermott JM, Davies J, Axelrood PE (2002) Molecular characterization of bacterial diversity in Lodgepole pine (*Pinus contorta*) rhizosphere soils from British Columbia forest soils differing in disturbance and geographic source. *FEMS Microbiol Ecol* 42(3):347–357
- da Rocha UN, van Elsas JD, van Overbeek LS (2010) Real-time PCR detection of *Holophagae* (*Acidobacteria*) and *Verrucomicrobia* subdivision 1 groups in bulk and leek (*Allium porrum*) rhizosphere soils. *J Microbiol Methods* 83(2):141–148
- da Rocha UN, Plugge CM, George I, van Elsas JD, van Overbeek LS (2013) The rhizosphere selects for particular groups of *Acidobacteria* and *Verrucomicrobia*. *PLoS One* 8(12):e82443
- Eichorst SA, Kuske CR, Schmidt TM (2011) Influence of plant polymers on the distribution and cultivation of bacteria in the phylum *Acidobacteria*. *Appl Environ Microbiol* 77(2):586–596
- Fierer N, Bradford MA, Jackson RB (2007) Toward an ecological classification of soil bacteria. *Ecology* 88(6):1354–1364
- Filion M, Hamelin R, Bernier L, St-Arnaud M (2004) Molecular profiling of rhizosphere microbial communities associated with healthy and diseased black spruce (*Picea mariana*) seedlings grown in a nursery. *Appl Environ Microbiol* 70(6):3541–3551
- Hayat R, Ali S, Amara U, Khalid R, Ahmed I (2010) Soil beneficial bacteria and their role in plant growth promotion: a review. *Ann Microbiol* 60(4):579–598
- Herder GD, Van Isterdael G, Beeckman T, De Smet I (2010) The roots of a new green revolution. *Trends Plant Sci* 15(11):600–607
- Idris R, Trifonova R, Puschenreiter M, Wenzel WW, Sessitsch A (2004) Bacterial communities associated with flowering plants of the Ni hyperaccumulator *Thlaspi goesingense*. *Appl Environ Microbiol* 70(5):2667–2677
- Janssen PH (2006) Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Appl Environ Microbiol* 72(3):1719–1728
- Kielak AM, Pijl AS, van Veen JA, Kowalchuk GA (2008) Differences in vegetation composition and plant species identity lead to only minor changes in soil-borne microbial communities in a former arable field. *FEMS Microbiol Ecol* 63(3):372–382
- Kielak AM, Barreto CC, Kowalchuk GA, van Veen JA, Kuramae EE (2016) The ecology of *Acidobacteria*: moving beyond genes and genomes. *Front Microbiol*. doi:10.3389/fmicb.2016.00744
- Koch IH, Gich F, Dunfield PF, Overmann J (2008) *Edaphobacter modestus* gen. nov., sp. nov., and *Edaphobacter aggregans* sp. nov., *Acidobacteria* isolated from alpine and forest soils. *Int J Syst Evol Microbiol* 58(5):1114–1122
- Kulichevskaya IS, Kostina LA, Valášková V, Rijkstra WIC, Damsté JSS, de Boer W, Dedys SN (2012) *Acidicapsa borealis* gen. nov., sp. nov. and *Acidicapsa ligni* sp. nov., subdivision 1 *Acidobacteria* from Sphagnum peat and decaying wood. *Int J Syst Evol Microbiol* 62(7):1512–1520
- Lee SH, Ka JO, Cho JC (2008) Members of the phylum *Acidobacteria* are dominant and metabolically active in rhizosphere soil. *FEMS Microbiol Lett* 285(2):263–269
- Mao Y, Li X, Smyth EM, Yannarell AC, Mackie RI (2014) Enrichment of specific bacterial and eukaryotic microbes in the rhizosphere of switchgrass (*Panicum virgatum* L.) through root exudates. *Environ Microbiol Rep* 6(3):293–306
- Mendes LW, Kuramae EE, Navarrete AA, van Veen JA, Tsai SM (2014) Taxonomical and functional microbial community selection in soybean rhizosphere. *ISME J* 8:1577–1587
- Morrissey J, Guerinot ML (2009) Iron uptake and transport in plants: the good, the bad, and the ionome. *Chem Rev* 109(10):4553–4567
- Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol Plant* 15(3):473–497
- Nautiyal CS (1999) An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. *FEMS Microbiol Lett* 170(1):265–270
- Nissinen RM, Männistö MK, Elsas JD (2012) Endophytic bacterial communities in three arctic plants from low arctic fell tundra are cold-adapted and host-plant specific. *FEMS Microbiol Ecol* 82(2):510–522
- Pankratov TA, Dedys SN (2010) *Granulicella paludicola* gen. nov., sp. nov., *Granulicella pectinivorans* sp. nov., *Granulicella aggregans* sp. nov. and *Granulicella rosea* sp. nov., acidophilic, polymer-degrading *Acidobacteria* from Sphagnum peat bogs. *Int J Syst Evol Microbiol* 60(12):2951–2959
- Pankratov TA, Serkebaeva YM, Kulichevskaya IS, Liesack W, Dedys SN (2008) Substrate-induced growth and isolation of *Acidobacteria* from acidic Sphagnum peat. *ISME J* 2(5):551–560
- Pankratov TA, Kirsanova LA, Kaparullina EN, Kevbrin VV, Dedys SN (2012) *Telmatobacter bradus* gen. nov., sp. nov., a cellulolytic facultative anaerobe from subdivision 1 of the *Acidobacteria*, and emended description of *Acidobacterium capsulatum* Kishimoto et al. 1991. *Int J Syst Evol Microbiol* 62(2):430–437
- Poosakkannu A, Nissinen R, Kytöviita MM (2015) Culturable endophytic microbial communities in the circumpolar grass, *Deschampsia flexuosa* in a sub-Arctic inland primary succession are habitat and growth stage specific. *Environ Microbiol Rep* 7(1):111–122
- Richardson AE, Simpson RJ (2011) Soil microorganisms mediating phosphorus availability update on microbial phosphorus. *Plant Physiol* 156(3):989–996
- Saha R, Saha N, Donofrio RS, Bestervelt LL (2013) Microbial siderophores: a mini review. *J Basic Microbiol* 53(4):303–317
- Schwyn B, Neilands J (1987) Universal chemical assay for the detection and determination of siderophores. *Anal Biochem* 160(1):47–56
- Shi S, Richardson AE, O’Callaghan M, DeAngelis KM, Jones EE, Stewart A, Firestone MK, Condon LM (2011) Effects of selected root exudate components on soil bacterial communities. *FEMS Microbiol Ecol* 77(3):600–610
- Singh BK, Munro S, Potts JM, Millard P (2007) Influence of grass species and soil type on rhizosphere microbial community structure in grassland soils. *Appl Soil Ecol* 36(2–3):147–155
- Valášková V, de Boer W, Gunnewiek PJK, Pospíšek M, Baldrian P (2009) Phylogenetic composition and properties of bacteria coexisting with the fungus *Hypholoma fasciculare* in decaying wood. *ISME J* 3(10):1218–1221

- Ward NL, Challacombe JF, Janssen PH, Henrissa B, Coutinho PM, Wu M, Xie G, Haft DH, Sait M, Badger J et al (2009) Three genomes from the phylum *Acidobacteria* provide insight into the lifestyles of these microorganisms in soils. *Appl Environ Microbiol* 75(7):2046–2056
- Whang K-S, Lee J-C, Lee H-R, Han S-I, Chung S-H (2014) *Terriglobus tenax* sp. nov., a novel exopolysaccharide producing acidobacterium isolated from rhizosphere soil of a medicinal plant. *Int J Syst Evolut Microbiol* 64(2):431–437