

1 **Fungus-associated bacteriome in charge of their host behaviour**

2

3 **Running title:** Fungal behaviour affected by its bacteriome

4

5 Kristin Schulz-Bohm^{1,3,§}, Olaf Tyc^{1,3,§}, Wietse de Boer^{1,3}, Nils Peereboom^{1,2}, Fons Debets²,
6 Niels Zaagman⁴, Thierry K.S. Janssens⁴ and Paolina Garbeva^{1*}

7

8 ¹ Department Microbial Ecology, Netherlands Institute of Ecology, NIOO-KNAW, PO Box 50,
9 6700 AB, Wageningen, Netherlands

10 ² Laboratory of Genetics, Wageningen University & Research Centre (WUR), PO Box 16, 6700
11 AA, Wageningen, Netherlands

12 ³ Department of Soil Quality, Wageningen University & Research Centre (WUR), PO Box 47,
13 6700 AA, Wageningen, Netherlands

14 ⁴MicroLife Solutions b.v., Science Park 406, 1098 XH Amsterdam, The Netherlands

15

16 * Correspondence: Paolina Garbeva, email: p.garbeva@nioo.knaw.nl

17

18 § These two authors contributed equally to this work

19 **Keywords:** *Mucor hiemalis*, fungal bacteriome, fungal behaviour, volatiles, interactions

20 **Abbreviations:** ARE (Artificial Root Exudates), CFU (Colony Forming Units), MA (antibiotic-
21 treated *M. hiemalis* isolate), M0 (original *Mucor hiemalis* isolate), OTU (Operational
22 Taxonomic Unit), PDA (Potato Dextrose Agar), TSB (Tryptic Soya Broth), VOCs (Volatile
23 Organic Compounds), WA (Water Agar), WYA (Water Yeast Agar)

24

25 **Abstract**

26 Bacterial-fungal interactions are widespread in nature and there is a growing number of
27 studies reporting distinct fungus-associated bacteria. However, little is known so far about how
28 shifts in the fungus-associated bacteriome will affect the fungal host's lifestyle. In the present
29 study, we describe for the first time the bacterial community associated with the saprotrophic
30 fungus *Mucor hiemalis*, commonly found in soil and rhizosphere. Two broad-spectrum
31 antibiotics that strongly altered the bacterial community associated with the fungus were
32 applied. Our results revealed that the antibiotic treatment did not significantly reduce the
33 amount of bacteria associated to the fungus but rather changed the community composition
34 by shifting from initially dominating *Alpha-Proteobacteria* to dominance of *Gamma-*
35 *Proteobacteria*. A novel approach was applied for the isolation of fungal-associated bacteria
36 which also revealed differences between bacterial isolates obtained from the original and the
37 antibiotic-treated *M. hiemalis*. The shift in the composition of the fungal-associated bacterial
38 community led to significantly reduced fungal growth, changes in fungal morphology, behavior
39 and secondary-metabolites production. Furthermore, our results showed that the antibiotic-
40 treated isolate was more attractive and susceptible to mycophagous bacteria as compared to
41 the original isolate. Overall, our study highlights the importance of the fungus-associated
42 bacteriome for the host's lifestyle and interactions and indicate that isolation with antibacterials
43 is not sufficient to eradicate the associated bacteria.

44

45 **1. Introduction**

46 Bacteria and fungi are widespread in nature and play important roles in many ecological
47 processes. Similar to other organisms, many fungi have an associated bacteriome and there
48 is a growing number of known endosymbionts where bacteria dwell within fungal hyphae
49 (Bonfante and Anca, 2009; Kobayashi and Crouch, 2009). Another type of fungal-bacterial
50 interaction is ectosymbiosis, where the bacterial partner is adhered to fungal hyphae
51 (Stopnisek et al., 2016; Warmink et al., 2009).

52 The symbioses with ecto- and endofungal bacteria are often overlooked, yet they may have a
53 profound effect on the fungus behavior and lifestyle. For example, in *Aspergillus nidulans* and
54 *A. niger*, the ectobacteria actinomycetes and *Bacillus subtilis* respectively, were shown to
55 affect fungal primary and secondary metabolism (Benoit et al., 2015; Schroeckh et al., 2009).

56 It is plausible that there is a relationship between fungi and bacteria of which both partner
57 profit. For example, bacteria or fungi can benefit from specific compounds that are produced
58 by the other partner if they cannot produce it themselves. Several mycorrhizal helper bacteria
59 secrete citric and malic acids that are metabolized by *Laccaria bicolor*, promoting its growth
60 (Duponnois and Garbaye, 1990). Conversely, ectomycorrhizal fungi may produce organic
61 acids or sugars that can affect the composition and growth of associated bacterial
62 communities. For example, the helper bacterial isolate *P. fluorescens* BBc6R8 can be
63 chemoattracted by the hyphae of the ectomycorrhizal fungus *L. bicolor* S238N and the
64 trehalose accumulated within the mycelium that promotes the growth of the helper bacterium
65 (Deveau et al., 2010). In the interaction between *S. cerevisiae* and several *Acinetobacter*
66 species, ethanol secreted by the yeast was shown to stimulate the growth of the bacterial
67 species and it can act as a signaling molecule, altering cell physiology (Smith et al., 2004).

68 Bacterial endosymbionts *Candidatus Glomeribacter gigasporarum* of arbuscular mycorrhizal
69 fungi *Gigaspora margarita* can be involved in the vitamin B12 provision for the fungus
70 (Ghignone et al., 2012). The co-adaptation between fungal hosts and bacterial endosymbionts

71 can become so tight that the genome size and the gene content of the endosymbionts is
72 reduced. Mycorrhizal endosymbionts from Mollicutes lineages were suggested to depend
73 metabolically on their host, and additionally have taken up regulatory eukaryotic genes
74 horizontally (Naito et al., 2015). Clear evidence for gene transfer between Mollicute-related
75 endobacteria and their mycorrhizal host *Dentiscutata heterogamma* was recently reported
76 (Torres-Cortes et al., 2015). Furthermore, endobacteria can improve the fitness of their host
77 by e.g. increasing the fungal sporulation success and raising the fungal bioenergetics capacity
78 (Salvioli et al., 2016). Another well-studied example is the rice seedling blight pathogen
79 *Rhizopus microsporus* (Lackner et al., 2009; Partida-Martinez and Hertweck, 2005). This
80 fungus contains endobacteria named *Burkholderia rhizoxinica* and *Burkholderia*
81 *endofungorum* (Partida-Martinez et al., 2007a), which have been shown to produce a potent
82 toxin involved in host pathogenesis (Gee et al., 2011; Partida-Martinez and Hertweck, 2005).
83 Interestingly, the endobacteria enforce their vertical transmission by controlling host
84 sporulation making use of a hrp type III secretion system (Lackner et al., 2011; Partida-
85 Martinez et al., 2007b). As a result, the host is not able to reproduce in absence of its
86 endosymbiont, thereby ensuring maintenance of the symbiosis (Partida-Martinez et al.,
87 2007b).

88 Microscopic and molecular analysis showed that several nitrous oxide-producing fungal
89 isolates of *Mortierella elongata* harboured endobacteria in their mycelia (Sato et al., 2010).
90 The sequencing of 16S rRNA genes revealed that the N₂O-producing fungus *Mortierella*
91 *elongate* harbored endobacteria belonging to the family *Burkholderiaceae*, however, the
92 significance of this fungal-bacterial association is unknown (Sato et al., 2010). Recently, an
93 endophyphal bacterium living in association with *Mortierella elongate* was isolated and on the
94 basis of phenotypic, chemotaxonomic and phylogenetic characteristics it was identified as a
95 novel genus and species, for which the name *Mycoavidus cysteinexigens* gen. nov., sp., nov
96 was proposed (Ohshima et al., 2016).

97 It is a common practice prior performing experiments with soil-borne fungi to pre-culture them
98 on a media supplemented with antibiotics (Ballhausen et al., 2016; Singh et al., 2015). In our
99 lab working with the saprotrophic fungus *Mucor hiemalis* (Zygomycota) isolated from the
100 rhizosphere of *Carex arenaria* (sand sedge) (De Rooij-van der Goes et al., 1995), we observed
101 that treatment with broad-spectrum antibiotics strongly altered the fungal morphology and
102 hyphal extension. Based on this observation, we aimed first to determine the bacterial
103 community associated with *Mucor hiemalis* and to test if the antibiotic treatment resulted in
104 bacteria-free fungus. Furthermore, we aimed to reveal how changes in the bacterial
105 community affect fungal fitness, behavior, metabolites production and interactions.

106

107 **2. Material and Methods**

108 **2.1 Fungal strains and growth conditions**

109 The *Mucor hiemalis* isolate M0 obtained from the rhizosphere of *Carex arenaria* (Sand sedge)
110 collected from sandy dune soil in the Netherlands was originally isolated on malt extract agar
111 supplemented with 50 ppm oxytetracycline (De Rooij-van der Goes et al., 1995). Here, the
112 antibiotic-treated isolate MA derived from the isolate M0, after plating on King's B agar with
113 antibiotics (rifampicin and kanamycin 50 mg/ml, final concentration), and transferred to Water
114 Yeast Agar (WYA) (Garbeva et al., 2011) and Oatmeal agar OA (24 g/L Difco, France)
115 supplemented with rifampicin and kanamycin (50 mg/ml final concentration). The spores were
116 washed in rifampicin and kanamycin solution (50 mg/ml final concentration) and collected over
117 glass wool with sterile deionized water and stored at -80°C. Spores of the M0 and MA isolates
118 were grown on nutrient-rich 0.5 strength Potato dextrose agar (PDA, Oxoid, England; pH 6),
119 and nutrient-poor Water agar + (NH₄)₂SO₄, pH 6.7 (Garbeva et al., 2011).

120

121 **2.2 Fungal identification**

122 The identity of the isolate M0 and MA was confirmed using the ITS sequences targeted by the
123 primers ITS1 and ITS4 (White et al., 1990). Fungal DNA was extracted using the ZR
124 Fungal/Bacterial DNA MiniPrep (Zymo Research, USA) according to the manufacturer's
125 instructions. Extracted DNA was used for PCR amplification with a master mix containing 1x
126 FastStart High Fidelity Reaction Buffer (Roche) with 18 mM MgCl₂ (Roche), 0.04 U FastStart
127 High Fidelity Enzyme Blend (Roche), 200 µM of each dNTP, 0.6 µM ITS1 and ITS4 primer.
128 The thermal protocol was as follows: initial denaturation at 95°C for 5 minutes, and 35 cycles
129 of 95°C for 30 seconds, 55°C for 30 seconds, 72°C for 1 minute, and a final elongation at 72°C
130 for 10 minutes. The PCR product was cleaned using a PCR purification kit (QIAGEN Benelux
131 B.V., The Netherlands) and sent for Sanger sequencing to MacroGen Europe (Amsterdam,

132 The Netherlands). Obtained sequences were checked for quality using BioEdit (Hall, 1999)
133 and identified using NCBI nucleotide database BLASTn (Altschul et al., 1990).

134

135 **2.3 Hyphal extension of fungal isolates M0 and MA**

136 For measuring hyphal extension, 6-mm-diameter agar disks taken from the edge of the fungal
137 hyphae of M0 or MA (pre-grown on 0.5 strength PDA) were plated in the middle of nutrient-
138 rich 0.5 strength PDA or nutrient-poor WA plates (8.5 cm diameter). Per fungal isolate and
139 agar type (i.e. PDA and WA) three plates were set-up. The plates were sealed with parafilm
140 and incubated at 20 °C for 5 days. On the fifth day, for each plate the extension of fungal
141 hyphae was measured with a ruler in three coordinates and the average extension was
142 calculates.

143

144 **2.4 Competition assay of fungal isolates M0 and MA**

145 Two plugs taken from the edge of the (on 0.5 strength PDA) pre-grown fungal hyphae of M0
146 or MA were placed with 3 cm distance from each other on a new 0.5 strength PDA plate (8.5
147 cm diameter) and incubated at 20°C. As control, both fungal isolates were incubated
148 separately on 0.5 strength PDA at the same time. For each treatment three replicates were
149 set-up. The hyphal extension was monitored at day 7, 14 and 21 after inoculation. Pictures
150 were taken with a Panasonic DMC-FZ200 digital camera. The area of M0 and MA hyphal
151 extension was measured with AxioVision V 4.9.1.0 (Carl Zeiss Microscopy GmbH, Germany).
152 A relative area for the hyphal extension per replicate was calculated by dividing the area
153 obtained in the competition experiment per fungal isolate by the average area obtained for
154 each control.

155

156 **2.5 Bioassay for testing the fungal growth inhibition by bacterial secondary** 157 **metabolites**

158 The bacterial strains *Burkholderia sp.* AD024 (De Ridder-Duine et al., 2005, Schulz-Bohm et
159 al., 2015) and *Collimonas fungivorans* Ter 331 (De Boer et al., 2004) were incubated overnight
160 at 22°C in 0.1 Tryptic Soya Broth (TSB) (Tyc et al., 2014), washed with phosphate buffer
161 (10mM KH₂PO₄, pH 6.5), and the OD₆₀₀ was adjusted to 10⁸ cells per ml. 50 µl of the bacterial
162 suspension was spread horizontally in the middle of a Petri-dish containing 1.5 % (v/v) water
163 agar (Schmidt et al., 2015) supplied with artificial root exudates (WA + ARE) as described
164 previously by Schulz-Bohm et al. (2015). Per liter WA, 18 ml ARE stock solution was added,
165 implying that 134 µg carbon per ml agar was additionally supplied. Bacteria were incubated
166 for three days at 20°C and a plug of 8 mm diameter with fungal hyphae (pre-grown on 0.5
167 strength PDA) was transferred to the top of the Petri-dish. Agar plates were wrapped with two
168 layer of parafilm and incubated at 20°C. After four days, the distances between the edge of
169 the fungal hyphae and bacterial biofilm (refer to as inhibition zone) were measured. Pictures
170 of agar plates were taken with a Panasonic DMC-FZ200 digital camera. Pictures of fungal
171 hyphae were taken and processed with a Olympus SZX12-ILLK200 stereo microscope
172 (Olympus Cooperations, Japan) in combination with AxioVision V 4.9.1.0 (Carl Zeiss
173 Microscopy GmbH, Germany).

174

175 **2.6 Bacterial and yeast isolation and identification**

176 Mycelium of M0 and MA (pre-grown on 0.5 strength PDA) was collected in 7 ml sterile
177 phosphate buffer (10 mM KH₂PO₄, pH 6.5). The mycelium was sheared using a MICRA D-9
178 homogenizer (MICCRA GmbH, Germany) for 5 minutes. The resulting suspension was
179 shaken for 45 minutes at 22°C and filtrated using membrane pore sizes of 10µM (Millipore,
180 The Netherlands) as well as 3µM (Schleicher and Schuell, Germany). 350 µL aliquots of the
181 filtrate were plated on R2A medium (Difco, France), WYA and 0.1 TSB (Tyc et al., 2014)
182 containing 100 mg/L filter sterilized cycloheximide (Sigma-Aldrich, The Netherlands) and 50
183 mg/L thiabendazole (Sigma-Aldrich, The Netherlands) to inhibit the growth of M0 and MA.
184 Three volatile compounds: dimethyldisulfide, dimethyltrisulfide and benzonitrile (Sigma-

185 Aldrich, The Netherlands) with known antifungal but no antibacterial activity (Garbeva et al.,
186 2014a), were added in concentrations of 160 μmol to a sterile filter paper positioned at the
187 edge of the plate as described previously by Garbeva et al. (2014a). Plates supplemented
188 with volatiles were kept in a protective cabinet at room temperature ($\pm 20^\circ\text{C}$).

189 PCR amplification of 16S rRNA genes from the isolates was either performed directly with
190 colony material diluted in sterile deionized water or with isolated DNA using the QIAamp DNA
191 Mini Kit (Qiagen, The Netherlands) according to the manufacturer's protocol. The PCR mix
192 contained 0.17 mg/ml BSA, 0.33 μM of each primer (27F or 515F and 1492r; Lane, 1991), 1.5-
193 2 μl template, and 1x PCR Mastermix (Thermo Scientific, The Netherlands) containing
194 0.05U/ μl Taq DNA polymerase, reaction buffer, 4 mM MgCl_2 , and 0.4 mM of each dNTP. The
195 thermal protocol was as followed: initial denaturation at 95°C for 8 min, 5 cycles at 95°C for
196 60 sec, at 40°C for 60 sec, and at 72°C for 90 sec, and 35 subsequent cycles at 95°C for 60
197 sec, at 53°C for 30 sec, and 72°C for 70 sec. The final elongation was at 72°C for 5 min. The
198 16S rRNA PCR product of each (bacterial) isolate was purified and sent for sequencing to
199 LCG genomics (Berlin, Germany). Sequences were identified using NCBI nucleotide database
200 and BLASTn algorithm (Altschul et al., 1990). Microscopic pictures to describe the cell
201 morphology were taken and processed with Axio Imager M1 (Carl Zeiss Microscopy GmbH,
202 Germany) in combination with AxioVision V 4.9.1.0 (Carl Zeiss Microscopy GmbH, Germany).
203 All isolates were tested on the resistance against kanamycin and rifampicin by plating them
204 on 0.1 TSB containing 50 $\mu\text{g/ml}$ (final concentration) of each antibiotic. The agar plates were
205 incubated for at least three weeks at 20°C and regularly checked for potential growth.

206

207 **2.7 DNA extraction from fungal hyphae**

208 Hyphae (pre-grown on 0.5 strength PDA) were transferred into Lysin-Matrix E tubes
209 (Biomedicals, The Netherlands), weighted (refer to as fresh-weight), and freeze-dried in liquid
210 nitrogen. Subsequently, fungal-bacterial DNA was extracted with phenol-chloroform as

211 described by Schulz-Bohm et al. (2015). Nucleic acid extracts were stored at -80°C or were
212 immediately used for PCR amplification.

213

214 **2.8 Quantitative PCR of bacterial *rpoB* genes**

215 Primers to quantify the gene of the β subunit of the bacterial RNA polymerase (*rpoB*) were
216 designed based on an alignment of 118 *rpoB* sequences of phylogenetic different bacteria
217 available at NCBI (latest visit: 30/06/15, Table S3). The designed primers RpoB-fw1 (5'-
218 GAAGGTCCGAACATCGGTCT-3') and RpoB-r1 (5'-TGCATGTTTCGAGCCCATCA-3') amplify
219 a fragment of 370 bp in the conserved region of *rpoB*. Quantitative PCR (qPCR) of the *rpoB*
220 gene was performed with a BioRad C1000 Touch™ Thermal Cycler (Bio-Rad, The
221 Netherlands) whereas each template was quantified in triplicates resulting in a total of nine
222 reactions per fungal isolate (3 technical x 3 biological replicates). The 20- μ l reaction mixture
223 consisted of 1-fold SensiFAST™ SYBR® No-ROX Kit (Bioline GmbH, The Netherlands), BSA
224 (0.5 μ g μ l⁻¹), 500 nM RpoB-fw1, 500 nM RpoB-r1, and 5 μ l of diluted template DNA (2-5 ng/
225 μ l) or nuclease-free water in case of the negative control. Conditions for the quantification
226 were as follows: 5 min initial denaturation at 95°C, ensued by 40 cycles of denaturation for
227 30 sec at 95°C, annealing for 30 sec at 63°C, elongation for 30 sec at 72°C, and fluorescence
228 signal detection for 15 sec at 82°C. Immediately after the 40th PCR cycle, a melting curve
229 analyses (Fig. S5) from 63°C to 95°C with increments of 0.5°C was followed. Agarose gel
230 electrophoreses of qPCR products displayed single bands of the expected size. Furthermore,
231 sequencing of cloned *rpoB* qPCR products revealed that solely bacterial *rpoB* genes were
232 targeted by RpoB-fw1/RpoB-r1 despite a huge background of fungal DNA (Fig. S2). Gene
233 copy numbers were calculated according to a standard curve (Fig. S4) and corrected for
234 potential inhibition (Zaprasis et al., 2010). A standard curve was set up by serially diluting of a
235 pGEM-T vector containing an *rpoB* gene fragment of the strain *Burkholderia sp.* AD024.

236

237 **2.9 Cloning of *rpoB* genes and sequence analysis**

238 The *rpoB* qPCR products obtained for DNA extracted from the original fungal isolate M0 or
239 the antibiotic-treated fungal isolate MA were pooled and purified with 20 % polyethylene glycol
240 6000 (AppliChem, Germany) and 2.5M sodium chloride after following protocol. One volume
241 of PCR product was mixed with one volume of the polyethylene glycol solution, incubated for
242 15 min at 37°C, and centrifuged for 10 min at 10,000 x g. The resulting pellet was washed with
243 70 % ethanol and resuspended in DNase/RNase-free water. Purified qPCR products were
244 ligated into a pGEM-T vector plasmid (Promega, USA) and transformed in *Escherichia coli*
245 JM109 competent cells (Promega, USA) according to the manual instructions. Clones were
246 picked and checked for the correct insert by M13 PCR (Messing, 1993). In total 70 M13 PCR
247 products with the correct length (35 per fungal isolate) were purified and sequenced by LCG
248 genomics (Berlin, Germany).

249 Sequences were analyzed with MEGA 6 (Tamura et al., 2011) and BLASTn. Based on a
250 pairwise distance matrix of aligned *rpoB* nucleotide sequences (Kimura 2-parameter model,
251 substitution includes transitions and transversions, pairwise deletion for gaps/missing data
252 treatment), sequences were assigned to different Operational Taxonomic Units (OTUs) and
253 the diversity index CHAO was calculated with DOTUR (Schloss and Handelsman, 2005). A
254 threshold value of 97 % was used to define species-level OTUs (Adékambi et al., 2009). The
255 coverage of the gene libraries was calculated according to Schloss et al. (2004) and
256 rarefaction curves were constructed after the method of Hurlbert (Heck et al., 1975; Hurlbert,
257 1971).

258

259 **2.10 Phylogenetic analyses**

260 All phylogenetic trees were constructed with MEGA 6. The algorithms Maximum-Likelihood
261 (Kimura 2-parameter model, partial deletion, 10,000 bootstraps), Neighbour-Joining (Saitou
262 and Nei, 1987; Kimura 2-parameter model, complete deletion, 1000 bootstraps), and

263 Maximum-Parsimony (Subtree-pruning-regrafting search method, complete deletion, 1000
264 bootstraps) were applied. Trees for 16S rRNA gene sequences from bacterial isolates and
265 *rpoB* clone sequences were based on an alignment of 800 and 398 nucleotide positions,
266 respectively.

267

268 **2.11 Nucleotide sequence accession numbers**

269 Sequences were submitted to GenBank. Accession numbers: KX057404 - KX057471 (*rpoB*
270 clones sequences) and KX057472 - KX057478 (16S rRNA gene sequences of bacterial and
271 yeast isolates associated to M0 and MA).

272

273 **2.12 Volatile trapping and measurement**

274 For the collection of volatiles, glass petri dishes were used with lids with an exit to which a
275 steel trap with 150 mg Tenax TA and 150 mg Carbopack B (Markes International Ltd, United
276 Kingdom) was fixed (Garbeva et al., 2014b). Volatiles were collected for 20 hours at days 3
277 and 6. The traps were closed and stored at 4°C until analysis. Incubations were done in
278 triplicates, including medium controls.

279 Trapped volatiles were desorbed using an automated thermodesorption unit (model UnityTD-
280 100, Markes International Ltd., United Kingdom) at 210°C for 12 min (Helium flow 50 ml/min)
281 and trapped on a cold trap at -10°C. The trapped volatiles were introduced into the GC-QTOF
282 (model Agilent 7890B GC and the Agilent 7200A QTOF, USA) by heating the cold trap for 3
283 min to 280°C. Split ratio was set to 1:20, and the column used was a 30 × 0.25 mm ID RXI-
284 5MS, film thickness 0.25 µm (Restek 13424-6850, USA). Temperature program used was as
285 follows: 39°C for 2 min, from 39 to 95°C at 3,5 °C/min, then to 165°C at 6°C/min, to 250°C at
286 15°C/min and finally to 300°C at 40°C/min, hold 20 min. The VOCs were detected by the MS
287 operating at 70 eV in EI mode. Mass-spectra were extracted with MassHunter Qualitative
288 Analysis Software V B.06.00 Build 6.0.633.0 (Agilent Technologies, USA) using the GC-Q-

289 TOF qualitative analysis module. The obtained mass spectra were exported as mzData files
290 for further processing in MZmine. The files were imported to MZmine V2.14.2 (Pluskal et al.,
291 2010) and compounds were identified via their mass spectra using deconvolution function
292 (Local-Maximum algorithm) in combination with two mass-spectral-libraries: NIST 2014 V2.20
293 (National Institute of Standards and Technology, USA <http://www.nist.gov>) and Wiley 9th
294 edition mass spectral libraries and by their linear retention indexes (LRI). The LRI values were
295 calculated using an alkane calibration mix before the measurements in combination with
296 AMDIS 2.72 (National Institute of Standards and Technology, USA). The calculated LRI were
297 compared with those found in the NIST and in the in-house NIOO-KNAW LRI database. After
298 deconvolution and mass identification peak lists containing the mass features of each
299 treatment (MZ-value/Retention time and the peak intensity) were created and exported as
300 CSV files for statistical processing via MetaboAnalyst V3.0 (www.metaboanalyst.ca; Xia et al.,
301 2015; Xia et al., 2012).

302

303 **2.13 Test on the effect of fungal VOCs on bacterial growth and motility**

304 A double plate-within-a-plate system (Schmidt et al., 2015b) was used to test the effect of
305 fungal volatiles emitted by the M0 and MA isolates on bacterial growth and motility. A 6 mm
306 plug with fungal hyphae (pre-grown on 0.5 strength PDA) was transferred on 1.5 % WA or 0.5
307 strength PDA (Schmidt et al., 2015b) on a small Petri-dish (35 mm diameter) located in one
308 half of a bipartite Petri-dish (Fig. S3). The other half of the bipartite Petri-dish was filled with
309 12 ml 1.5 % WA + ARE (described above) or with 0.3 % WA + ARE for the motility assay. After
310 the fungus was incubated for three days at 20°C, 10 µl droplets of bacterial suspensions of
311 different cell concentration (10^3 to 10^6 CFU per ml) were added on WA + ARE, i.e. per replicate
312 four droplets (from lower to higher cell concentration) were placed horizontally to the border
313 of the two compartments (Fig. S3A). In case of the motility assay, one droplet of 10 µl bacterial
314 suspension (10^7 CFU per ml) was placed in the middle of the second compartment filled with
315 0.3 % motility agar (Fig. S3B). The bacterial suspension consisted either of *Burkholderia sp.*

316 AD024 or *Collimonas fungivorans* Ter331 which were grown overnight at 22°C in 0.1 TSB and
317 washed with phosphate buffer (10mM KH₂PO₄, pH 6.5). Bipartite Petri-dishes were sealed
318 two-times with parafilm and incubated for six days at 20°C.

319 Colonies at the spot where the lowest dilution of bacterial cells per ml was added were used
320 for the analysis of bacterial growth. In case of bacterial motility, the diameter of the bacterial
321 colony was measured at four different spots (Fig. S3B) and a mean value was calculated (refer
322 to as colony extension) for further analyses.

323

324 **2.14 Statistical analysis**

325 Statistical analysis on volatolomic data was performed using MetaboAnalyst V3.0 (Xia et al.,
326 2012; Xia et al., 2015). To identify significant mass features, One-way-ANOVA with post-hoc
327 Tukey test (HSD- test) were performed. To identify important mass features, PLS-D analysis
328 was performed. Mass features were considered to be statistical relevant if p- and FDR- values
329 were ≤ 0.05 . Statistical relevant mass features were further used for compound identification.

330 The statistical analysis on hyphal extension of *M. hiemalis* as well as on the number of
331 bacterial colonies and on colony extension when they were grown in presence of fungal VOCs
332 was performed with R 3.1.1 (<http://www.r-project.org/>) using One-way-ANOVA Turkey's HSD
333 test (De Mendiburu, 2014). Student's t-tests on data obtained by qPCR as well as for
334 bioassays on fungal growth competition and on fungal growth inhibition by bacterial secondary
335 metabolites were conducted with SigmaPlot 12.5 (Systat Software). The 5% level was taken
336 as threshold for significance.

337 3. Results

338 3.1 Composition and abundance of the bacterial community associated with the fungus

339 *M. hiemalis* based on *rpoB* gene

340 The composition of bacterial community associated with the fungus *M. hiemalis* was
341 determined by targeting the gene of the β subunit of the bacterial RNA polymerase (*rpoB*). For
342 this purpose, primers were newly designed with the aim to amplify a broad spectrum of
343 phylogenetic different bacteria but not fungal DNA. In total 68 *rpoB* clone sequences were
344 retrieved from both the original M0 isolated and antibiotic-treated MA isolate (Fig. 1A and Fig.
345 S2). The total coverage was 94 % and rarefaction analysis showed an out-plateauing curve
346 (Fig. 1B) indicating that sequencing was sufficient for the coverage of fungus-associated
347 bacterial species. For M0, the coverage of *rpoB* sequences was lower than for MA (91 % and
348 97 %, respectively). This is reflected by a stronger out-plateauing rarefaction curve (Fig. 1C)
349 and a lower CHAO diversity index for MA compared to M0 (6 and 19, respectively). In general,
350 *rpoB* clone sequences could be assigned to the phyla *Actinobacteria*, *Bacteroidetes*, and
351 *Proteobacteria* (classes: *Alpha*-, *Beta*-, and *Gamma-Proteobacteria*) (Fig. S2). Clone
352 sequences retrieved from M0 were affiliated to *Actinobacteria* (3 %) and *Proteobacteria* (97 %,
353 dominated by *Alpha-Proteobacteria* [53 %], followed by *Gamma*- and *Beta-Proteobacteria* [29
354 % and 15 %, respectively]). *RpoB* clone sequences from MA were affiliated with *Bacteroidetes*
355 (3%), and *Proteobacteria* (97 %, dominated by *Gamma-Proteobacteria* [65 %], followed by
356 *Alpha*- and *Beta-Proteobacteria* [24 % and 9 %, respectively]). Within the *rpoB* clone library of
357 M0, most sequences could be assigned to *Bradyrhizobium oligotrophicum* (41 %) and for MA,
358 sequences were mostly related to *Serratia marcescens subsp. sakuensis* (47 %) (Fig. 1A).

359 In total, the bacterial community between M0 and MA appeared to be distinct. More diverse
360 bacteria at the species-level were associated to M0. The bacterial community of the original
361 M0 isolate was dominated by *Alpha-Proteobacteria* (*Bradyrhizobium* and *Brevundimonas*), the
362 community associated to the antibiotic-treated isolate MA appeared to be dominated by
363 *Gamma-Proteobacteria* (*Serratia*, *Pseudomonas* and *Rhodanobacter*) (Fig. 1A). After the

364 antibiotic treatment sequences of *Mycobacterium* spp., *Stenotrophomonas maltophilia*,
365 *Bradyrhizobium oligotrophicum* and *Bradyrhizobium liaoningense* were not detected.

366 The PCR quantification revealed that the *rpoB* gene copy number per gram fresh fungal
367 hyphae for M0 ($3.47 \times 10^4 \pm 1.09 \times 10^4$) was about 1.4 fold higher compared to MA (2.64×10^4
368 $\pm 8.53 \times 10^3$). However, the difference was not significant, indicating that the treatment with
369 broad-spectrum antibiotics reduced the number of associated bacteria to the fungus *M.*
370 *hiemalis* only slightly.

371

372 **3.2 Isolation of *M. hiemalis* associated bacteria and yeast**

373 Fungus-associated bacteria were isolated from both the original M0 and the antibiotic-treated
374 MA isolate by using a combination of three different volatile organic compounds (VOCs) with
375 known antifungal activity (Garbeva et al., 2014a). Only by applying a combination of all three
376 volatiles the growth of *Mucor* hyphae could be successfully inhibited while the growth of
377 fungus-associated bacteria was enhanced. In total, seven isolate types with different colony
378 and cell morphologies were obtained (Fig. 2). Sequences of those isolates were related to
379 different bacteria and to one yeast. Isolates retrieved from M0 were related to the *Alpha-*
380 *Proteobacterium Ochrobactrum intermedium* strain AG1 (100% identity), the *Gamma-*
381 *Proteobacterium Stenotrophomonas maltophilia* strain Zunyi-F (99 % identity), and the
382 *Firmicute Bacillus* sp. A-3-15 (100 % identity). Most isolates were related to *S. maltophilia*
383 (Fig. 3). Sequences related to this species as well as sequences related to *Alpha-*
384 *Proteobacteria* were also found for the *rpoB* clone library of M0 (Fig. 1 and Fig. S1). Isolates
385 obtained from MA were related to the *Actinobacterium Kocuria kristinae* strain VTT E-82147
386 (99 % identity) and *Micrococcus* sp. O-1 (99 % ident), the *Firmicute Staphylococcus* sp. ccc_1
387 (99 % identity) as well as to the yeast *Meyerozyma guilliermondii* strain Nc49HB-1 (99 %
388 identity). Sequences related to *Staphylococcus* and *Kocuria* were also exclusively detected
389 for MA by next generation sequencing of 16S rRNA (Table S2) but not by cloning of *rpoB*
390 qPCR products. All isolates were tested on resistance to rifampicin and kanamycin –

391 antibiotics previously applied to *M. hiemalis*. Only *S. maltophila* and the yeast *Meyerozyma*
392 *guilliermondii* (obtained from M0 and MA, respectively) grew on TSB with rifampicin and
393 kanamycin (Fig. 2).

394

395 **3.3 Comparison of the hyphal extension**

396 The comparison of hyphal extension revealed differences between the original M0 isolate and
397 the antibiotic-treated MA isolate on both media, with significantly higher hyphal extension of
398 M0 (Fig. 3). Besides hyphal extension, morphological differences in pigmentation were
399 observed between the two isolates under both nutrient conditions (Fig.S 4 and 5). On the
400 nutrient-rich media, M0 had yellow pigmentation and MA had brown pigmentation (Fig. 4).

401

402 **3.4 Competition between the fungal isolates M0 and MA**

403 By incubating both fungal isolates M0 and MA together in Petri-dishes, the area of hyphal
404 extension as compared to the monoculture was significantly lower for MA (57 ± 6 % [day7],
405 35 ± 4 % [day14]; n= 3) than for M0 (74 ± 4 % [day7], 65 ± 3 % [day14]; n = 3). After seven
406 days of incubation, the hyphal extension of both competing isolates did not further increase.
407 Only the hyphal network became denser, and a clear separation zone between both fungal
408 isolates was visible (Fig. 4). For the control (i.e. the monocultures of both fungal isolates), the
409 Petri-dishes were fully covered with mycelium after 14 days of incubation (Fig. 4).

410

411 **3.5 Growth of M0 and MA in presence of rhizobacteria**

412 In the presence of the rhizobacterium *Burkholderia sp.* AD024, the hyphal extension of MA
413 towards the bacterial biofilm was more inhibited as compared to M0 (Fig. 5C and I). The
414 inhibition zone, i.e. the distance between bacterial biofilm and the fungal hyphae, was
415 significantly bigger for MA (73 ± 6 mm) than for M0 (43 ± 12 mm).

416 In case of *Collimonas fungivorans* Ter 331, there was no significant difference between M0
417 and MA. However, the fungal hyphae of MA were visibly affected by the presence of
418 *Collimonas* compared to M0 (Fig. 5B, F, H, and L). Such difference was not observed for
419 incubations with *Burkholderia sp.* AD024 (Fig. 5B, D, H, and J).

420

421 **3.6 Volatiles produced by *M. hiemalis* isolate M0 and MA**

422 Headspace volatiles produced by *M. hiemalis* M0 and MA on nutrient-rich PDA and nutrient-
423 poor media WA were collected at day 6 after inoculation and subsequently analysed using
424 GC/MS. Different blends of volatiles were produced by the two isolates under both nutrient
425 conditions (Fig. 6) with more volatile compounds produced on the nutrient-rich PDA media.
426 Interestingly the volatile blend of the MA isolate on nutrient-poor WA media did not differ from
427 the control without fungus (Fig. 6). Several compounds, such as trifluorobenzene, 1 butanole
428 and 1-butanol-2methyl, were detected only in the headspace of M0 isolate, while heptane-2,4-
429 dimethyl was produced only by the MA isolate (Table 1). Numerous compounds could not be
430 assigned with certainty to a volatile organic compound and remained unknown. Some of these
431 unknown compounds were also discriminative between M0 and MA.

432

433 **3.7 Effect of fungal volatiles on rhizobacteria**

434 By exposure to fungal volatiles emitted by MA on PDA, the numbers of colonies of
435 rhizobacteria *Burkholderia sp.* AD024 and *Collimonas fungivorans* Ter 331 were significantly
436 increased in comparison to M0. Such significant difference was not observed when the fungal
437 isolates were incubated on WA (Fig. 7A and B). In addition, the number of *C. fungivorans* Ter
438 331 colonies was significantly reduced by exposure to volatiles of M0 growing on PDA (Fig.
439 7B).

440 The motility of both bacteria (reflected in colony extension) was differently affected by the
441 volatiles of M0 and MA (Fig. 7C and D). The swimming motility of *C. fungivorans* Ter 331 was

442 significantly higher in presence of volatiles produced by MA growing on PDA as compared to
443 M0 (Fig. 7D).

444 The swimming motility of the bacterial strain *Burkholderia* sp. AD24 was significantly increased
445 by exposure to volatiles of both fungal isolates whereas volatiles of M0 seemed to stimulate
446 more the motility of this strain compared to volatiles of MA (Fig. 7C).

447

448 **4. Discussion**

449 The fates of bacteria and fungi are ecologically intimately connected in soil and rhizosphere.
450 Close association of bacteria with fungi, both endo- and ectosymbionts is well known (Frey-
451 Klett et al., 2011). Most methods for fungal isolation and purification are involving application
452 of broad-spectrum antibiotics (Singh et al., 2015; Oliveira et al., 2013). The model soil-borne
453 fungus *Mucor hiemalis* used in our study was originally isolated by plating on media
454 supplemented with oxytetracycline (De Rooij-van der Goes et al., 1995). To ensure that we
455 are using bacteria-free fungus, spores of *Mucor hiemalis* were washed with antibiotics and
456 plated on a media supplemented with rifampicin and kanamycin. The treatment with these
457 broad-spectrum antibiotics strongly altered the fungal morphology and hyphal extension.
458 Hence, the purpose of our study was to determine the bacterial community associated with
459 the fungus *M. hiemalis* and to examine the effect of the two broad-spectrum antibiotics on the
460 fungus-associated bacteriome and, consequently, on fungal behavior and interactions.

461 Our results revealed that a high diversity of bacteria is associated to *M. hiemalis*. Interestingly,
462 the antibiotic treatment did not significantly reduce the amount of bacteria associated to the
463 fungus but rather changed the community composition by shifting from initially dominating
464 *Alpha-Proteobacteria* to dominance by *Gamma-Proteobacteria*. The antibiotic-treated *M.*
465 *hiemalis* isolate revealed less diverse bacterial community as compared to the original isolate.
466 Interestingly, several OTUs (*Rhodanobacter* spp., *Pseudomonas* sp., and *Bacteroides* spp)
467 not detected in the original M0 isolate were detected in the antibiotic treated MA isolates. It is
468 plausible that by affecting some of the antibiotic sensitive strains other “silent” bacteria carried
469 by the fungus became more dominant. The composition of the non-culturable bacterial
470 community associated with the fungus *M. hiemalis* is determined based on housekeeping *rpoB*
471 *gene* which allowed to target bacteria and, in addition, to avoid co-amplification of fungal DNA.
472 Our attempt to determine the composition of the bacterial community based on 16S rRNA next
473 generation sequencing demonstrated also differences between the bacterial communities

474 (supplementary material Table S2 and Fig. S1). However, the majority (more than 90%) of
475 OTUs were assigned to fungal mitochondrial DNA (Fig. S1A and B).

476 Plausible explanation for bacterial survival after antibiotic treatment is the resistance to the
477 applied antibiotic or protection by the fungal host. This is in line with another study that failed
478 to obtain bacteria-free fungi (Sharma et al., 2008). Analogous, it is also challenging to isolate
479 bacteria that are free from fungal mycelium as often both partners are dependent on each
480 other (Sato et al., 2010). Here, we applied a novel method for the isolation of fungal-associated
481 bacteria by the application of three volatile compounds (namely benzonitril, dimethyldisulfide
482 and dimethyltrisulfide) with strong antifungal activity and stimulating effect on bacterial growth
483 (Garbeva et al., 2014a). The high fungal sensitivity to volatile compounds (Effmert et al., 2012;
484 Garbeva et al., 2014a) compared to high resistance or even growth promotion for bacteria
485 (Garbeva et al., 2014b; Schmid et al., 2015a) make volatiles suitable compounds for isolation
486 of fungal-associated bacteria. However, difference in fungal susceptibility to volatiles was
487 observed and reported in several independent studies (Effmert et al., 2012 Schmid et al.,
488 2015). The yeast *Meyerozyma guilliermondii* was not affected by the volatile treatment
489 indicating that yeast may be less susceptible to volatiles. The results of the bacterial isolation
490 likewise revealed differences in the communities between the original and antibiotic-treated
491 *M. hiemalis*. While isolates obtained from the original M0 isolate belonged to *Alpha*- and
492 *Gamma-Proteobacterium* and *Firmicutes*, the isolates obtained from the antibiotic-treated MA
493 belonged to the phyla *Actinobacteria*, *Firmicutes* and yeast. Interestingly, the test for
494 resistance to antibiotics revealed that only *Stenotrophomonas* and the yeast were resistant to
495 rifampicin and kanamycin, indicating that the fungal host may indeed play a major role in
496 protecting their associated bacteria. Several sequences related to the bacterial isolates such
497 as for *Stenotrophomonas*, *Staphylococcus* and *Kocuria* were also detected in the *rpoB* clone
498 library and in the next generation 16S rRNA sequencing approach.

499 The shifts in fungal-associated bacterial community led to clear changes in morphology and
500 behaviour of *M. hiemalis*. For instance, the hyphal extension of the original isolate was

501 significantly faster as compared to the antibiotic-treated isolate. The direct competition
502 experiment between the two isolates revealed an advantage in plate colonization for the
503 original M0 isolates indicating that the fast hyphal extension can be beneficial for *M. hiemalis*
504 in terms of resource competition. Whereas bacterial motility in water-unsaturated soil is
505 assumed to be highly restricted, mycelia spread efficiently in the soil, penetrate air-water
506 interfaces and cross over air-filled pores (Jimenez-Sanchez et al., 2015; Warmink et al., 2011).
507 Hence, fast hyphal extension may be beneficial for bacteria and, therefore, may be stimulated
508 by the *M. hiemalis*-associated bacterial community. Similar stimulation of hyphal growth by
509 bacteria was reported for the interaction of *Amanita muscaria* and *Streptomyces* sp. Ach505
510 where bacteria showed an enhanced production of the secondary metabolite auxofuran, which
511 promotes the extension of the fungal mycelium (Frey-Klett et al., 2011).

512 Besides hyphal extension, susceptibility to antifungal compounds produced by bacteria was
513 also significantly affected by changes in the *M. hiemalis* associated bacterial community,
514 thereby the antibiotic-treated isolate appeared to be more sensitive to antifungal compounds
515 as compared to the original isolate. This may point at role of the fungus-associated bacteriome
516 in the protection of its host.

517 Fungal-associated bacteria have been shown to affect secondary metabolism of the fungi
518 including the production of volatiles (Minerdi *et al.*, 2008; Splivallo et al., 2015; Vahdatzadeh
519 et al., 2015). Due to the shifts in bacterial community, we observed changes in fungal
520 pigmentation and volatile emission. In the last years, it has become evident that microbial
521 volatiles can play major roles in long-distance interactions within soil microbial communities
522 acting as infochemicals or antimicrobial compounds (Effmert et al., 2012; Schmidt et al.,
523 2015a; Schmidt et al., 2015b; Schulz-Bohm et al., 2015). The shifts in fungal-associated
524 bacterial community led to shifts in volatile production of the original and the antibiotic-treated
525 *M. hiemalis* isolates which, consequently, affected their interactions. The volatiles emitted by
526 the original M0 isolate on the nutrient-rich PDA inhibited the growth of *C. fungivorans* Ter 331,
527 whereas the antibiotic-treated MA isolate stimulated the growth of this strain. Bacteria from

528 the genus *Collimonas*, were previously shown to colonize and grow on living fungal hyphae,
529 a phenomenon called mycophagy (De Boer et al., 2004; Leveau et al., 2010). Interestingly,
530 the motility of *C. fungivorans* Ter 331 was significantly stimulated by the volatiles emitted by
531 the antibiotic-treated *M. hiemalis* implying that volatiles might play a role as long-distance
532 signals for attracting such mycophagous bacteria. Both, volatile-mediated interactions and
533 direct interactions may indicate that the antibiotic-treated isolate is more attractive and
534 susceptible to mycophagous bacterium as compared to the original isolate.

535 The composition and abundance of volatiles was affected by the nutrient conditions with more
536 volatiles produced on the nutrient-rich PDA media. Several independent studies have reported
537 that the volatile profiles of bacteria and fungi are strongly dependent on growth conditions,
538 interactions and nutrient availability (Garbeva et al., 2014b; Schmidt et al., 2015b; Schulz-
539 Bohm et al., 2015; Tyc et al., 2015; Weikl et al., 2016). Hence, it is questionable whether the
540 same volatile-mediated interactions will occur in nature under nutrient-limited conditions. In
541 soil, the rhizosphere is a “hot-spot” of microbial activity, where approximately 20 to 40% of the
542 photosynthetic carbon fixed by a plant is released as root exudates (Jones et al., 2009;
543 Philippot et al., 2013). For a long time, it has been assumed that the rhizosphere is mainly
544 dominated by bacteria, however, recent studies revealed significant utilization of root exudates
545 by saprotrophic fungi (Hannula et al., 2012). Therefore fungal-bacterial interactions as the one
546 described in this study may take place in the rhizosphere and have an effect on the
547 rhizosphere microbiome, which consequently, may play an important role for plant growth and
548 health.

549 Overall, in the present study we described diverse bacterial community associated with the
550 saprotrophic fungus *M. hiemalis* most probably carried as a contaminant from the natural
551 environment. Our results revealed that antibiotic treatments can cause shifts in this bacterial
552 community that consequently affect the host in terms of morphology, behavior, secondary
553 metabolite production and interactions.

554 Evidence has emerged over the past years that endo- and ectosymbiotic bacteria are
555 widespread in fungi. Hence, similar to animals and plants, fungi are never alone as they
556 constantly carry their bacteriome. It remains questionable if the fungal isolates in the pure
557 culture collections are really free of bacteria, as the isolation with antibacterials is clearly not
558 sufficient to eradicate the associated bacteria. Future research should be directed to study
559 fungal bacteriome, bacterial localization, the mechanisms of interactions, and its significance
560 for evolution, trophic interactions and ecosystem functioning.

561

562

563 **5. Acknowledgments**

564 We thank Saskia Gerards, Hans Zweers, Mattias Hollander and Cornelis Hordijk (Netherlands
565 Institute of Ecology) for technical assistance, bioinformatics support and helpful discussions.

566 This study is financed by the Netherlands Organization for Scientific Research (NWO), VIDI
567 personal grant (864.11.015). This is publication XXXX of the NIOO-KNAW.

568

569 6. References

- 570
- 571 Adékambi, T., et al., 2009. The *rpoB* gene as a tool for clinical microbiologists. Trends Microbiol. 17,
572 37-45.
- 573 Altschul, S. F., et al., 1990. Basic local alignment search tool. J Mol Biol. 215, 403-410.
- 574 Ballhausen, M. B., et al., 2016. Trait Differentiation within the Fungus-Feeding (Mycophagous)
575 Bacterial Genus *Collimonas*. Plos One. 11.
- 576 Benoit, I., et al., 2015. *Bacillus subtilis* attachment to *Aspergillus niger* hyphae results in mutually
577 altered metabolism. Environmental Microbiology. 17, 2099-2113.
- 578 Bonfante, P., Anca, I. A., 2009. Plants, mycorrhizal fungi, and bacteria: A network of interactions.
579 Annual Review of Microbiology. 63, 363-383.
- 580 De Boer, W., et al., 2008. Rhizosphere bacteria from sites with higher fungal densities exhibit greater
581 levels of potential antifungal properties. Soil Biology and Biochemistry. 40, 1542-1544.
- 582 De Boer, W., et al., 2004. *Collimonas fungivorans* gen. nov., sp nov., a chitinolytic soil bacterium with
583 the ability to grow on living fungal hyphae. International Journal of Systematic and Evolutionary
584 Microbiology. 54, 857-864.
- 585 De Mendiburu, F., 2014. agricolae : statistical procedures for agricultural research. R package version
586 1.2-1. <http://CRAN.R-project.org/package=agricolae>.
- 587 De Ridder-Duine, A. S., et al., 2005. Rhizosphere bacterial community composition in natural stands
588 of *Carex arenaria* (sand sedge) is determined by bulk soil community composition. Soil Biology and
589 Biochemistry. 37, 349-357.
- 590 De Rooij-van der Goes, P. C. E. M., et al., 1995. Analysis of nematodes and soil-borne fungi from
591 *Ammophila arenaria* (Marram grass) in Dutch coastal foredunes by multivariate techniques. European
592 Journal of Plant Pathology. 101, 149-16.
- 593 Deveau, A., et al., 2010. Role of fungal trehalose and bacterial thiamine in the improved survival and
594 growth of the ectomycorrhizal fungus *Laccaria bicolor* S238N and the helper bacterium *Pseudomonas*
595 *fluorescens* BBc6R8. Environmental Microbiology Reports. 2, 560-568.
- 596 Duponnois, R., Garbaye, J., 1990. Some mechanisms involved in growth-stimulation of
597 ectomycorrhizal fungi by bacteria. Canadian Journal of Botany-Revue Canadienne de Botanique. 68,
598 2148-2152.
- 599 Effmert, U., et al., 2012. Volatile mediated interactions between bacteria and fungi in the soil. J Chem
600 Ecol. 38, 665-703.
- 601 Frey-Klett, P., et al., 2011. Bacterial-fungal interactions: Hyphens between agricultural, clinical,
602 environmental, and food microbiologists. Microbiology and Molecular Biology Reviews. 75, 583-+.
- 603 Garbeva, P., et al., 2014a. Volatile-mediated interactions between phylogenetically different soil
604 bacteria. Front Microbiol. 5, 289.
- 605 Garbeva, P., et al., 2014b. Volatiles produced by the mycophagous soil bacterium *Collimonas*. FEMS
606 Microbiology Ecology. 87, 639-649.
- 607 Garbeva, P., et al., 2011. Transcriptional and antagonistic responses of *Pseudomonas fluorescens*
608 Pf0-1 to phylogenetically different bacterial competitors. The ISME Journal. 5, 973-85.
- 609 Gee, J. E., et al., 2011. Characterization of *Burkholderia rhizoxinica* and *B. endofungorum* isolated
610 from clinical specimens. Plos One. 6.

- 611 Ghignone, S., et al., 2012. The genome of the obligate endobacterium of an AM fungus reveals an
612 interphyllum network of nutritional interactions. *The ISME Journal*. 6, 136-145.
- 613 Hall, T. A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program
614 for Windows 95/98/NT. *Nucleic Acids Symposium Series*. 41, 95-98.
- 615 Hannula, S. E., et al., 2012. ¹³C pulse-labeling assessment of the community structure of active fungi
616 in the rhizosphere of a genetically starch-modified potato (*Solanum tuberosum*) cultivar and its
617 parental isoline. *New Phytol.* 194, 784-99.
- 618 Heck, K. L., et al., 1975. Explicit calculation of rarefaction diversity measurement and determination of
619 sufficient sample size. *Ecology*. 56, 1459-1461.
- 620 Hurlbert, S. H., 1971. The nonconcept of species diversity: a critique and alternative parameters.
621 *Ecology*. 52, 577-586.
- 622 Jimenez-Sanchez, C., et al., 2015. Impact of Dissolved Organic Matter on Bacterial Tactic Motility,
623 Attachment, and Transport. *Environmental Science & Technology*. 49, 4498-4505.
- 624 Jones, D. L., et al., 2009. Carbon flow in the rhizosphere: Carbon trading at the soil-root interface.
625 *Plant and Soil*. 321, 5-33.
- 626 Kobayashi, D. Y., Crouch, J. A., 2009. Bacterial/fungal interactions: From pathogens to mutualistic
627 endosymbionts. *Annual Review of Phytopathology*. 47, 63-82.
- 628 Lackner, G., et al., 2011. Endofungal bacterium controls its host by an hrp type III secretion system.
629 *The ISME Journal*. 5, 252-261.
- 630 Lackner, G., et al., 2009. Endofungal bacteria as producers of mycotoxins. *Trends in Microbiology*.
631 17, 570-576.
- 632 Lane, D. J., 16S/23S rRNA sequencing. In: E. Stackebrandt, M. Goodfellow, (Eds.), *Nucleic acid*
633 *techniques in bacterial systematics*. John Wileys and Sons Ltd., Chichester, United Kingdom, 1991,
634 pp. 115–175.
- 635 Leveau, J. H., et al., 2010. The bacterial genus *Collimonas*: mycophagy, weathering and other
636 adaptive solutions to life in oligotrophic soil environments. *Environ Microbiol.* 12, 281-92.
- 637 Messing, J., 1993. M13 cloning vehicles. Their contribution to DNA sequencing. *Methods Mol Biol.* 23,
638 9-22.
- 639 Naito, M., et al., 2015. Minimal genomes of mycoplasma-related endobacteria are plastic and contain
640 host-derived genes for sustained life within *Glomeromyco*. *Proc Natl Acad Sci U S A*. 112, 7791-6.
- 641 Ohshima, S., et al., 2016. *Mycoavidus cysteinexigens* gen. nov., sp nov., an endohyphal bacterium
642 isolated from a soil isolate of the fungus *Mortierella elongata*. *International Journal of Systematic and*
643 *Evolutionary Microbiology*. 66, 2052-2057.
- 644 Oliveira, L. G., et al., 2013. Diversity of filamentous fungi isolated from the soil in the semiarid area,
645 Pernambuco, Brazil. *Journal of Arid Environments*. 95, 49-54.
- 646 Partida-Martinez, L. P., et al., 2007a. *Burkholderia rhizoxinica* sp. nov. and *Burkholderia*
647 *endofungorum* sp. nov., bacterial endosymbionts of the plant-pathogenic fungus *Rhizopus*
648 *microsporus*. *International Journal of Systematic and Evolutionary Microbiology*. 57, 2583-2590.
- 649 Partida-Martinez, L. P., Hertweck, C., 2005. Pathogenic fungus harbours endosymbiotic bacteria for
650 toxin production. *Nature*. 437, 884-888.

- 651 Partida-Martinez, L. P., et al., 2007b. Endosymbiont-dependent host reproduction maintains bacterial-
652 fungal mutualism. *Current Biology*. 17, 773-777.
- 653 Philippot, L., et al., 2013. Going back to the roots: The microbial ecology of the rhizosphere. *Nature*
654 *Reviews: Microbiology*. 11, 789-799.
- 655 Pluskal, T., et al., 2010. MZmine 2: Modular framework for processing, visualizing, and analyzing
656 mass spectrometry-based molecular profile data. *BMC Bioinformatics*. 11, 1-11.
- 657 Saitou, N., Nei, M., 1987. The neighbor-joining method: a new method for reconstructing phylogenetic
658 trees. *Mol Biol Evol*. 4, 406-425.
- 659 Salvioli, A., et al., 2010. Endobacteria affect the metabolic profile of their host *Gigaspora margarita*,
660 an arbuscular mycorrhizal fungus. *Environmental Microbiology*. 12, 2083-2095.
- 661 Salvioli, A., et al., 2016. Symbiosis with an endobacterium increases the fitness of a mycorrhizal
662 fungus, raising its bioenergetic potential. *ISME J*. 10, 130-44.
- 663 Sato, Y., et al., 2010. Detection of Betaproteobacteria inside the mycelium of the fungus *Mortierella*
664 *elongata*. *Microbes and Environments*. 25, 321-324.
- 665 Schloss, P. D., Handelsman, J., 2005. Introducing DOTUR, a computer program for defining
666 operational taxonomic units and estimating species richness. *Appl Environ Microbiol*. 71, 1501-1506.
- 667 Schloss, P. D., et al., 2004. Integration of microbial ecology and statistics: a test to compare gene
668 libraries. *Appl Environ Microbiol*. 70, 5485-5492.
- 669 Schmidt, R., et al., 2015a. Volatile affairs in microbial interactions. *The ISME Journal*. 9, 2329-2335.
- 670 Schmidt, R., et al., 2015b. Microbial Small Talk: Volatiles in Fungal-Bacterial Interactions. *Front*
671 *Microbiol*. 6, 1495.
- 672 Schroeckh, V., et al., 2009. Intimate bacterial-fungal interaction triggers biosynthesis of archetypal
673 polyketides in *Aspergillus nidulans*. *Proceedings of the National Academy of Sciences of the United*
674 *States of America*. 106, 14558-14563.
- 675 Schulz-Bohm, K., et al., 2015. A fragrant neighborhood: volatile mediated bacterial interactions in soil.
676 *Front Microbiol*. 6, 1212.
- 677 Sharma, M., et al., 2008. Detection and identification of bacteria intimately associated with fungi of the
678 order Sebaciales. *Cell Microbiol*. 10, 2235-46.
- 679 Singh, M., et al., 2015. Soil fungi for mycoremediation of arsenic pollution in agriculture soils. *Journal*
680 *of Applied Microbiology*. 119, 1278-1290.
- 681 Smith, M. G., et al., 2004. Microbial synergy via an ethanol-triggered pathway. *Molecular and Cellular*
682 *Biology*. 24, 3874-3884.
- 683 Splivallo, R., et al., 2015. Bacteria associated with truffle-fruited bodies contribute to truffle aroma.
684 *Environmental Microbiology*. 17, 2647-2660.
- 685 Stopnisek, N., et al., 2016. Molecular mechanisms underlying the close association between soil
686 *Burkholderia* and fungi. *The ISME Journal*. 10, 253-264.
- 687 Tamura, K., et al., 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood,
688 evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 28, 2731-2739.

689 Torres-Cortes, G., et al., 2015. Mosaic genome of endobacteria in arbuscular mycorrhizal fungi:
690 Transkingdom gene transfer in an ancient mycoplasma-fungus association. Proceedings of the
691 National Academy of Sciences of the United States of America. 112, 7785-7790.

692 Tyc, O., et al., 2014. Impact of interspecific interactions on antimicrobial activity among soil bacteria.
693 Front Microbiol. 5, 567.

694 Tyc, O., et al., 2015. Volatiles in Inter-Specific Bacterial Interactions. Front Microbiol. 6, 1412.

695 Vahdatzadeh, M., et al., 2015. The Role of the Microbiome of Truffles in Aroma Formation: a Meta-
696 Analysis Approach. Appl Environ Microbiol. 81, 6946-52.

697 Warmink, J. A., et al., 2011. Hitchhikers on the fungal highway: The helper effect for bacterial
698 migration via fungal hyphae. Soil Biology and Biochemistry. 43, 760-765.

699 Warmink, J. A., et al., 2009. Universal and species-specific bacterial 'fungiphiles' in the mycospheres
700 of different basidiomycetous fungi. Environmental Microbiology. 11, 300-312.

701 Weigl, F., et al., 2016. Sesquiterpene emissions from *Alternaria alternata* and *Fusarium oxysporum*:
702 Effects of age, nutrient availability, and co-cultivation. Scientific Reports. 6, 22152.

703 White, T. J., et al., Amplification and direct sequencing of fungal ribosomal RNA genes for
704 phylogenetics. In: M. A. Innis, et al., Eds.), PCR protocols: A guide to methods and applications.
705 Academic Press Inc, New York, 1990, pp. 315–322.

706 Xia, J., et al., 2012. MetaboAnalyst 2.0-a comprehensive server for metabolomic data analysis.
707 Nucleic Acids Research. 40, W127-W133.

708 Xia, J., et al., 2015. MetaboAnalyst 3.0-making metabolomics more meaningful. Nucleic Acids
709 Research. 43, W251-W257.

710 Zapras, A., et al., 2010. Abundance of novel and diverse tfdA-like genes, encoding putative
711 phenoxyalkanoic acid herbicide-degrading dioxygenases, in soil. Applied and Environmental
712 Microbiology. 76, 119-128.

713

714 7. Figures and Tables

715

716 **Figure 1.** Phylogenetic distribution (A) and rarefaction analysis (B and C) of *rpoB* gene
717 sequences from bacteria associated to *M. hiemalis*. Sequences retrieved from *M. hiemalis*,
718 original isolate (M0) and the isolate treated with antibiotics (MA), were assigned to species-
719 level OTUs based on a threshold value of 97% identity. Data corresponding to M0 and MA are
720 indicated in black and grey, respectively. Rarefaction analysis was performed for all
721 sequences obtained from both isolates (B) as well as for the subset per fungal isolate (C).
722 Solid lines in panel B and C represent 95% confidence intervals.

723

724 **Figure 2.** Phylogenetic association and characteristics of bacterial and yeast isolates
725 associated to *M. hiemalis*. Bacteria and yeast were isolated from *M. hiemalis*, original isolate
726 (M0) and isolate treated with antibiotics (MA). A Maximum-Likelihood tree of representative
727 16S rRNA gene sequences was calculated. Percentages of replicate trees (> 50%) in which
728 the associated taxa clustered together in the bootstrap test (10,000 bootstraps) are shown
729 next the branches. Filled squares indicate confirmed tree topology with Maximum-Parsimony
730 and Neighbor Joining (bootstraps > 70 %) calculations with the same dataset. Accession
731 numbers are included in parentheses. Table next to the tree represents general characteristics
732 of the bacterial isolates such as resistance to rifampicin (Rif) and kanamycin (Kan). Scale bar
733 in microscopic pictures represent 2 μm .

734

735 **Figure 3.** Hyphal extension of *M. hiemalis* isolates. Both, the original (M0) and antibiotic-
736 treated isolated (MA) were incubated on 0.5 strength Potato dextrose agar (PDA) or water-
737 agar (WA).

738

739 **Figure 4.** Growth of the *M. hiemalis* isolates M0 and MA (original isolate and antibiotic- treated,
740 respectively) on 0.5 strength Potato dextrose agar for 7 (A, D, G), 14 (B, E, H) and 21 days

741 (C, F, I). Fungal isolates were incubated either alone (M0: G-I; MA: D-F) or together (left: MA;
742 right: M0) on one Petri-dish (A-C).

743

744 **Figure 5.** Effect of two soil bacteria strains on hyphal extension of the *M. hiemalis* isolates M0
745 (A-F) and MA (G-L). The bacteria were pre-incubated for three days without fungus followed
746 by a four days incubation with the fungus. M0 represents the original and MA the antibiotic-
747 treated fungal isolate. Overviews of the assays are presented in panels A,C, E, G, I and K and
748 detailed views of mycelia growth in panel B, D, F, H, J and L.

749

750 **Figure 6.** PLS-DA score plot of volatile compounds produced by the original (M0) and
751 antibiotic-treated (MA) *M. hiemalis* isolates. A multivariate analysis was performed on mass
752 features of detected compounds which were emitted by the fungi on water-agar (WA; panel
753 A) or Potato dextrose agar (PDA; panel B) Controls, volatiles released by the agar media
754 without *M.hiemalis*, are included in the analysis..

755

756 **Figure 7.** Effect of fungal volatiles on the development (A and B) and on swimming motility (C
757 and D) of two rhizobacterial strains. The rhizobacteria were exposed to volatiles produced by
758 original (M0) and antibiotic-treated *M. hiemalis* isolate (MA) growing on 0.5 Potato dextrose
759 agar (PDA) or water-agar (WA). Bacterial growth and motility were assessed by counting of
760 colonies appearing at the highest dilution and measuring of the colony extension, respectively.
761 Different letters represent statistical different values tested by One-way-ANOVA.

762

763 **Table 1.** Volatile organic compounds produced by *Mucor hiemalis* on potato dextrose agar.