



Royal Netherlands Academy of Arts and Sciences (KNAW) KONINKLIJKE NEDERLANDSE AKADEMIE VAN WETENSCHAPPEN

Zymoseptoria gen. nov.: a new genus to accommodate Septoria-like species occurring on graminicolous hosts

Quaedvlieg, W.; Kema, G.H.J.; Groenewald, J.Z.; Verkley, G.J.M.; Seifbarghi, S.; Razavi, M.; Gohari, A.M.; Mehrabi, R.; Crous, P.W.

published in

Persoonia

2011

DOI (link to publisher)

[10.3767/003158511X571841](https://doi.org/10.3767/003158511X571841)

document version

Publisher's PDF, also known as Version of record

[Link to publication in KNAW Research Portal](#)

citation for published version (APA)

Quaedvlieg, W., Kema, G. H. J., Groenewald, J. Z., Verkley, G. J. M., Seifbarghi, S., Razavi, M., Gohari, A. M., Mehrabi, R., & Crous, P. W. (2011). *Zymoseptoria gen. nov.*: a new genus to accommodate Septoria-like species occurring on graminicolous hosts. *Persoonia*, 26, 57-69. <https://doi.org/10.3767/003158511X571841>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the KNAW public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the KNAW public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

pure@knaw.nl



Zymoseptoria gen. nov.: a new genus to accommodate Septoria-like species occurring on graminicolous hosts

W. Quaedvlieg^{1, 2}, G.H.J. Kema³, J.Z. Groenewald¹, G.J.M. Verkley¹, S. Seifbarghi⁴,
M. Razavi⁴, A. Mirzadi Gohari⁴, R. Mehrabi^{5, 6}, P.W. Crous^{1, 2, 5}

Key words

Hordeum vulgare
ITS
LSU
multilocus sequence typing
Mycosphaerella
Septoria
systematics
Triticum aestivum

Abstract The *Mycosphaerella* complex is both poly- and paraphyletic, containing several different families and genera. The genus *Mycosphaerella* is restricted to species with *Ramularia* anamorphs, while *Septoria* is restricted to taxa that cluster with the type species of *Septoria*, *S. cytisi*, being closely related to *Cercospora* in the *Mycosphaerellaceae*. Species that occur on graminicolous hosts represent an as yet undescribed genus, for which the name *Zymoseptoria* is proposed. Based on the 28S nrDNA phylogeny derived in this study, *Zymoseptoria* is shown to cluster apart from *Septoria*. Morphologically species of *Zymoseptoria* can also be distinguished by their yeast-like growth in culture, and the formation of different conidial types that are absent in *Septoria* s.str. Other than the well-known pathogens such as *Z. tritici*, the causal agent of septoria tritici blotch on wheat, and *Z. passerinii*, the causal agent of septoria speckled leaf blotch of barley, both for which epitypes are designated, two leaf blotch pathogens are also described on graminicolous hosts from Iran. *Zymoseptoria brevis* sp. nov. is described from *Phalaris minor*, and *Z. halophila* comb. nov. from leaves of *Hordeum glaucum*. Further collections are now required to elucidate the relative importance, host range and distribution of these species.

Article info Received: 28 January 2011; Accepted: 12 March 2011; Published: 5 April 2011.

INTRODUCTION

More than 10 000 names have been described in the genus *Mycosphaerella* (*Capnodiales*, *Dothideomycetes*) and its associated anamorph genera (*Cercospora*, *Pseudocercospora*, *Septoria*, *Ramularia*, etc.) (Crous et al. 2009a), making it one of the largest genera of plant pathogenic Ascomycetes known to date (Crous 2009). However, in contrast to earlier phylogenetic studies based on the ITS region (Stewart et al. 1999, Crous et al. 1999, 2000, 2001, Goodwin et al. 2001), more robust multi-gene phylogenies have revealed *Mycosphaerella* to be polyphyletic (Crous et al. 2007, 2009b, Schoch et al. 2009a, b), suggesting that *Mycosphaerella* s.l. should be subdivided to reflect natural groups (genera) as defined by their anamorphs.

The genus *Mycosphaerella* is typified by *M. punctiformis*, which has a *Ramularia* anamorph, *R. endophylla* (Verkley et al. 2004a). Ever since it was established, the name *Mycosphaerella* has been used to describe related and unrelated, small loculoascmycetes (in some cases even asexual coelomycetes) (Aptroot 2006), prompting Crous et al. (2009b), to suggest that the older generic name *Ramularia* (1833), rather than the confused name *Mycosphaerella* (1884) should be used for this well-defined morphologic (Braun 1998) and phylogenetic clade of fungi (Crous et al. 2009b, Kirschner 2009).

The genus *Septoria* Sacc. (1884) currently contains almost 3 000 species (Verkley & Priest 2000, Verkley et al. 2004b), several of which have *Mycosphaerella*-like teleomorphs. The type species is *Septoria cytisi* (Fig. 1), a pathogen of *Cytisus laburnum* (= *Laburnum anagyroides*). *Septoria* represents a polyphyletic assembly of anamorph genera that cluster mostly in the *Mycosphaerellaceae* (a family incorporating many plant pathogenic coelomycetes), although *Septoria*-like anamorphs have also evolved outside this family (Crous et al. 2009b). In this regard some *Septoria* species on graminicolous hosts (e.g. *S. passerinii* and *S. tritici*) have a distinct dimorphic lifestyle. Besides their mycelial state, they can exhibit a yeast-like growth in culture via microcyclic conidiation, distinguishing them from *Septoria* s.str. Furthermore, phylogenetically the *Septoria*-like species occurring on graminicolous hosts have also been found to cluster apart from *Septoria* species occurring on other hosts (Crous et al. 2001, Verkley et al. 2004b). This clear phylogenetic separation, together with the unique yeast-like growth for *S. tritici* and *S. passerinii*, led to the hypothesis that the *S. tritici* clade did not belong to *Septoria* s.str., but should be classified as a separate genus. In order to prove this hypothesis, the phylogenetic relationship of the type species of the genus *Septoria* (*S. cytisi*) needs to be determined. However these data are not currently available, as other than herbarium material, we have not been able to recollect or locate any living strains of *S. cytisi*.

The aims of this study were thus to isolate and sequence part of the nuclear ribosomal DNA operon from *S. cytisi* herbarium material, and to test the hypothesis whether the *S. tritici* clade can represent a new genus of fungi. A further aim was to resolve the identity of *Septoria*-like species occurring on graminicolous hosts. To this end partial gene sequences of five loci viz. actin (ACT), calmodulin (CAL), β -tubulin (TUB), RNA polymerase II second largest subunit (RPB2) and 28S nuclear ribosomal RNA gene (LSU) were generated and analysed.

¹ CBS-KNAW Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands;

corresponding author e-mail: w.quaedvlieg@cbs.knaw.nl.

² Microbiology, Department of Biology, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands.

³ Plant Research International, Biointeractions and Plant Health, P.O. Box 16, 6700 AA, Wageningen, The Netherlands.

⁴ Iranian Research Institute of Plant Protection, Department of Plant Pathology, P.O. Box 19395-1454, Tehran, Iran.

⁵ Wageningen University and Research Centre (WUR), Laboratory of Phytopathology, Droevendaalsesteeg 1, 6708 PB Wageningen, The Netherlands.

⁶ Seed and Plant Improvement Institute, P.O. Box 4119, Karaj 31585, Iran.

Table 1 Details of cultures subjected to DNA sequencing.

Species	Isolate no ¹	Host	Location	Collected by	GenBank Accession no ²						LSU	
					ACT	CAL	ITS	TUB	RPB2	RPB2		
<i>Cercospora apii</i>	CBS 118712	<i>Hedysarum coronarium</i>	Fiji	P. Tyler	–	–	–	–	–	–	–	GQ852583
<i>C. arminensis</i>	CBS 137.56	<i>Beta vulgaris</i>	Italy	M. Ribaldi	–	–	–	–	–	–	–	JF700933
<i>C. beticola</i>	CBS 124.31		Romania	–	–	–	–	–	–	–	–	JF700934
<i>Cladosporium brutnei</i>	CBS 188.54		–	–	–	–	–	–	–	–	–	JF700935
	CBS 115683	Douglas-fir pole	USA	–	–	–	–	–	–	–	–	JF700936
<i>Dissoconium australiensis</i>	CBS 120729	<i>Eucalyptus platyphylla</i>	Australia	P.W. Crous	–	–	–	–	–	–	–	GQ852588
<i>D. commune</i>	CPC 12397	<i>Eucalyptus globulus</i>	Australia	I.W. Smith	–	–	–	–	–	–	–	GQ852591
<i>D. dekkeri</i>	CPC 13479	<i>Eucalyptus camaldulensis</i>	Thailand	W. Himaman	–	–	–	–	–	–	–	GQ852595
<i>Dohistroma pini</i>	CBS 116484	<i>Pinus nigra</i>	USA	G. Adams	–	–	–	–	–	–	–	JF700937
<i>D. septosporum</i>	CPC 16799	<i>Pinus mugo uncinata</i>	The Netherlands	W. Quaedvlieg	–	–	–	–	–	–	–	JF700938
	CPC 3779 (= 112498)	<i>Pinus radiata</i>	Ecuador	–	–	–	–	–	–	–	–	JF700939
<i>Lecanosticta acicola</i>	CBS 871.95	<i>Pinus radiata</i>	France	M. Morelet	–	–	–	–	–	–	–	GU214663
	CPC 17940	<i>Pinus</i> sp.	Mexico	M. de Jesus	–	–	–	–	–	–	–	JF700940
	IMI 281598	<i>Pinus oocarpa</i>	Guatemala	H.C. Evans	–	–	–	–	–	–	–	JF700941
<i>Mycosphaerella ellipsoidea</i>	CBS 111167	<i>Eucalyptus cladocalyx</i>	South Africa	A.R. Wood	–	–	–	–	–	–	–	GU214450
<i>M. elongata</i>	CBS 120735	<i>Eucalyptus camaldulensis</i>	South Africa	M.J. Wingfield	–	–	–	–	–	–	–	JF700942
<i>M. marksi</i>	CBS 110981	<i>Eucalyptus</i> sp.	Venezuela	M.J. Wingfield	–	–	–	–	–	–	–	JF700943
<i>Mycosphaerella</i> sp.	CBS 110843	<i>Eucalyptus cladocalyx</i>	Tanzania	P.W. Crous	–	–	–	–	–	–	–	GQ852602
<i>M. vietnamensis</i>	CBS 119974	<i>Eucalyptus grandis</i>	South Africa	P.W. Crous	–	–	–	–	–	–	–	JF700944
<i>Passalora eucalypti</i>	CBS 111318	<i>Eucalyptus saligna</i>	Vietnam	T.I. Burgess	–	–	–	–	–	–	–	JF700945
<i>Phaeophloeospora eugeniae</i>	CPC 15143	<i>Eugenia uniflora</i>	Brazil	P.W. Crous	–	–	–	–	–	–	–	GU214458
<i>P. eugenicola</i>	CPC 2557	<i>Eugenia</i> sp.	Brazil	A.C. Alfenas	–	–	–	–	–	–	–	FJ493206
<i>Pseudocercospora gracilis</i>	CPC 11144	<i>Eucalyptus</i> sp.	Brazil	A.C. Alfenas	–	–	–	–	–	–	–	JF700946
<i>P. heimii</i>	CPC 11716		Indonesia	M.J. Wingfield	–	–	–	–	–	–	–	JF700947
<i>P. heimii</i> oides	CBS 111190		Brazil	A.C. Alfenas	–	–	–	–	–	–	–	JF700947
<i>P. irregulariramosa</i>	CBS 111211	<i>Eucalyptus</i> sp.	Indonesia	M.J. Wingfield	–	–	–	–	–	–	–	GU214439
<i>P. pseudo-eucalyptorum</i>	CPC 13769	<i>Eucalyptus saligna</i>	South Africa	M.J. Wingfield	–	–	–	–	–	–	–	GQ852609
<i>P. robusta</i>	CBS 111175	<i>Eucalyptus punctata</i>	South Africa	P.W. Crous	–	–	–	–	–	–	–	GQ852642
<i>P. stomatosa</i>	CBS 101953	<i>Eucalyptus robur</i>	Malaysia	M.J. Wingfield	–	–	–	–	–	–	–	JF700948
<i>Ramularia endophylla</i>	CBS 113265	<i>Protea</i> sp.	South Africa	S. Denman	–	–	–	–	–	–	–	EU167598
<i>R. eucalypti</i>	CBS 120726	<i>Quercus robur</i>	South Africa	G. Verkleij	–	–	–	–	–	–	–	DQ470968
<i>R. lamii</i>	CPC 11312	<i>Corymbia grandifolia</i>	Italy	W. Gams	–	–	–	–	–	–	–	JF700949
<i>Ramulispora sorghi</i>	CBS 110578	<i>Leonurus sibiricus</i>	Korea	H.D. Shin	–	–	–	–	–	–	–	JF700950
	CBS 110579	<i>Sorghum</i> sp.	South Africa	D. Nowell	–	–	–	–	–	–	–	JF700951
<i>Septoria azaleae</i>	CBS 352.49	<i>Rhododendron</i> sp.	South Africa	D. Nowell	–	–	–	–	–	–	–	GQ852654
<i>S. betulae</i>	CBS 116724	<i>Betula pubescens</i>	Belgium	J. van Holder	–	–	–	–	–	–	–	JF700952
<i>S. cytsi</i>	USO 378994 (Herbarium specimen)	<i>Laburnum anagyroides</i>	Czechoslovakia	J. A. Baumler	JF700932	–	–	–	–	–	–	JF700953
<i>S. gerberae</i>	CBS 410.61	<i>Gerbera jamesonii</i>	Italy	W. Gerlach	–	–	–	–	–	–	–	JF700954
<i>S. menthae</i>	CBS 404.34		Japan	T. Hemmi	–	–	–	–	–	–	–	JF700955
<i>S. rosae</i>	CBS 355.58		–	–	–	–	–	–	–	–	–	JF700956
<i>S. rubi</i>	CBS 102327	<i>Rosa</i> sp.	–	–	–	–	–	–	–	–	–	JF700957
<i>S. verbinae</i>	CBS 113481	<i>Rubus</i> sp.	The Netherlands	G. Verkleij	–	–	–	–	–	–	–	JF700958
<i>Teratosphaeria fibrillosa</i>	CBS 121707	<i>Septoria</i> sp.	New Zealand	G. Verkleij	–	–	–	–	–	–	–	JF700959
<i>T. molleirana</i>	CBS 117926	<i>Protea</i> sp.	South Africa	P.W. Crous & L. Mostert	–	–	–	–	–	–	–	GU323213
<i>T. nubifosa</i>	CPC 12830	<i>Eucalyptus globulus</i>	Australia	–	–	–	–	–	–	–	–	JF700960
<i>T. pseudo-cryptica</i>	CPC 11264	<i>Eucalyptus globulus</i>	Portugal	A. Philips	–	–	–	–	–	–	–	GQ852697
<i>T. secundaria</i>	CBS 115608	<i>Eucalyptus</i> sp.	New Zealand	J. Stalpers	–	–	–	–	–	–	–	JF700961
<i>T. suberosa</i>	CPC 13090	<i>Eucalyptus grandis</i>	Brazil	A.C. Alfenas	–	–	–	–	–	–	–	JF700962
<i>Verrucisporota daviesiae</i>	CBS 116002	<i>Eucalyptus agglomerata</i>	Australia	A.J. Cargenie	–	–	–	–	–	–	–	JF700963
<i>V. proteacearum</i>	CBS 116003	<i>Daviesia latifolia</i>	Australia	V. beilhartz	–	–	–	–	–	–	–	GQ852730
<i>Zasmidium anthuriticola</i>	CBS 118742	<i>Grevillea</i> sp.	Australia	J.L. Alcom	–	–	–	–	–	–	–	GQ852731
<i>Z. citri-grisea</i>	CPC 13467	<i>Anthurium</i> sp.	Thailand	C.F. Hill	–	–	–	–	–	–	–	GQ852732
<i>Z. nabiacense</i>	CBS 125010	<i>Eucalyptus</i> sp.	Thailand	W. Himaman	–	–	–	–	–	–	–	GQ852733
<i>Z. pseudoparkii</i>	CBS 110999	<i>Eucalyptus</i> sp.	Australia	A.J. Cargenie	–	–	–	–	–	–	–	JF700964
<i>Z. xenoparkii</i>	CBS 111185	<i>Eucalyptus grandis</i>	Colombia	M.J. Wingfield	–	–	–	–	–	–	–	JF700965
<i>Zymoseptoria brevis</i>	IRANI485C (= CPC 18102)	<i>Phalaris paradoxa</i>	Indonesia	M.J. Wingfield	JF701035	JF701103	JF700866	JF700967	JF700798	–	–	JF700966
	CPC 18106 (= 8S) = CBS 128853	<i>Phalaris minor</i>	Iran	–	JF701036	JF701104	JF700867	JF700968	JF700799	–	–	–
	IRANI486C (= CPC 18107)	<i>Phalaris minor</i>	Iran	–	JF701037	JF701105	JF700868	JF700969	JF700800	–	–	–
	CPC 18109 (= 81)	<i>Phalaris paradoxa</i>	Iran	–	JF701038	JF701106	JF700869	JF700970	JF700801	–	–	–
	CPC 18110 (= 83)	<i>Phalaris paradoxa</i>	Iran	–	JF701039	JF701107	JF700870	JF700971	JF700802	–	–	–

<i>Zymoseptoria halophila</i>	CPC 18111 (= 84)	Iran	JF701040	JF701108	JF700871	JF700972	JF700803
	CPC 18112 (= 85)	Iran	JF701041	JF701109	JF700872	JF700973	JF700804
	CPC 18113 (= 86)	Iran	JF701042	JF701110	JF700873	JF700974	JF700805
	CPC 18114 (= 87)	Iran	JF701043	JF701111	JF700874	JF700975	JF700806
	CPC 18115 (= 88)	Iran	JF701044	JF701112	JF700875	JF700976	JF700807
	IRAN1483C (= CPC 18105) = CBS 128854	Iran	JF701045	JF701113	JF700876	JF700977	JF700808
	CBS 120382	USA	JF701046	JF701114	JF700877	JF700978	JF700809
	CBS 120384	P71 × P83A, USA	JF701047	JF701115	JF700878	JF700979	JF700810
	CBS 120385	S. Ware	JF701048	JF701116	JF700879	JF700980	JF700811
	IRAN1489C (= CPC 18099)	Iran	JF701049	JF701117	CJF700880	JF700981	JF700812
	CPC 18100	Iran	JF701050	JF701118	JF700881	JF700982	JF700813
	CPC 18101	Iran	JF701051	JF701119	JF700882	JF700983	JF700814
	IRAN1484C (= CPC 18103)	Iran	JF701052	JF701120	JF700883	JF700984	JF700815
	CPC 18116	Iran	JF701053	JF701121	JF700884	JF700985	JF700816
	CPC 18117	Iran	JF701054	JF701122	JF700885	JF700986	JF700817
	CBS 392.59	E. Becker	JF701055	JF701123	AY152603	JF700987	JF700818
	CBS 398.52	Switzerland	JF701056	JF701124	JF700886	JF700988	JF700819
	IPO 01001	New Zealand	JF701057	JF701125	JF700887	JF700989	JF700820
	IPO 02158	Iran	JF701058	JF701126	JF700888	JF700990	JF700821
	IPO 03008	Germany	JF701059	JF701127	JF700889	JF700991	JF700822
	IPO 320	Romania	JF701060	JF701128	JF700890	JF700992	JF700823
	IPO 323	The Netherlands	JF701061	JF701129	AF181692	JF700993	JF700824
IPO 86013	Turkey	JF701062	JF701130	JF700891	JF700994	JF700825	
IPO 86015	Morocco	JF701063	JF701131	JF700892	JF700995	JF700826	
IPO 86036	Israel	JF701064	JF701132	JF700893	JF700996	JF700827	
IPO 87016	Uruguay	JF701065	JF701133	JF700894	JF700997	JF700828	
IPO 88004	Ethiopia	JF701066	JF701134	JF700895	JF700998	JF700829	
IPO 90012	Mexico	JF701067	JF701135	JF700896	JF700999	JF700830	
IPO 90015	Peru	JF701068	JF701136	JF700897	JF701000	JF700831	
IPO 91009	Tunisia	JF701069	JF701137	JF700898	JF701001	JF700832	
IPO 91010	Tunisia	JF701070	JF701138	JF700899	JF701002	JF700833	
IPO 91012	Tunisia	JF701071	JF701139	JF700900	JF701003	JF700834	
IPO 91014	Tunisia	JF701072	JF701140	JF700901	JF701004	JF700835	
IPO 91016	Tunisia	JF701073	JF701141	JF700902	JF701005	JF700836	
IPO 91020	Morocco	JF701074	JF701142	JF700903	JF701006	JF700837	
IPO 92002	Portugal	JF701075	JF701143	JF700904	JF701007	JF700838	
IPO 92003	Portugal	JF701076	JF701144	JF700905	JF701008	JF700839	
IPO 92005	Portugal	JF701077	JF701145	JF700906	JF701009	JF700840	
IPO 92032	Algeria	JF701078	JF701146	JF700907	JF701010	JF700841	
IPO 92050	Kenya	JF701079	JF701147	JF700908	JF701011	JF700842	
IPO 94231	USA	JF701080	JF701148	JF700909	JF701012	JF700843	
IPO 94236	USA	JF701081	JF701149	JF700910	JF701013	JF700844	
IPO 95001	Switzerland	JF701082	JF701150	JF700911	JF701014	JF700845	
IPO 95006	Syria	JF701083	JF701151	JF700912	JF701015	JF700846	
IPO 95013	Syria	JF701084	JF701152	JF700913	JF701016	JF700847	
IPO 95025	Syria	JF701085	JF701153	JF700914	JF701017	JF700848	
IPO 95026	Syria	JF701086	JF701154	JF700915	JF701018	JF700849	
IPO 95027	Syria	JF701087	JF701155	JF700916	JF701019	JF700850	
IPO 95028	Syria	JF701088	JF701156	JF700917	JF701020	JF700851	
IPO 95031	Syria	JF701089	JF701157	JF700918	JF701021	JF700852	
IPO 95046	Syria	JF701090	JF701158	JF700919	JF701022	JF700853	
IPO 95047	Algeria	JF701091	JF701159	JF700920	JF701023	JF700854	
IPO 95050	Algeria	JF701092	JF701160	JF700921	JF701024	JF700855	
IPO 95052	Algeria	JF701093	JF701161	JF700922	JF701025	JF700856	
IPO 95054	Algeria	JF701094	JF701162	JF700923	JF701026	JF700857	
IPO 95062	Algeria	JF701095	JF701163	JF700924	JF701027	JF700858	
IPO 95071	Algeria	JF701096	JF701164	JF700925	JF701028	JF700859	
IPO 95072	Algeria	JF701097	JF701165	JF700926	JF701029	JF700860	
IPO 95073	Algeria	JF701098	JF701166	JF700927	JF701030	JF700861	
IPO 95074	Algeria	JF701099	JF701167	JF700928	JF701031	JF700862	
IPO 97016	Italy	JF701100	JF701168	JF700929	JF701032	JF700863	
IPO 98115	Hungary	JF701101	JF701169	JF700930	JF701033	JF700864	
IPO 99048	France	JF701102	JF701170	JF700931	JF701034	JF700865	

1 CBS: Centraalbureau voor Schimmcultures, Utrecht, The Netherlands; CPC: Pedro Crous working collection housed at CBS; IMI: International Mycological Institute; USO: United States Department of Agriculture, National Fungus Collections (BPI); IPO: Research Institute for Plant Protection, Wageningen (IRAN); Iranian Fungal Culture Collection, Iranian Research Institute of Plant Protection. *ACT = Actin, TUB = β-tubulin, CAL = Calmodulin, LSU = 28S large subunit of the nrRNA gene, RPB2 = RNA polymerase II second largest subunit.




Fig. 1 *Septoria cytisi* (BPI 378994). a. Leaf with leaf spots; b. lesion with pycnidia oozing conidial cirrhi; c. conidiogenous cells showing sympodial and percurrent proliferation; d. conidia. — Scale bars = 10 µm.

MATERIALS AND METHODS

Isolates

Symptomatic leaves were collected from several localities (Table 1), and leaves with visible asexual fruiting bodies were immediately subjected to direct fungal isolation, or alternatively were first incubated in moist chambers to stimulate sporulation. Single-conidial isolates were established on malt extract agar (MEA; 20 g/L Biolab malt extract, 15 g/L Biolab agar) using the previously described procedure (Crous et al. 2009c). Cultures were later plated on fresh MEA, 2 % tap water agar supplemented with green, sterile barley leaves (WAB), 2 % potato-dextrose agar (PDA), and oatmeal agar (OA) (Crous et al. 2009c), and subsequently incubated at 25 °C under near-ultraviolet light to promote sporulation. Reference strains are maintained in the culture collection of the CBS-KNAW Fungal Biodiversity Centre, Utrecht, the Netherlands, the Plant Research Institute, Wageningen, the Netherlands, and the Iranian Research Institute of Plant Protection, Tehran, Iran (Table 1), and supplemented with other relevant isolates present in the CBS collection. Descriptions, nomenclature, and illustrations were deposited in MycoBank (www.mycobank.org, Crous et al. 2004).

DNA extraction, amplification and sequencing

Herbarium specimens

Ten *S. cytisi* herbarium specimens occurring on *Cytisus laburnum* (= *Laburnum anagyroides*), were obtained from the U.S.

National Fungus Collections (BPI) in Beltsville, Maryland, USA (Table 2). After microscopic inspection, the five specimens with the least amount of surface contamination (yeast and saprobes) where selected for DNA extraction (Table 2). Using a stereo microscope, ± 25 pycnidia, including their dried conidial cirrhi, where excised from each respective herbarium specimen, and suspended in tubes with 20 µL STL buffer from an E.Z.N.A.® Forensic DNA Kit (Omegabiotek, Norcross). Special care was taken to keep the amount of contaminant leaf material, excised together with the fungal tissue, as low as possible. The fungal material was kept in STL buffer to rehydrate for 24 h at 4 °C, after which the fungal cell walls were degraded by two cycles of freezing with liquid nitrogen and immediate re-heating to 99 °C.

Table 2 Herbarium specimens of *Laburnum anagyroides* infected with *Septoria cytisi*, obtained from the U.S. National Fungus Collections (BPI), Maryland, USA. Specimens marked with an asterisk were selected for DNA extraction.

BPI accession number	Host	Year collected	Location
0378986	<i>Laburnum anagyroides</i>	1913	France
0378987	<i>Laburnum anagyroides</i>	1933	Romania
0378988	<i>Laburnum anagyroides</i>	1893	Italy
0378989*	<i>Laburnum anagyroides</i>	1929	'Czechoslovakia'
0378990*	<i>Laburnum anagyroides</i>	1874	Italy
0378991*	<i>Laburnum anagyroides</i>	1885	'Czechoslovakia'
0378992	<i>Laburnum anagyroides</i>	1903	Italy
0378993*	<i>Laburnum anagyroides</i>	1929	Austria
0378994*	<i>Laburnum anagyroides</i>	1884	'Czechoslovakia'
0378995	<i>Laburnum anagyroides</i>	1876	Italy




Fig. 2 A diagrammatic representation of part of the nrDNA operon indicating the positions of the *Septoria*-specific primers used to generate ITS and LSU sequences of *S. cytisi*.

The genomic DNA extraction was performed using the 'Isolation of DNA from dried blood' protocol available in the E.Z.N.A.® Forensic DNA Kit with one modification: in order to increase the final DNA concentration, only 50 µL of preheated (70 °C) elution buffer was used to elude the DNA from the column.

Genus-specific primers had to be designed because the use of generic fungal ITS and LSU primers only generated sequences of contaminants (mostly yeasts). For the amplification reactions concerning the herbarium specimens, the Verbatim High Fidelity DNA Polymerase Kit (Thermo Scientific) was used in combination with the *Septoria*-specific S18S-2 forward primer (annealing to the nuclear rDNA operon at the 3'-end of the 18S nrRNA gene (SSU); Table 2), together with the *Septoria*-specific SITS2_Fd reverse primer (annealing to the nuclear rDNA operon at the 5'-end of the 28S nrRNA gene (LSU); Table 2), in order to amplify a region spanning the 5.8S nrRNA gene and the first and second internal transcribed spacer regions (Fig. 2). This amplification reaction was set up in a volume of 12.5 µL using 5× High Fidelity buffer (with MgCl₂), 0.8 µM of each primer, 2 µL of gDNA, 150 µM dNTP mix and 0.1 unit of Verbatim polymerase using a MyCycler thermal cycler (Bio-Rad). PCR amplification conditions were set as follows: an initial denaturation temperature at 98 °C for 2 min, followed by 50 cycles of denaturation temperature at 98 °C for 30 s, primer annealing at 52 °C for 30 s, primer extension at 72 °C for 30 s and final extension at 72 °C for 2 min. The resulting PCR products were then size-fractionated on a 3 % (w/v) agarose gel stained with ethidium bromide, excised from the gel and subsequently sequenced as described by Cheewangkoon et al. (2008).

Degradation and shearing of the *S. cytisi* herbarium gDNA made it impossible to directly amplify and sequence the approximate 1 300 bp needed to cover both the ITS and D1–D3 domains of the 28S nrDNA in a single reaction. Therefore, specific primers were developed from the obtained *S. cytisi* ITS1 sequence, spaced about 300 bp apart (Table 2, Fig. 2), which made it possible to sequentially amplify and sequence the entire regions of both the ITS, and the D1–D3 domains of the LSU of *S. cytisi* sequentially, and later to sequence it as described by Cheewangkoon et al. (2008).

Fungal cultures

Genomic DNA was extracted from mycelium growing on MEA (Table 1), using the UltraClean® Microbial DNA Isolation Kit (Mo Bio Laboratories, Inc., Solana Beach, CA, USA). These strains were screened for five loci, namely ITS, Actin (ACT), calmodulin (CAL), RNA polymerase II second largest subunit (RPB2) and β-tubulin (TUB) (Table 3). DNA amplification and sequencing reactions were performed as described by Cheewangkoon et al. (2008).

Phylogenetic analysis

To determine whether the multi-locus DNA sequence datasets were congruent, a partition homogeneity test (Farris et al. 1994) of all possible combinations was performed in PAUP v4.0b10

(Swofford 2003) with 1 000 replications. Parallel to this, a 70 % Neighbour-Joining (NJ) reciprocal bootstrap method with Maximum Likelihood distance (Mason-Gamer & Kellogg 1996, Lombard et al. 2010) was also employed to check congruency. The models of evolution for the NJ tree were estimated with Modeltest v3.7 (Posada & Crandall 1998) and bootstrap analyses (10 000 replicates) were performed in PAUP. Resulting NJ tree topologies were visually compared for conflicts between the individual gene regions. Maximum-parsimony genealogies for individual datasets and the combined dataset were estimated in PAUP using heuristic searches based on 1 000 random taxon addition sequences and the best trees were saved. All characters were weighted equally and alignment gaps were treated as missing data. Branches of zero length were collapsed and all multiple, equally most parsimonious trees were saved. Tree length (TL), consistency index (CI), retention index (RI) and the rescaled consistency index (RC) were calculated in PAUP for the equally most parsimonious trees and the resulting trees were printed with TreeView (Page 1996) and the alignments and phylogenetic trees were lodged in TreeBASE (www.treebase.org). All novel sequences derived from this study were deposited in GenBank (Table 1). Trees were either rooted to *Cladosporium bruhnei* for the LSU tree, or to *Mycosphaerella punctiformis* for the multigene tree.

Morphology

Descriptions were based on fungal cultures sporulating in vitro on WAB, incubated under continuous near-ultraviolet light for 2–4 wk. Wherever possible, 30 measurements (×1 000 magnification) were made of structures mounted in lactic acid, with the extremes of spore measurements given in parentheses. Colony colours (surface and reverse) were assessed after 1 mo on MEA, PDA and OA at 25 °C in the dark, using the colour charts of Rayner (1970).

RESULTS

ITS and LSU amplification and sequencing of *S. cytisi*

The gDNA extractions from the *S. cytisi* herbarium samples were performed on the herbarium specimens indicated in Table 2, and both the ITS and a partial LSU regions were targeted for these isolates using *Septoria*-specific primers (Table 4). An ITS amplicon length of 486 bp was achieved from herbarium sample US0378993 while the other samples yielded only partial ITS amplicons varying in length from 440 bp in sample US0378994 to ± 200 bp in sample US0378990; amplicons of sample US0378991 only yielded contamination sequences with general primers and did not amplify with either *Septoria*- or *S. cytisi*-specific primers.

A comparison between the full-length *S. cytisi* ITS sequence and 287 other *Septoria* ITS sequences that were generated as part of a larger unpublished study, broadly linked *S. cytisi* to a distinct ITS clade containing *S. astralagi* and *S. hippocastani*, basal to a clade consisting of the majority of sequenced *Septoria*

Table 3 Primer combinations used during this study for generic amplification and sequencing.

Locus	Primer	Primer sequence 5' to 3'	Orientation	Reference
Actin	ACT-512F	ATGTGCAAGGCCGTTTCGC	Forward	Carbone & Kohn (1999)
Actin	ACT2Rd	ARRTCRCGDCCRGCCATGTC	Reverse	Groenewald, unpubl. data
Calmodulin	CAL-228F	GAGTTC AAGGAGGCCTCTCC	Forward	Carbone & Kohn (1999)
Calmodulin	CAL2Rd	TGRTCNCGCTCDCGGATCATCTC	Reverse	Groenewald, unpubl. data
β-tubulin	TUB2Fd	GTBCACCTYCARACCGGYCARTG	Forward	Aveskamp et al. (2009)
β-tubulin	TUB4Rd	CCRGAYTGRCCRAARACRAAGTTGTC	Reverse	Aveskamp et al. (2009)
RPB2	fRPB2-5F	GAYGAYMGWGATCAYTTYGG	Forward	Liu et al. (1999)
RPB2	fRPB2-5F+414R	ACMANNCCCCARTGNGWRTTRTG	Reverse	Present study
LSU	LSU1Fd	GRATCAGGTAGGRATACCCG	Forward	Crous et al. (2009a)
LSU	LR5	TCCTGAGGGAAACTTCG	Reverse	Vilgalys & Hester (1990)

Table 4 *Septoria cytisi*-specific ITS and LSU primers used for amplification and sequencing. Nucleotide positions were determined relative to the ITS/LSU sequence of *Zymoseptoria tritici* (GenBank accession FN428877).

Primer name	Primer sequence 5' to 3'	Orientation	Relative position
S18S-2	CGTAGGTGAACYTGCGRAGGGATCATTACYGAGTGA	Forward	7
5.8S1Fd	CTCTTGGTTTCBVCATCG	Forward	240
SITS2_Fwd	CCGCCCGCACTCCGAAGCGATTAATGAAATC	Forward	459
SITS2_Rev	GATTTTCAATCGCTTCGGAGTGCGGGCGG	Reverse	459
LSU_Sep_230_Fwd	TATGTGACCGGCCCGCACCCCTTTAC	Forward	710
LSU_Sep_230_Rev	GTAAGGGTGGCGGGCCGGTCACATA	Reverse	710
LSU_Sep_530_Fwd	AAGACCTTAGGAATGTAGCTCACCT	Forward	999
LSU_Sep_530_Rev	AGGTGAGCTACATTCTAAGGTCTT	Reverse	999
LSU_Sep_575_Fwd	CTTGGGCGAGGTCCGCGCT	Forward	1059
LSU_Sep_575_Rev	AGCGCGACCTCGCCCAAG	Reverse	1059
LSU_Sep_785_Rev	AGGACATCAGGATCGGTGCGAT	Reverse	1225

Annotation: ITS1 = 1–172 bp, 5.8S = 173–330 bp, ITS2 = 331–525 bp, LSU D1 & D2 domain = 525–1110 bp.

Table 5 Polymorphisms found in the ITS and LSU sequence between the *S. cytisi* herbarium specimens. Data marked with – are not available.

BPI specimen	Collection year	ITS position (bp)			LSU position (bp)					
		93	219	411	176	377	446	536	561	563
USO 378989	1929	A	–	–	T	T	G	T	T	C
USO 378993	1929	A	C	C	C	C	C	A	G	–
USO 378994	1884	C	G	T	C	C	G	A	G	G
USO 378990	1874	A	G	C	–	–	–	–	–	–

species (data not shown). Interspecific variation in the *S. cytisi* ITS sequences were present; however, it was limited to a few nucleotides per isolate sequenced (Table 5).

Amplification of the D1–D3 domains of the LSU region was attempted on the same *S. cytisi* gDNA extracts as mentioned before. A full-length sequence read of the *S. cytisi* D1–D3 domains (the first ± 900 bp of the 28S nrRNA gene) was only obtained from a single sample (USO378994). The four remaining herbarium specimens only yielded LSU sequences varying in length from 500–800 bp. Interspecific variation in the LSU nucleotide sequences was limited to a few nucleotides per sequenced isolate (Table 5).

Phylogenetic analyses

LSU dataset

During phylogenetic analyses, the *S. cytisi* LSU sequence was aligned with LSU sequence data of 64 *Capnodiales* taxa, including 19 representative *Septoria* taxa, in order to determine which of these *Septoria* isolates belonged to *Septoria* s.str. (i.e. high association with *S. cytisi*) and to establish how this clade is related to other well-established genera within the *Capnodiales*. For the LSU tree, ± 759 characters were determined for 64 *Capnodiales* taxa, including 19 *Septoria* taxa as well as the two *Cladosporium bruhnei* isolates that were used as outgroups (CPC 5101 and CBS 188.54). The phylogenetic analysis showed that 164 characters were parsimony-informative, 38 were variable and parsimony-uninformative and 557 were constant. Thirty-two equally most parsimonious trees were obtained from the heuristic search, the first of which is shown in Fig. 3 (TL = 574, CI = 0.495, RI = 0.848, RC = 0.419). The phylogenetic analysis of the *Capnodiales* LSU dataset, including *S. cytisi*, showed this species clustering in a well-defined clade incorporating the majority of the *Septoria* spp. used in this analysis, clearly delineating this clade as *Septoria* s.str. These results also show a distinct monophyletic clade that are referred to as *Zymoseptoria* gen. nov. below, which contains *S. tritici* and *S. passerinii* together with two other species in this genus.

Multi-locus dataset

For the multi-locus phylogenetic analyses of the gramincolous isolates, ± 220 nucleotides were determined for ACT, 345 for CAL, 513 for ITS, 350 for TUB, and 305 for RPB2 (see Table 3 for detailed primer description). The adjusted sequence alignment for each locus consisted of 69 ingroup taxa with *Ramularia endophylla* (*Mycosphaerella punctiformis*; strain CBS 113265) as outgroup.

The strict consensus tree (Fig. 4) based on the multi-locus maximum-parsimony analysis had an identical topology to those of the strict consensus trees obtained for the individual loci. The partition homogeneity tests for all of the possible combinations of the five gene regions consistently yielded a P-value of 0.001, and were therefore incongruent. However, the 70 % reciprocal bootstrap trees of the individual gene regions showed no conflicting tree topologies between the separate datasets. Based on the result of the 70 % reciprocal bootstrap trees (Mason-Gamer & Kellogg 1996, Cunningham 1997), the DNA sequences of the five gene regions (ACT, CAL, RPB2, TUB and ITS) were concatenated for the phylogenetic analyses.

The concatenated and manually aligned multi-locus alignment contained 70 taxa (including the outgroup sequence) and, out of the 1 723 characters used in the phylogenetic analysis, 233 were parsimony-informative, 291 were variable and parsimony-uninformative and 1 199 were constant. 810 equally parsimonious trees were obtained from the heuristic search, the first of which is shown in Fig. 4 (TL = 768, CI = 0.815, RI = 0.922, RC = 0.751). Phylogenetic results showed two well-supported new species emerging besides the conserved *S. tritici* and *S. passerinii* clades, with a significant amount of genetic variation within the *S. tritici* clade as previously found by Goodwin et al. (2007). This intraspecific variation is most likely the cause of the partition homogeneity test failure.

The overall genetic diversity of *S. tritici*, examined over five loci, was found to be quite significant within the 54 global isolates of *S. tritici* used for this study. Most of the existing phylogenetic variation observed between the *S. tritici* isolates used in the combined tree (Fig. 4) was caused by single insertion and deletion events of triplets within tandem repeats inside the ACT and RPB2 intron sequences of these isolates. The most significant




Fig. 3 The first of 32 equally most parsimonious trees obtained from a heuristic search with 1 000 random taxon additions of the LSU alignment containing representative species that currently form well-supported clades within the *Capnodiales*. The scale bar indicates 10 changes and bootstrap support values from 1 000 replicates are indicated at the nodes. Thickened lines indicate conserved branches present in the strict consensus tree.

impact of these indel events can be seen in the phylogenetic cluster containing CPC 18099–18101 (on *Aegilops tauschii*, Iran), that arises in the *S. tritici* clade of the combined tree (Fig. 4). This small clade has a bootstrap support value of 94 %, suggesting that it could represent a cryptic or ancestral lineage of what is currently considered to be *S. tritici*. Further study using more isolates would be required to address this issue.

Taxonomy

Based on the LSU dataset (Fig. 3), *S. cytisi* was shown to cluster within the major *Septoria* clade, while the taxa occurring on graminicolous hosts clustered in a separate clade, distinct from

Septoria (*S. cytisi*) and *Mycosphaerella* (*M. punctiformis*, represented by *R. endophylla*), suggesting that they represented a distinct genus in the *Mycosphaerellaceae*. Morphologically these phylogenetic differences were supported by the distinct yeast-like growth exhibited in culture by the graminicolous species, as well as their mode of conidiogenesis, e.g. phialidic, with periclinal thickening and occasional inconspicuous percurrent proliferation(s), but lacking blastic sympodial proliferation which occurs in many species of *Septoria* s.str. Based on these differences in culture, morphology and phylogeny, a new genus is hereby introduced for the taxa occurring on graminicolous hosts.




Fig. 4 The first of 810 equally most parsimonious trees obtained from a heuristic search with 1 000 random taxon additions of the combined ACT, CAL, TUB, RPB2 and ITS sequence alignment of *Zymoseptoria* spp. The scale bar indicates 10 changes and bootstrap support values from 1 000 replicates are indicated at the nodes. Thickened lines indicate conserved branches present in the strict consensus tree.

***Zymoseptoria* Quaedvlieg & Crous, gen. nov.** — MycoBank MB517922

Septoriae similis, sed aducto fermentoide, sine formatione blastica-symphodiali conidiorum, in cultura typis conidiorum usque ad 3.

Type species. *Zymoseptoria tritici* (Desm.) Quaedvlieg & Crous.

Etymology. *Zymo* = yeast-like growth; *Septoria* = *Septoria*-like in morphology.

Conidiomata pycnidial, semi-immersed to erumpent, dark brown to black, subglobose, with central ostiole; wall of 3–4 layers of brown *textura angularis*. *Conidiophores* hyaline, smooth, 1–2-septate, or reduced to conidiogenous cells, lining the inner cavity. *Conidiogenous cells* tightly aggregated, ampulliform to doliiform or subcylindrical, phialidic with periclinal thickening, or with 2–3 inconspicuous, percurrent proliferations at apex. *Type I conidia* solitary, hyaline, smooth, guttulate, narrowly




Fig. 5 *Zymoseptoria brevis* (CPC 18106) a. Pycnidium forming on barley leaves in vitro; b. colony sporulation on potato-dextrose agar; c. conidiogenous cells; d. colony on synthetic nutrient-poor agar, showing yeast-like growth; e. conidium undergoing microcyclic conidiation (arrows; Type III); f–h. pycnidiospores (Type I). — Scale bars = 10 µm.

cylindrical to subulate, tapering towards acutely rounded apex, with bluntly rounded to truncate base, transversely euseptate; hila not thickened nor darkened. On OA and PDA aerial hyphae disarticulate into phragmospores (*Type II conidia*), that again give rise to *Type I conidia* via microcyclic conidiation; yeast-like growth and microcyclic conidiation (*Type III conidia*) common on agar media.

Zymoseptoria brevis M. Razavi, Quaedvlieg & Crous, sp. nov.
— MycoBank MB517923; Fig. 5

Zymoseptoriae passerinii similis, sed conidiis minoribus, (12–)13–16(–17) × 2(–2.5) µm.

Etymology. Named after its conidia, which are shorter (*brevis*) than those of the other species.

On sterile barley leaves on WA: *Conidiomata* pycnidial, substomatal, immersed to erumpent, globose, dark brown, up to 200 µm diam, with central ostiole, 5–10 µm diam; wall of 3–4 layers of brown *textura angularis*. *Conidiophores* reduced to conidiogenous cells, or with one supporting cell, lining the inner cavity. *Conidiogenous cells* hyaline, smooth, tightly aggregated, subcylindrical to ampulliform, straight to curved, 7–15 × 2–4 µm, with 1–2 inconspicuous, percurrent proliferations at apex, 1–1.5 µm diam. *Type I conidia* solitary, hyaline, smooth, guttulate, subcylindrical to subulate, tapering towards bluntly rounded apex, with truncate base, 0–1-septate, (12–)13–16(–17) × 2(–2.5) µm; on PDA, 9–21 × 2–3.5 µm; hila not thickened nor darkened, 1–2 µm. On OA and PDA yeast-like growth and microcyclic conidiation (*Type III conidia*) common, also forming on aerial hyphae via solitary conidiogenous loci.

Culture characteristics — Colonies on PDA flat, spreading, with moderate aerial mycelium and feathery, lobate margins; surface olivaceous-grey, outer region dirty white, reverse iron-grey; on MEA more erumpent, with less aerial mycelium; surface iron-grey with patches of white, reverse greenish black; on OA somewhat fluffy with dirty white to pale olivaceous aerial

mycelium, and submerged, olivaceous-grey margin; reaching 15 mm diam after 1 mo at 25 °C; fertile.

Specimen examined. IRAN, Ilam province, Dehloran, on living leaves of *Phalaris minor*, M. Razavi, holotype CBS H-20542, cultures ex-type No 8S = CPC 18106 = CBS 128853.

Notes — *Zymoseptoria brevis* can easily be distinguished from the other taxa presently known within the genus based on its shorter conidia.

Zymoseptoria halophila (Speg.) M. Razavi, Quaedvlieg & Crous, comb. nov. — MycoBank MB517924; Fig. 6

Basionym: *Septoria halophila* Speg., Anales Mus. Nac. Hist. Nat. Buenos Aires, Ser. 3, 13: 382. 1910.

Initial symptoms of the disease were dark-brown lesions which soon became pale buff in the centre. The leaves were heavily mottled later, and the solitary, sometimes aggregated pycnidia formed on the lesions. The disease was more severe on the lower leaves. *Pycnidia* were observed on adaxial surface of the infected leaves, and were dark-brown, globose, measuring 90–150 µm, with an ostiole ± 10 µm diam. **On sterile barley leaves on WA:** *Conidiomata* pycnidial, semi-immersed to erumpent, dark brown to black, subglobose, up to 300 µm diam, with central ostiole, up to 30 µm diam; wall of 3–4 layers of brown *textura angularis*. *Conidiophores* reduced to conidiogenous cells, lining the inner cavity. *Conidiogenous cells* hyaline, smooth, tightly aggregated, ampulliform to doliiform, 10–15 × 4–7 µm, with 2–3 inconspicuous, percurrent proliferations at apex, 1–2 µm diam. *Type I conidia* solitary, hyaline, smooth, guttulate, narrowly cylindrical to subulate, tapering towards acutely rounded apex, with bluntly rounded to truncate base; basal cell long obconically truncate, 1(–3)-septate, (30–)33–38(–50) × 2(–3) µm; conidia in vivo 1–2-septate, 36–45 × 1.5–2 µm; hila not thickened nor darkened, 1–2 µm. On OA and PDA conidia can be up to 62 µm long, and aerial hyphae disarticulate into




Fig. 6 *Zymoseptoria halophila* (CPC 18105). a. Pycnidia forming on barley leaves in vitro, with oozing conidia cirrus; b–e. conidiogenous cells formed in pycnidia; f. conidia (Type I); g. colony with yeast-like growth on synthetic nutrient-poor agar; h, j–l. conidia formed as phragmospores in aerial hyphae (Type II); i, m. conidia formed via microcyclic conidiation (Type III). — Scale bars = 10 µm.

phragmospores (*Type II conidia*), that again give rise to *type I conidia* via microcyclic conidiation; yeast-like growth and microcyclic conidiation (*Type III conidia*) common on agar media.

Culture characteristics — Colonies on PDA flat, spreading, with sparse aerial mycelium and feathery, lobate margins; centre olivaceous-grey, outer region iron-grey; reverse iron-grey; on MEA surface and reverse greenish black; on OA iron-grey, reaching 20 mm diam after 1 mo at 25 °C; fertile.

Specimen examined. IRAN, Ilam province, Dehloran, on living leaves of *Hordeum glaucum*, 25 Apr. 2007, M. Razavi, specimens IRAN12892F, CBS H-20543, cultures ex-type GLS1 = IRAN1483C = CPC 18105 = CBS 128854.

Notes — The present collection of *Z. halophila* was initially reported from Iran as *S. halophila* by Seifbarghi et al. (2009) (GenBank HM100267, HM100266), based on the description of *S. halophila* provided by Priest (2006). *Zymoseptoria halophila* was originally described from *Hordeum halophilum* collected in Argentina, with conidia being (0–)1(–2)-septate, 36–58 × 1.5(–2) µm, and conidiogenous cells being 8–10 × 2.5–3.5 µm. It is likely that the various collections on *Hordeum* and *Poa* spp. from Australia listed by Priest (2006) could represent different species, but this can only be resolved once additional collections and cultures have been obtained to facilitate further molecular comparisons.

Zymoseptoria halophila is closely related to *Z. passerinii*, which is also reflected in its conidial size, which overlaps in length, but

can only be distinguished based on their difference in width. It is possible that some published records of *Z. passerinii* could in fact represent *Z. halophila*, but more collections would be required to resolve its host range and geographic distribution.

***Zymoseptoria passerinii* (Sacc.) Quaedvlieg & Crous, comb. nov.** — MycoBank MB517925; Fig. 7

Basionym: *Septoria passerinii* Sacc., Syll. Fung. (Abellini) 3: 560. 1884.

Specimens examined. ITALY, Vigheffio, near Parma, on *Hordeum murinum*, June 1879 (F. von Thümen, Mycotheca Univ. No. 1997, isotype in MEL, see Priest 2006, f. 107). — USA, North Dakota, Foster county, on *Hordeum vulgare*, coll. S. Goodwin, isol. D. Long, epitype designated here CBS H-20544, culture ex-epitype P83 = CBS 120382.

Notes — Priest (2006) reported *Z. passerinii* from several *Hordeum* species collected in Western Australia and deposited them at IMI (now in Kew), and found them to be identical to type material examined, suggesting that this pathogen is widely distributed along with its host. Ware et al. (2007) reported a *Mycosphaerella*-like teleomorph from a heterothallic mating of isolates of *Z. passerinii*. Single ascospore isolates have been deposited as CBS 120384 (P71 × P83A) and CBS 120385 (P71 × P83B). Isolate P63, which is genetically similar to P83 on the loci sequenced in this study, has been used for whole genome analysis of *Z. passerinii* (E.H. Stukenbrock, pers. comm.).




Fig. 7 *Zymoseptoria passerinii* (CBS 120382). a. Colony sporulating on potato-dextrose agar; b. colony sporulating on synthetic nutrient-poor agar; c. conidiogenous cells formed inside pycnidia; e, f. conidia from pycnidia (Type I). — Scale bars = 10 µm.




Fig. 8 *Zymoseptoria tritici* (CBS 115943). a. Conidiogenous cells formed inside pycnidia; b. conidia from pycnidia (Type I); c. colony sporulating on synthetic nutrient-poor agar, showing yeast-like growth; d, e. conidia formed via microcyclic conidiation (Type III). — Scale bars = 10 µm.

Zymoseptoria tritici (Desm.) Quaedvlieg & Crous, *comb. nov.*
— MycoBank MB517926; Fig. 8

Basionym: *Septoria tritici* Desm., Ann. Sci. Nat., Bot., sér. 2, 17: 107 (1842).

Teleomorph: '*Mycosphaerella*' *graminicola* (Fuckel) J. Schröt., in Cohn, Krypt.-Fl. Schlesien 3, 2: 340. 1894 ('1893').

Basionym: *Sphaeria graminicola* Fuckel, Fungi Rhenani Exsicc.: no. 1578. 1865.

≡ *Sphaerella graminicola* (Fuckel) Fuckel, Jahrb. Nassauischen Vereins Naturk. 23–24: 101. 1870.

Specimens examined. FRANCE, on *Triticum* sp. (holotype of *Septoria tritici*; PC). — GERMANY, Oestrich, on *Triticum repens*, Fuckel, Fungi Rhenani Exsiccati no. 1578 (L, isotype of *Mycosphaerella graminicola*). — NETHERLANDS, Brabant West, on *Triticum aestivum*, coll. R. Daamen, 6 May 1981, isol. as single conidium, W. Veenbaas, 810507/1, 7 May 1981, epitype designated here CBS H-20545, including teleomorph material on *Triticum* leaf of heterothallic mating IPO 323 (MAT 1-1) × IPO 94269 (MAT 1-2), culture ex-epitype IPO 323 = CBS 115943.

Notes — The isolate designated here as ex-epitype (IPO 323 = CBS 115943) is also the strain used in the whole genome amplification and sequencing of this species (<http://genome.jgi-psf.org/Mycgr3/Mycgr3.download.html>).

DISCUSSION

For many years the genus *Mycosphaerella* has been treated as a wide general concept to accommodate a range of related and unrelated species and genera that have small ascomata, and hyaline, 1-septate ascospores, without pseudoparaphyses (Aptroot 2006). The observation that *Mycosphaerella*-like teleomorphs were linked to more than 40 different anamorphs (Crous 2009) was thus seen as rather odd, though acceptable within this wider concept used to accommodate these thousands of mostly phytopathogenic fungi. It was only in recent years when the higher order phylogenetic relationships of *Mycosphaerella* was addressed as part of the Assembling the Fungal Tree of Life initiative (Schoch et al. 2006), that *Mycosphaerella* was shown to be polyphyletic (Crous et al. 2007), even containing different families within the *Dothideomycetes* (Crous et al. 2009a, b, Schoch et al. 2009a, b).

The fact that *Septoria* also contains significant morphological variation was commented on by Sutton (1980), who stated that the genus is heterogeneous, and should be revised, containing conidiomata that ranged from acervuli to pycnidia, and conidiogenesis that ranged from blastic sympodial to annellidic (percurrent proliferation) or phialidic (with periclinal thickening). As can be seen with the taxa treated to date, however, these characters alone are also insufficient to delineate all natural

genera, as several modes of conidiogenesis or conidiomatal types occur within the same genus in the *Septoria*-like complex. Part of the reason for the confusion surrounding the genus *Septoria* is based on the fact that until now no DNA sequence data were available for the type species, *S. cytisi*. Due to the lack of cultures of this species, DNA was subsequently extracted from several herbarium specimens. Using this technique, however, some intraspecific variation was observed in both the LSU and ITS sequences of *S. cytisi*. This could possibly be explained by geographical and temporal spread in the sampling sites, spanning 54 years from a region encompassing South and Central Europe, making some sequence variation within these specimens probable. Even if one or two nucleotides might actually be scored wrong in the US0378994-derived LSU sequence for *S. cytisi*, this would not have any impact on the phylogenetic position of *S. cytisi* within the *Septoria* s.str. clade, its nearest sister genus being *Cercospora* in the *Mycosphaerellaceae* (Groenewald et al. 2006).

As shown in the present study (Fig. 2), the genus *Mycosphaerella* is unavailable to accommodate the taxa occurring on graminicolous hosts, as *Mycosphaerella* is restricted to species with *Ramularia* anamorphs (Verkley et al. 2004a, Crous et al. 2009b). Furthermore, *Septoria* s.str. also clusters apart from the species on cereals (Fig. 3), making the name *Septoria* unavailable for these pathogens.

In the present study we introduce a novel genus *Zymoseptoria* to accommodate the *Septoria*-like species occurring on graminicolous hosts. Although species of *Zymoseptoria* tend to have phialides with apical periclinal thickening, this mode of conidiogenesis has also evolved in *Septoria* s.str. (e.g. *S. apiicola*), and is not restricted to *Zymoseptoria*. More importantly, species of *Zymoseptoria* exhibit a yeast-like growth in culture, and have up to three different conidial types that can be observed, namely Type I (pycnidial conidia), Type II (phragmospores on aerial hyphae), and Type III (yeast-like growth proliferating via microcyclic conidiation). Introducing a novel genus for this group of important plant pathogens was not taken lightly, as *Z. passerinii* causes septoria speckled leaf blotch (SSLB) on barley (*Hordeum vulgare*), and has been reported around the globe on this crop (Mathre 1997, Cunfer & Ueng 1999, Goodwin & Zismann 2001, Ware et al. 2007). *Septoria tritici* blotch (STB) is caused by *Z. tritici* (teleomorph '*Mycosphaerella*' *graminicola*), and is currently present in all major wheat growing areas. This disease is consistently ranked amongst the most damaging wheat diseases in Australia, Europe, North and South America, and in Europe more than 70 % of all the fungicides applied to wheat are to control STB (Eyal et al. 1987). Wheat, together with maize and rice directly contribute 47 % to global human consumption (Tweeten & Thompson 2009). Since 1961, wheat production has increased globally with almost 300 % on a virtually stable cultivation area of 200 M ha. This progress was largely achieved by increased average yields (FAO 2010). However, the annual growth rate of global wheat production cannot meet the global market requirements in the coming four decades (Fischer et al. 2009, Fischer & Edmeades 2010).

Although *Z. passerinii* and *Z. tritici* share many similarities (Goodwin et al. 2001) (Fig. 3, 4), both pathogens having a dimorphic lifestyle (Mehrabi et al. 2006); one major difference between them is that *Z. tritici* has a year-round and very active sexual cycle (Shaw & Royle 1993, Kema et al. 1996, Zhan et al. 2003), whereas there have been no reports of a sexual cycle for *S. passerinii* observed in nature, despite isolates of *S. passerinii* having opposite mating types being commonly found in natural populations, even on the same leaf (Goodwin et al. 2003), suggesting cryptic sex does exist for *Z. passerinii* (Ware et al. 2007). With respect to the two additional species treated in the present study, *Z. brevis* and *Z. halophila*, almost

nothing is known about their relative importance, geographical distribution, host range or sexual behaviour. Given the importance of their known host crops, however, this complex is in dire need of further study.

Acknowledgements The curator of the US National Fungus Collection in Beltsville, Maryland USA (BPI) is gratefully acknowledged for permission to extract DNA from the herbarium specimens of *S. cytisi*, without which this study would not have been possible. The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/2007-2013)/grant agreement no. 226482 (Project: QBOL - Development of a new diagnostic tool using DNA barcoding to identify quarantine organisms in support of plant health) by the European Commission under the theme 'Development of new diagnostic methods in support of Plant Health policy' (no. KBBE-2008-1-4-01). Drs M. Abbasi, R. Zare and Mr M. Aminirad helped us in providing herbarium and fungal culture collection numbers, and identifying plant species at the Department of Botany, Iranian Research Institute of Plant Protection.

REFERENCES

- Aptroot A. 2006. *Mycosphaerella* and its anamorphs: 2. Conspectus of *Mycosphaerella*. CBS Biodiversity Series 5: 1–231.
- Aveskamp MM, Verkley GJM, Gruyter J de, Murace MA, Perello A, et al. 2009. DNA phylogeny reveals polyphyly of *Phoma* section *Peyronellaea* and multiple taxonomic novelties. *Mycologia* 101: 363–382.
- Braun U. 1998. A monograph of *Cercosporiella*, *Ramularia* and allied genera (Phytopathogenic Hyphomycetes). Vol. 2. IHW-Verlag, Eching.
- Carbone I, Kohn LM. 1999. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 91: 553–556.
- Cheewangkoon R, Crous PW, Hyde KD, Groenewald JZ, To-anan C. 2008. Species of *Mycosphaerella* and related anamorphs on *Eucalyptus* leaves from Thailand. *Persoonia* 21: 77–91.
- Crous PW. 2009. Taxonomy and phylogeny of the genus *Mycosphaerella* and its anamorphs. *Fungal Diversity* 38: 1–24.
- Crous PW, Aptroot A, Kang J-C, Braun U, Wingfield MJ. 2000. The genus *Mycosphaerella* and its anamorphs. *Studies in Mycology* 45: 107–121.
- Crous PW, Braun U, Groenewald JZ. 2007. *Mycosphaerella* is polyphyletic. *Studies in Mycology* 58: 1–32.
- Crous PW, Gams W, Stalpers JA, Robert V, Stegehuis G. 2004. MycoBank: an online initiative to launch mycology into the 21st century. *Studies in Mycology* 50: 19–22.
- Crous PW, Hong L, Wingfield MJ, Wingfield BD, Kang J. 1999. *Uwebraunia* and *Dissoconium*, two morphologically similar anamorph genera with distinct teleomorph affinity. *Sydowia* 52: 155–166.
- Crous PW, Kang JC, Braun U. 2001. A phylogenetic redefinition of anamorph genera in *Mycosphaerella* based on ITS rDNA sequence and morphology. *Mycologia* 93: 1081–1101.
- Crous PW, Schoch CL, Hyde KD, Wood AR, Gueidan C, et al. 2009a. Phylogenetic lineages in the Capnodiales. *Studies in Mycology* 64: 17–47.
- Crous PW, Summerell BA, Carnegie AJ, Wingfield MJ, Hunter GC, et al. 2009b. Unravelling *Mycosphaerella*: do you believe in genera? *Persoonia* 23: 99–118.
- Crous PW, Verkley GJM, Groenewald JZ, Samson RA (eds). 2009c. *Fungal Biodiversity. CBS Laboratory Manual Series 1: 1–269*. Centraalbureau voor Schimmelcultures, Utrecht, Netherlands.
- Cunfer BM, Ueng PP. 1999. Taxonomy and identification of *Septoria* and *Stagonospora* species on small grain cereals. *Annual Review of Phytopathology* 37: 267–284.
- Cunningham CW. 1997. Can three incongruence tests predict when data should be combined? *Molecular Biology and Evolution* 14: 733–740.
- Eyal Z, Sharen AL, Prescott JM, Ginkel M van. 1987. The *Septoria* diseases of wheat: concepts and methods of disease management. Mexico, DF, CIMMYT.
- FAO. 2010. <http://faostat.fao.org/site/567/DesktopDefault.aspx?PageID=567#ancor>.
- Farris JS, Källersjö M, Kluge AG, Bult C. 1994. Testing significance of incongruence. *Cladistics* 10: 315–319.
- Fischer RA, Byerlee D, Edmeades GO. 2009. Can technology deliver on the yield challenge to 2050? Prepared for UN FAO Expert Meeting on How to Feed the World in 2050. Rome, 24–26 June 2009.
- Fischer RA, Edmeades GO. 2010. Breeding and cereal yield progress. *Crop Science* 50: S85–S98.
- Goodwin SB, Dunkle LD, Zismann VL. 2001. Phylogenetic analysis of *Cercospora* and *Mycosphaerella* based on the internal transcribed spacer region of ribosomal DNA. *Phytopathology* 91: 648–658.

- Goodwin SB, Lee TAJ van der, Cavaletto JR, Hekkert BTL, Crane CF, Kema GHJ. 2007. Identification and genetic mapping of highly polymorphic micro-satellite loci from an EST database of the septoria tritici blotch pathogen *Mycosphaerella graminicola*. *Fungal Genetics and Biology* 44: 398–414.
- Goodwin SB, Waalwijk C, Kema GHJ, Cavaletto JR, Zhang G. 2003. Cloning and analysis of the mating-type idiomorphs from the barley pathogen *Septoria passerinii*. *Molecular Genetics and Genomics* 269: 1–12.
- Goodwin SB, Zismann VL. 2001. Phylogenetic analyses of the ITS region of ribosomal DNA reveal that *Septoria passerinii* from barley is closely related to the wheat pathogen *Mycosphaerella graminicola*. *Mycologia* 93: 934–946.
- Groenewald M, Groenewald JZ, Braun U, Crous PW. 2006. Host range of *Cercospora apii* and *C. beticola*, and description of *C. apiicola*, a novel species from celery. *Mycologia* 98: 275–285.
- Kema GHJ, Verstappen ECP, Todorova M, Waalwijk C. 1996. Successful crosses and molecular tetrad progeny analyses demonstrate heterothallism in *Mycosphaerella graminicola*. *Current Genetics* 30: 251–258.
- Kirschner R. 2009. *Cercospora* and *Ramularia*. *Mycologia* 101: 110–119.
- Liu Y, Whelen S, Hall B. 1999. Phylogenetic relationships among ascomycetes: evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16: 1799–1808.
- Lombard L, Crous PW, Wingfield BD, Wingfield MJ. 2010. Multigene phylogeny and mating tests reveal three cryptic species related to *Calonectria pauciramosa*. *Studies in Mycology* 66: 15–30.
- Mason-Gamer RJ, Kellogg EA. 1996. Testing for phylogenetic conflict among molecular data sets in the tribe Triticeae (Gramineae). *Systematic Biology* 45: 524–545.
- Mathre DE. 1997. *Compendium of barley diseases*. Second edn. American Phytopathological Society Press, St. Paul, Minnesota, USA.
- Mehrabi R, Zwiers LH, Waard M de, Kema GHJ. 2006. MgHog1 regulates dimorphism and pathogenicity in the fungal wheat pathogen *Mycosphaerella graminicola*. *Molecular Plant-Microbe Interactions* 19: 1262–1269.
- Page RDM. 1996. TREEVIEW: An application to display phylogenetic trees on personal computers. *Computer Applications in the Biosciences* 12: 357–358.
- Posada D, Crandall KA. 1998. MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14: 817–818.
- Priest MJ. 2006. *Septoria*. *Fungi of Australia*. ABRIS, Canberra: CSIRO Publishing, Melbourne, Australia.
- Rayner RW. 1970. *A mycological colour chart*. CMI and British Mycological Society. Kew, Surrey, England.
- Schoch CL, Crous PW, Groenewald JZ, Boehm EWA, Burgess TI, et al. 2009a. A class-wide phylogenetic assessment of Dothideomycetes. *Studies in Mycology* 64: 1–15.
- Schoch CL, Shoemaker RA, Seifert KA, Hambleton S, Spatafora JW, Crous PW. 2006. A multigene phylogeny of the Dothideomycetes using four nuclear loci. *Mycologia* 98: 1041–1052.
- Schoch CL, Sung GH, López-Giráldez F, Townsend JP, Miadlikowska J, et al. 2009b. The Ascomycota tree of life: a phylum wide phylogeny clarifies the origin and evolution of fundamental reproductive and ecological traits. *Systematic Biology* 58: 224–239.
- Seifbarghi S, Razavi M, Aminian H, Zare R, Etebarian H. 2009. Studies on the host range of *Septoria* species on cereals and some wild grasses in Iran. *Phytopathologia Mediterranea* 48: 422–429.
- Shaw MW, Royle DJ. 1993. Factors determining the severity of epidemics of *Mycosphaerella graminicola* (*Septoria tritici*) on winter wheat in the UK. *Plant Pathology* 42: 882–899.
- Stewart EL, Liu Z, Crous PW, Szabo L. 1999. Phylogenetic relationships among some cercosporoid anamorphs of *Mycosphaerella* based on rDNA sequence analysis. *Mycological Research* 103: 1491–1499.
- Sutton BC. 1980. *The coelomycetes. Fungi imperfecti with Pycnidia, Acervuli and Stromata*. Commonwealth Mycological Institute, Kew, Surrey, England.
- Swofford DL. 2003. PAUP*. *Phylogenetic Analysis Using Parsimony* (*and other methods). Version 4. Sinauer Associates, Sunderland, Massachusetts, USA.
- Tweeten L, Thompson S. 2009. Long-term global agricultural output supply-demand balance and real farm and food prices. *Farm Policy Journal* 6.
- Verkley GJM, Crous PW, Groenewald JZ, Braun U, Aptroot A. 2004a. *Mycosphaerella punctiformis* revisited: morphology, phylogeny, and epitypification of the type species of the genus *Mycosphaerella* (Dothideales, Ascomycota). *Mycological Research* 108: 1271–1282.
- Verkley GJM, Priest MJ. 2000. *Septoria* and similar coelomycetous anamorphs of *Mycosphaerella*. *Studies in Mycology* 45: 123–128.
- Verkley GJM, Starink-Willemse M, Iperen A van, Abeln ECA. 2004b. Phylogenetic analyses of *Septoria* species based on the ITS and LSU-D2 regions of nuclear ribosomal DNA. *Mycologia* 96: 558–571.
- Vilgalys R, Hester M. 1990. Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246.
- Ware SB, Verstappen ECP, Breeden J, Cavaletto JR, Goodwin SB, et al. 2007. Discovery of a functional *Mycosphaerella* teleomorph in the presumed asexual barley pathogen *Septoria passerinii*. *Fungal Genetics and Biology* 44: 389–397.
- Zhan J, Pettway RE, McDonald BA. 2003. The global genetic structure of the wheat pathogen *Mycosphaerella graminicola* is characterized by high nuclear diversity, low mitochondrial diversity, regular recombination, and gene flow. *Fungal Genetics and Biology* 38: 286–297.