



K O N I N K L I J K E N E D E R L A N D S E
A K A D E M I E V A N W E T E N S C H A P P E N

Climate change leads to differential shifts in the timing of annual cycle stages in a migratory bird

Tomotani, B.M.; van der Jeugd, H.P.; Gienapp, P.; De la Hera, I.; Pilzecker, J.; Teichmann, Corry; Visser, M.E.

published in

Global Change Biology
2018

DOI (link to publisher)

[10.1111/gcb.14006](https://doi.org/10.1111/gcb.14006)

document version

Peer reviewed version

document license

CC BY

[Link to publication in KNAW Research Portal](#)

citation for published version (APA)

Tomotani, B. M., van der Jeugd, H. P., Gienapp, P., De la Hera, I., Pilzecker, J., Teichmann, C., & Visser, M. E. (2018). Climate change leads to differential shifts in the timing of annual cycle stages in a migratory bird. *Global Change Biology*, 24(2), 823-835. <https://doi.org/10.1111/gcb.14006>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the KNAW public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the KNAW public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

pure@knaw.nl

DR. BARBARA TOMOTANI (Orcid ID : 0000-0002-8855-4803)

Article type : Primary Research Articles

Climate change leads to differential shifts in the timing of annual cycle stages in a migratory bird

Running head: Shifts in the annual cycle of migratory birds

Barbara M. Tomotani¹, Henk van der Jeugd^{1,2}, Phillip Gienapp¹, Iván de la Hera^{1,3}, Jos Pilzecker⁴, Corry Teichmann¹, Marcel E. Visser¹

¹ Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), P.O. Box 50, 6700 AB Wageningen, The Netherlands.

²Dutch Centre for Avian Migration and Demography, Netherlands Institute of Ecology (NIOO-KNAW), P.O. Box 50, 6700 AB Wageningen, The Netherlands.

³ School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Field Campus, Cork, Ireland.

⁴Vogelwerkgroep Arnhem, Oudeweg 35, 6974 AD Leuvenheim, The Netherlands.

Corresponding author: Barbara M. Tomotani, email: b.tomotani@nioo.knaw.nl

Keywords: breeding, European pied flycatcher, *Ficedula hypoleuca*, fitness, mark-recapture, migration, moult, recruitment

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/gcb.14006

This article is protected by copyright. All rights reserved.

Type of paper: Primary Research Article

Abstract

Shifts in reproductive phenology due to climate change have been well documented in many species but how, within the same species, other annual cycle stages (*e.g.*, moult, migration) shift relative to the timing of breeding has rarely been studied. When stages shift at different rates, the interval between stages may change resulting in overlaps, and as each stage is energetically demanding, these overlaps may have negative fitness consequences. We used long-term data of a population of European pied flycatchers (*Ficedula hypoleuca*) to investigate phenological shifts in three annual cycle stages: spring migration (arrival dates), breeding (egg-laying and hatching dates) and the onset of post-breeding moult. We found different advancements in the timing of breeding compared to moult (moult advances faster) and no advancement in arrival dates. To understand these differential shifts, we explored which temperatures best explain the year-to-year variation in the timing of these stages, and show that they respond differently to temperature increases in the Netherlands, causing the intervals between arrival and breeding and between breeding and moult to decrease. Next, we tested the fitness consequences of these shortened intervals. We found no effect on clutch size, but the probability of a fledged chick to recruit increased with a shorter arrival-breeding interval (earlier breeding). Finally, mark-recapture analyses did not detect an effect of shortened intervals on adult survival. Our results suggest that the advancement of breeding allows more time for fledgling development, increasing their probability to recruit. This may incur costs to other parts of the annual cycle but, despite the shorter intervals, there was no effect on adult survival. Our results show that to fully understand the consequences of climate

change, it is necessary to look carefully at different annual cycle stages, especially for organisms with complex cycles, such as migratory birds.

Introduction

During the past decades, increases in temperature caused by global warming have affected biological systems in various ways (Walther *et al.*, 2002). One of the most evident effects are the alterations in phenology or timing of annual events across different animal and plant taxa (Crick *et al.*, 1997; Brown *et al.*, 1999; Parmesan & Yohe, 2003; Parmesan, 2006). For example, timing of flowering, breeding and migration are some of the events known to have advanced in response to the increased temperatures (Crick *et al.*, 1997; Parmesan, 2007; Charmantier & Gienapp, 2014; Thackeray *et al.*, 2016). Such changes make important to understand whether shifts in timing are fast enough to also track changes in the optimal timing for stages to occur (Visser & Both, 2005; Visser, 2008).

The use of standardised long-term datasets, especially of the same population, has allowed researchers to track how organisms respond to climate change and to identify potential causes and consequences of these responses (Visser, 2008, 2016). A relatively large number of studies have reported the effects of climate change on timing of breeding, particularly in birds (*e.g.* Crick *et al.*, 1997; Both *et al.*, 2004). A few studies also have explored if optimal breeding dates shifted with climate change and if animals were able to respond accordingly (Visser *et al.*, 2006; Reed *et al.*, 2013a, 2013b; Plard *et al.*, 2014; Stoks *et al.*, 2014; Phillimore *et al.*, 2016). These studies are crucial to our understanding of whether species can adapt to such rapid changes in the environment or not (Gienapp *et al.*, 2013).

While some organisms present quite simple annual cycles with only a breeding/non-breeding transition, others have much more complex cycles (Jacobs & Wingfield, 2000; Wingfield, 2008). For example, many species of birds and mammals also migrate, moult their

Accepted Article

plumage/pelage and/or hibernate. These additional stages of the annual cycle are likewise reported to shift due to climate change (Both & te Marvelde, 2007; Ozgul *et al.*, 2010; Charmantier & Gienapp, 2014; Morrison *et al.*, 2015; Zimova *et al.*, 2016). However, all of these stages (including breeding) are not necessarily affected in the same way by changes in temperatures (Serreze & Francis, 2006; Visser *et al.*, 2006; Visser, 2008; Both *et al.*, 2009). Moreover, because temperatures do not change at the same rate in time or space (Easterling *et al.*, 1997; Vose *et al.*, 2005; Serreze & Francis, 2006; Stocker *et al.*, 2013), it is possible that within the same population, different parts of the annual cycle also change at different rates in response to the unequal increase in temperatures (Crozier *et al.*, 2008). In this scenario, we can expect increased or decreased intervals between annual cycle stages. Therefore, climate change will also alter the time constraints across the annual cycle. This could be particularly problematic for organisms with complex annual cycles since they are supposedly more time constrained than organisms with simpler cycles (Jacobs & Wingfield, 2000; Wingfield, 2008).

Analyses showing how intervals between annual cycle stages are changing are rare in comparison to studies dealing with changes in single stages. To detect such patterns it is necessary to collect long-term data of not only one, but multiple annual cycle stages within a population. It has been reported that some species have gained more time for key intervals. For example, yellow-bellied marmots (*Marmota flaviventris*) advanced both the termination of hibernation and weaning, resulting in more time for growth and fattening for their offspring (Ozgul *et al.*, 2010). In red deer (*Cervus elaphus*), on the other hand, reproductive stages of males and females unequally advanced suggesting that males are unable to track advancements in oestrus of females. Moreover, the termination dates of males' rut advanced more than the initiation dates, which resulted in a shorter breeding window and likely reduced breeding performance (Moyes *et al.*, 2011). Barnacle geese (*Branta leucopsis*) that

Accepted Article

rapidly expanded their breeding range to temperate areas more than doubled the interval between breeding and moult, resulting in changed body mass dynamics (Van der Jeugd *et al.*, 2009; Eichhorn *et al.*, 2010). Finally, long-term data on date of arrival and breeding of migratory birds suggest that timing of breeding and migration may not respond the same way to climate change. There are reports of both shorter (Both & Visser, 2001) and longer (Ahola *et al.*, 2004; Valtonen *et al.*, 2016) intervals between arrival and breeding. Thus, for animals with more complex annual cycles, the assessment of the impacts of climate change becomes considerably more challenging, because unequal changes in the timing of stages may impose fitness costs (*e.g.*, reduce the interval between moult and breeding thereby causing more overlap; Hemborg & Lundberg, 1998).

Here, we analysed how climate change affects the timing of different annual cycle stages of a long-distance migratory bird, the European pied flycatcher. We used long-term data (1980-2015) collected from a Dutch population of flycatchers, looking at three important annual cycle stages: spring migration (arrival dates), breeding (egg-laying and hatching dates) and post-breeding moult onset. Furthermore, we included initial information on the variability in timing of autumn migration. Apart from describing the changes in patterns through time, we also investigated the potential environmental causes and fitness consequences of the variation in timing of these important stages.

Materials and Methods

Study species and study area

Pied flycatchers (*Ficedula hypoleuca* ([Pallas], 1764); Muscicapidae) are long-distance migratory birds that reproduce in Europe and winter in West Africa (Ivory Coast in the case of our study populations)(Ouwehand *et al.*, 2016, see also geolocator data in results). These

Accepted Article

birds readily accept artificial nest boxes and have low nest desertion rates, allowing the precise monitoring of their reproduction. We used long-term data from the breeding population of the forests of the Hoge Veluwe National Park (The Netherlands; 5°51'E 52°02'N). Forested areas in the park are dominated by pedunculate oaks (*Quercus robur*), northern red oaks (*Quercus rubra*), Scots pines (*Pinus sylvestris*), Larches (*Larix* spp.) and birches (*Betula* spp.). For more than 60 years nest boxes are provided year-round in an area of 171 ha, they are occupied in spring by cavity-nesting passerines such as pied flycatchers, great tits (*Parus major*), blue tits (*Cyanistes caeruleus*), nuthatches (*Sitta europaea*) and coal tits (*Periparus ater*). Voucher material of this pied flycatcher population was deposited in the ornithology collection of the Naturalis Biodiversity Center (Leiden, The Netherlands) under the inventory numbers RMNH 592347, RMNH 592348 and RMNH 592349.

Data collection

Data on timing of breeding (egg-laying, chick hatching) was collected for this pied flycatcher population since 1959, timing of migration (arrival dates and, more recently, departure dates) and timing of moult since 2005. We also obtained 35 years of arrival data of pied flycatchers breeding in a nearby location (from the Vogelwerkgroep Arnhem, the dataset used in Both & Visser, 2001, including recent years). For our final dataset, we used 35 years of egg-laying, chick hatching dates and female arrival (nest building) dates (to match the number of years available for arrival dates from the Arnhem dataset), 11 years of male arrival dates collected on our study population, 9 years of moult onset dates and 3 years of departure dates and arrival at the wintering grounds.

a) Timing of breeding

Data for timing of breeding was available for 1959 (when the first pied flycatchers started to breed in our study area) and then from 1962 to 2015, but as mentioned above, we used data only from 1980 until 2015 because male arrival data (Arnhem) was available from that year onwards. Every year nest boxes were checked weekly from early March until late July and information on the progression of nest building and on the date when first eggs were laid was collected. Since eggs are laid in one-day intervals and pied flycatcher clutches typically have six eggs in the Netherlands, we had accurate measurements of individual laying dates. Pied flycatcher females typically incubate for 12-14 days, usually starting at the date when the last egg is laid, thus, after around 13 days of incubation, nests were inspected daily to identify the hatching dates of chicks. In some cases when we missed the actual hatching date, the date could be determined by ageing the chicks based on size and plumage development. When nestlings were seven days old, they were identified with a uniquely numbered aluminium ring and parents were also caught and identified.

b) Arrival dates

From 2005 until 2015, individual arrival date of males was assessed by daily scoring newly arriving males in our study area from early April onwards (Potti, 1998; Visser *et al.*, 2015; Both *et al.*, 2016). Birds choose a territory upon arrival and advertise their cavity or nest box to the females by singing continuously at or close to the potential nest site. Two or three trained observers walked independently pre-established routes covering the whole study area and visiting all boxes. Routes and direction of the routes were alternated daily among observers in order to prevent any potential bias among them. Detected birds were described in terms of plumage and aluminium/colour ring combinations. Male pied flycatchers display relatively large individual variation in plumage characteristic which, combined with colour

ring combinations, allows an initial recognition in the field without the need of capturing the birds. In our study site, plumage coloration varied from female-brown or light grey to almost entirely black with intermediates of increasing blackness (Drost, 1936). The forehead white patch also varied in size, from absent to a large patch covering most of the forehead, and also in shape from two distinct dots to a rectangular-shaped patch. We associated singing males to the closest nest-box in the vicinity. During the chick-rearing phase, those males were caught and described again in terms of plumage characteristics and ring combination. In most years (except 2005 and 2012) we also collected data on “bachelors”, *i.e.*, males still singing one week after the first eggs were found. These latter males were captured, identified and blood sampled and a few became breeding birds later in the season.

Apart from this more detailed dataset on individual arrivals (henceforth “observed arrival dataset”), we also had a second dataset from the Vogelwerkgroep Arnhem, a bird observation group that collected data on arrival dates of male pied flycatchers from 1980 to 2015, allowing the analyses of much longer term temporal shifts. This data collection occurred around 10 km from our study area and there was a high correlation (Pearson correlation coefficient = 0.83, 95% CI [0.45, 0.95], n=11 years; Supporting information Fig. S1) between their mean arrival data and the mean arrival dates of the observed arrival dataset. We are confident that this is a reliable and representative dataset for longer term trends in the studied pied flycatcher population, however, because the intercept was larger than 0 and the slope not equal 1 (slope: 0.52 ± 0.12 ; intercept: 6.73 ± 2.87 ; p -value < 0.01 , Fig. S1), we used a calculated arrival date based on the Arnhem dataset (henceforth “calculated arrival dataset”). We estimated the arrival dates of the males in the Hoge Veluwe National Park for the years we do not have data (before 2005) using the slope and intercept obtained in the relation between Arnhem and Hoge Veluwe arrival dates as follows: Calculated arrival = Intercept +

Slope * (Arnhem arrival). This calculated arrival date was then used in subsequent statistical analyses.

Female individual arrival date was not obtained directly as in the males' case. Instead, we used data from the start of nest building as a proxy, following the same procedure of Visser *et al.* (2015). Female pied flycatcher nest building reflects well their arrival dates as shown in Visser *et al.* (2015) since they choose a male and start nest building shortly after arriving (Dale *et al.*, 1992; Dale & Slagsvold, 1995). Nest building of individual pied flycatcher females was collected in our study area from 1980 to 2015, allowing the analysis of potential long-term shifts in annual timing of arrival of females.

c) Timing of moult

From 2005 to 2015 we had information on whether birds were moulting or not when they were caught for identification (when they had 7 days-old chicks). From 2009 to 2015 we also had information on moult score, *i.e.*, which feather was dropped and how much its replacement feather had grown. Finally, from 2013 to 2015, we had the actual date when the first feather was dropped for most of the breeding birds (Tomotani *et al.*, submitted). We could thus use the latter to define the accuracy of each type of measurement. We only used male data for this, since for most years we only had data on one or two moulting females. In 2007 and 2012 no information on moult was collected and thus we could not use these years in our analysis.

We used the R “moult” package (Erni *et al.*, 2013) to calculate the population average moult onset based on 2 analyses: (1) using presence/absence data (moulting/non-moulting individuals), we calculated the annual average starting date with a “probit model”, a generalized linear model with a binomial distribution and probit link function (Erni *et al.*, 2013). (2) Using the moult scores of different individuals (based on which feather was

missing or re-growing and how much it had grown) converted to a value of new feather mass grown (Dawson & Newton, 2004; Erni *et al.*, 2013) and regressing a line through the values of all individual new feather mass grown to obtain the population moult onset per year (with feather mass as response and date as explanatory variable). When we compared these values with the averages of the actual observed individual moult onsets, we noticed that the first presence/absence model provided a better estimate of moult onset than the feather mass regression (2013: observed = 76.93 ± 1.22 , presence/absence model = 73.09 ± 0.05 , feather mass regression = 59.22 ± 3.69 ; 2014: observed = 71.70 ± 1.15 , presence/absence model = 66.65 ± 0.04 , feather mass regression = 61.81 ± 2.30 ; 2015: observed = 74.42 ± 1.02 , presence/absence model = 71.14 ± 0.17 , feather mass regression = 62.81 ± 4.39). This was probably due to the fact that moult scores were mostly collected early in the season, with very few individuals with large moult scores, making the linear regression less reliable. It is also important to mention that we could use this “probit model” because we have experimental data showing that, within a year, the male moult onset is independent of the termination of breeding. Thus, in a given year, most individuals moult more or less on the same calendar day (Tomotani *et al.*, submitted.). Therefore, it is not so problematic that we have to use mid-to late breeding birds to calculate the populational moult onset in the present paper as the majority of early breeders would not be moulting when captured at the standardized chick-day 7. We thus used the 9 years of moult onset based on data of presence/absence of moulting birds, even in the case of the 3 years we had the exact onsets in order to have comparable data over years.

d) Departure dates and arrival at African wintering grounds

From 2013 to 2015, a subsample of the males (38 in 2013, 30 in 2014 and 30 in 2015) was equipped with light-level geolocators (Intigeo-W50, Migrate Technology Ltd, Cambridge,

UK) using a leg loop harness built with an 0.7 mm elastane cord and adjusted to each bird with a knot in the posterior mounting tube, between the device's mounting loops.

Geolocators were recovered in the subsequent years (2014 to 2016) when birds returned for breeding and 26 tracks (out of 98, 26%) could be analysed (12 from 2013, 7 from 2014 and 7 from 2015). Data was processed similarly to Åkesson *et al.*(2012), but due to the imprecision of latitude data, we only used the information collected for longitude (inferred from local solar noon/midnight). Twilight transitions were determined using TransEdit (British Antarctic Survey, Cambridge) with a single threshold value of five, minimum daylight periods of one hour and minimum night period of four hours. Positions were obtained using the software BirdTracker, which gave us two positions per day (noon and midnight). Data was then visually inspected to detect large changes from the study area longitude, indicating a departure from breeding grounds and then arrival at the wintering grounds, since pied flycatchers move to the west, following the African coast during autumn migration (Ouwehand *et al.*, 2016).

If the logger was still working upon recapture and a full track could be downloaded, we used the data corrected for clock drift, otherwise, clock drift effects on longitude were tested as described in (Ouwehand *et al.*, 2016). There was no noticeable clock drift effect (always <1 min).

e) Temperatures

Daily temperature values were collected from the Dutch meteorological institute database (KNMI - <https://www.knmi.nl/nederland-nu/klimatologie/>, accessed February 2016) for the Dutch temperatures and from the US National Oceanic and Atmospheric Administration database (NOAA - <ftp://ftp.ncdc.noaa.gov/pub/data/g sod/>, accessed February 2016) for

African temperatures. Temperatures from the NOAA database were converted from Fahrenheit were converted to Celsius to match the KNMI database.

For the Dutch weather variables, we used data from the Deelen weather station which is directly adjacent to the study area. For African temperature data, we used information on the pied flycatchers' wintering location using the data from the geolocators that we deployed and also from the literature (Ouwehand *et al.*, 2016) to identify the closest weather station from their wintering grounds. Pied flycatchers from the Netherlands winter in the Ivory Coast, where Daloa (6°27'W 6°53'N) is the closest weather station with a reasonable amount of data. Because the dataset still had large gaps we also used information from two other nearby locations (Gagnoa, 5°56'W 6°08'N, and Yammousoukro, 5°17'W 6°49'N). We still ended up with a few gaps in the data, that we interpolated using the average of data at the boundaries of the gap. This was seen as a minimal issue due to the way this data is used in the statistical analysis (see section "causes of variation in timing"). We obtained complete data from 1980 until 2015 for Dutch temperatures.

We also obtained data on photoperiodic variation of the Netherlands from the NOAA. We considered the civil twilight as the boundary of the effective light phase important for the birds (Gwinner, 1989). Because photoperiodic variation of the breeding and wintering grounds has a high correlation and vary in the same direction, it was not necessary to obtain and model day length data for Africa separately.

f) Adult survival

We used two datasets with data on individual capture histories; one with 11 years for which we had data on the average interval between breeding and moult for each year (2005 to 2015, 1252 individuals) and a second one with 35 years for which we had data on the average interval between arrival and breeding for each year (1980 to 2015, 3887 individuals).

Individuals were not included in the analyses when nestlings, but only when breeding for the first time. Thus, we did not include the nestling survival in this analysis. We used these datasets to estimate adult survival and recapture probability and whether the change in the two intervals (arrival-breeding and breeding-moult) had an effect on adult survival, while taking into account effects of sex, capture occasion (two categories: first or later) and age at first capture (three categories: (i) second calendar year birds (hereafter **SY**), ringed as nestling in the previous year; (ii) after second calendar year birds (**ASY**), ringed as nestling two or more years prior to the capture; and (iii) unknown age (**Unknown**), for the birds that were not ringed as nestlings and consequently their age (SY or ASY) could not be reliably determined.

Data analysis

All analyses were performed in R version 3.2.1 (R Core Team, 2015) and focused on the annual means of each annual cycle stage. To define the minimal models, we always used backwards model selection, dropping non-significant terms in each step. Survival analyses were performed with program MARK (White & Burnham, 1999).

a) Shifts in timing

We used simple and multiple regressions fitting year as linear and quadratic terms to test for shifts of the annual average value of each annual cycle component across years (including both longer-calculated and shorter-observed datasets for male arrival dates). Then to test whether there has been any change in the amount of days available between arrival, breeding and moult, we calculated the differences between arrival and breeding dates (arrival date of males and females and egg-laying dates) and between chick fledging date and moult onset. Predicted chick fledging was calculated from egg-laying dates by adding 6 days of egg laying, 12 days of incubation and 15 days of chick care until fledging. In all analyses

involving timing of moult, we used the standard errors of the estimates as weights (weight = $1 / se$) in weighted analysis to account for the prediction error of the moult dates.

We also tested whether the different slopes that we obtained from the separate regressions were significantly different from each other. We used multiple regression analyses with date as the response variable and the interaction of annual cycle stage (arrival, moult, egg-laying and hatching dates) and year (linear and quadratic) as the explanatory variables. We ran two separate analyses, one including all stages, starting in 1980 but with missing values for some stages in earlier years, and a second one only including those stages that we had complete data since 1980 (calculated arrival, egg-laying and hatching dates).

b) Causes of variation in timing

We tested whether the variation of each of the annual cycle's components (with the exception of hatching dates, which depend mostly on egg-laying dates) could be explained by variation in temperature cues alone or in interaction with changes in day-length. As in other analyses we also used the annual means of each stage here. Following the method described in Gienapp *et al.* (2005), we used proportional hazard models (Cox, 1992) implemented in the R “survival” package (Therneau, 2015) to model the relationship between the climatic event and the occurrence of the event. Proportional hazard models calculate the daily probability of an event to occur. They, therefore, allow including time-dependent variables, i.e. variables that change their value during the time an individual is ‘at risk’. Modelling effects of weather variables on annual cycle stages, as arrival date, is biologically more realistic than using fixed time windows over which these variables are averaged (Gienapp *et al.*, 2005). The value of this time-dependent weather variable at day t was calculated as the average over periods of various lengths (5, 10, 15, 20, 25 or 30 days) ending at day t (see Supporting information Table S3 for more details). For African weather variables, we also used lagged shifting

Accepted Article

windows of the same length (of 20 days) but ending 20, 40, 60 or 80 days before day t (see Table S3). We defined intervals that we deemed biologically significant; thus African temperatures were only tested for arrival dates because it is unlikely that African temperatures would have affected the egg-laying dates and moult onset, as the birds were already in the Netherlands. Therefore, in total, we compared 24 possible combinations for arrival dates, 5 for egg-laying dates and 5 for moult (Table S3). Identification of the best period was done by comparing the log-likelihoods of the different models, however, their close values made such selection challenging in some cases (Table S3). After the best window was defined for each stage, we tested what temperatures significantly explained the variation in timing of different stages using different proportional hazard models for each annual cycle stage (compared with likelihood ratio tests). We fitted African temperatures (with and without lag), Dutch temperatures and the interactions of day-length and Dutch temperature and day-length and lagged African temperature, depending on the stage. The temperatures selected in the best models for each stage were also fitted against year as a trend (linear and quadratic) to test whether these temperatures also changed across years.

c) Consequences of variation in timing

We used multiple regressions to test whether the intervals between the arrival and breeding dates and breeding date and moult onset explained fitness components. We looked at two components related to breeding success (the annual average clutch size and the annual proportion of fledged chicks that recruited per year) and at adult survival.

We used generalised linear models with binomial (Bernoulli) response and logit link function to test whether the annual proportion of fledged chicks that recruited was explained by either the annual difference of the interval between arrival and breeding or the interval between breeding and moult. For the interval between arrival and breeding, we only used the longer

calculated arrival dataset, since it correlated with the shorter (Hoge Veluwe) dataset, showing a similar pattern, but included many more years. Clutch size was similarly analysed using multiple regressions testing for year and interval effects.

To analyse adult survival we performed a Cormack-Jolly-Seber (CJS) mark-recapture analysis using the software MARK (White & Burnham, 1999). The CJS model estimates annual local survival probabilities (Φ) based on live recaptures only, while controlling for capture probability (P). In this analysis, we used Akaike's Information Criterion (AICc) for the model selection and goodness of fit was tested using the bootstrap procedure in the MARK software. We first defined the best model including time ("year" as a factor), capture occasion (first or later), sex (male or female) and age at first capture (SY, ASY or unknown), for both survival and recapture probabilities. In the most complex models, birds that were in their SY or of unknown age at their first capture moved to the ASY age class in the following year and remained in this age class for the rest of the years. Birds first caught in their ASY never moved to another age class in subsequent captures. We could not use plumage characters as ageing criteria because they were not deemed as precise enough and also not always recorded for all captured individuals. On the other hand, in the simplified models that did not include the age at first capture, the capture occasion variable only explained differences between the first or later captures of the same individual independent on their age. We first fitted the model with the interaction between time, capture occasion and the interaction between sex and age at first capture and then used backwards comparison to define the best model. After the best model was defined we replaced the time variable by the interval between arrival and breeding (longer dataset) or breeding and moult (shorter dataset) and compared these new models with the best model to investigate specifically whether variation in survival among years could, in fact, be explained by variation in these intervals.

Results

Shifts in timing

Annual means of male and female arrival (nest building) date, egg-laying date and date of moult onset shifted at different rates, as shown by the significant interaction between stage and year both when all stages were considered ($F_{5,152} = 6.74$, p -value <0.01 ; Supporting information Table S1) and also when we only included stages for which we had data since 1980 ($F_{3,136} = 8.12$, p -value <0.01 ; Table S1). There were no significant advancements for male arrival (calculated arrival, estimate = -0.03 ± 0.03 Table S1, linear slope not significant in post hoc test, Supporting information Table S2) and male moult onset advanced faster than the other stages (estimate = -1.11 ± 0.29 ; significant slope $F_{1,7} = 12.90$, p -value = 0.01, Tables 1, S1 and S2), both in the long and short-term dataset. The rates of advancements for (observed) arrival dates of the males, arrival date of the females, egg-laying dates and hatching dates were strikingly similar (observed male arrival = -0.32 ± 0.28 , no significant slope; arrival females = -0.30 ± 0.05 , egg-laying = -0.30 ± 0.05 , hatching = -0.30 ± 0.05 , significant slopes $F_{1,34} = 29.55$, $F_{1,34} = 48.8$, $F_{1,34} = 29.72$, respectively, p -value <0.01 , Tables 1 and S2).

Because arrival date, egg-laying dates and moult dates shifted unequally, we calculated the interval between each stage and tested whether they changed across years. As expected from the previous analyses, the intervals between male arrival and breeding and between breeding and male moult changed across years. In contrast, the interval between female arrival (nest building) and egg-laying dates did not significantly change since 1980 (Table S2, Fig. 1e). The interval between male (calculated) arrival and egg-laying date changed non-linearly: it became shorter until around 2008 and then started to increase again (estimate for the quadratic term = 0.01 ± 0.01 , $F_{1,33} = 7.86$, p -value <0.01 ; Table S2). A post hoc broken stick analysis suggested that two separate regressions, one before 2008 and one after 2008, had a

Accepted Article

better fit than the quadratic model (adjusted R^2 for quadratic term = 0.53, adjusted R^2 for two regressions before and after 2008 = 0.59). Dividing the dataset before and after this year and testing separately we found that these two regressions were significant with distinct slopes (First: slope = -0.42 ± 0.07 , $F_{1,26} = 37.11$, p -value < 0.01 . Second: slope = 0.60 ± 0.20 , $F_{1,6} = 9.08$, p -value = 0.02) (Fig. 1e). Finally, the interval between calculated fledging date (termination of breeding) and date of male moult onset significantly decreased across years. As a consequence, there was a larger overlap between moult and breeding in recent years (slope = -1.38 ± 0.26 , $F_{1,7} = 27.87$, p -value < 0.01 ; Table S2, Fig. 1e).

Annual average departure dates from the breeding grounds varied between the 7th and 11th of August (with a significant advancement across years using a weak test that included the three years with available data; see Fig. 1, Table S2) and annual mean arrival dates in Africa varied between the 6th and 19th of September. Average wintering longitude was -7.46 degrees (range: -9.71 to -4.86 , 26 loggers), suggesting that this pied flycatcher population winters in Ivory Coast similarly to what was reported for other Dutch populations (Ouwehand *et al.*, 2016).

Causes of variation in timing

Variation in annual male arrival date was not explained by temperatures, moreover, these temperatures also did not significantly change since 1980 (Supporting information Tables S4, S5, Fig. S2).

Variation in annual female arrival (nest building) and egg-laying dates, on the other hand, were significantly explained by Dutch temperatures (arrival: coefficient = 0.23 ± 0.07 , $\chi^2 = 10.26$, p -value = 0.01; egg-laying: coefficient = 0.58 ± 0.12 , $\chi^2 = 25.89$, p -value < 0.01 ; Tables 1 and S4), with higher temperatures being related to earlier arrival and egg-laying. However, while the temperatures important for egg-laying significantly changed across years (estimate = 0.05 ± 0.07 , $F_{1,34} = 4.41$, p -value = 0.04), this was not the case for those important for arrival

(Table S5, Fig. S2). A post hoc test for the egg-laying date data, in which we included year and temperature in the same model to correct for the trends of temperature and year in the same direction, resulted in both temperature and year as significant.

Finally, variation in annual male moult onset was not related to Dutch temperatures (Tables 1 and S4) thus the advancement observed for the moult onset cannot be related to any recent increase in Dutch temperatures.

Consequences of variation in timing

There was no effect of the interval between arrival and breeding and breeding and moult on clutch size. Clutch size, however changed across years since 1980 in a non-linear way, increasing until around 2003 and later on decreasing again (quadratic estimate = -0.001 ± 0.0005 , $F_{1,33} = 9.50$, p -value < 0.01 ; Supporting information Table S6). A post hoc broken stick analysis suggests that two regressions before and after 2008 have a better fit than the model with the quadratic term (adjusted R^2 for quadratic term = 0.40, adjusted R^2 for two regressions before and after 2008 = 0.48). Dividing the dataset before and after this year and testing them separately we found that the first regression was significant (slope = 0.03 ± 0.01 , $F_{1,26} = 17.43$, p -value < 0.01), but the second was not (n.s. slope = -1.03 ± 0.05 , $F_{1,6} = 4.90$, p -value = 0.07) (Fig.2a).

The proportion of fledged chicks that recruited also changed non-linearly across years (quadratic estimate = -0.001 ± 0.0002 , $\chi^2 = 10.49$, p -value < 0.01 ; Fig.2b, Table S6), but it was also significantly related to the interval between arrival and breeding, even when year as a continuous term was included in the same model (Table S6). A higher proportion of chicks fledged when this interval was shorter (slope = -0.04 ± 0.01 , $\chi^2 = 6.99$, p -value = 0.01; Fig. 2c, Table S6). A shorter interval is also related to earlier egg-laying dates of the females, and earlier laying was significantly related to a higher proportion of fledged chicks that recruited

(slope = -0.03 ± 0.02 , $\chi^2 = 4.04$, p -value = 0.04; Fig. 2d, Table S6). Thus the improved chick recruitment may be related to the earlier breeding.

Finally, the proportion of fledged chicks was affected by the interval between breeding and moult, but the relation was non-linear (quadratic estimate = -0.02 ± 0.01 , $\chi^2 = 5.40$, p -value = 0.02; Table S6).

We did not observe a trend in adult survival across years and neither the interval between arrival and breeding nor breeding and moult explained the variation in adult survival. When analysing adult survival probability (Φ), none of the models including the interval between arrival and breeding and almost none containing the interval between breeding and moult was among the best models (delta AICc always larger than 2; Table 2). Even those models containing the breeding/moult interval, with a delta AICc smaller than 2, however, had negligible effects of this interval (beta estimate in the model only including that interval = 0.0001 ± 0.0008).

Discussion

Our results show that climate change affected the annual cycle stages of the European pied flycatcher, making them shift at different rates from one another. This was probably due to the different changes in the various temperatures that correlate with the phenology of these annual cycle stages. Although timing of moult advanced rapidly, we could not detect a correlating temperature period to explain this shift. Our results also suggest that the advancement of breeding allows more time for events such as fledgling development, which could explain at least in part the increase in the proportion of fledged chicks that recruited when the breeding – moult interval got longer. The shortening of the interval between arrival and breeding and the larger moult/breeding overlap would be expected to incur fitness costs, but we did not detect effects on adult survival. Therefore, at least for the aspects studied so

far, climate change has actually led to an improvement in breeding conditions for this species, potentially by allowing more time for fledgling development (Tomotani *et al.*, 2016).

Other consequences of unequal shifts

While costs of shortening the intervals between stages could not be detected in terms of adult survival, it is still possible that costs are present in other aspects we have not investigated. Females, for instance, may suffer costs that are much more subtle. For example, a decrease in the number of days between arrival and breeding may also mean that females need to be much faster in choosing a male (Dale *et al.*, 1992; Dale & Slagsvold, 1995), building the nest and initiation of egg-laying. Indeed, in recent years male and female arrival are happening almost at the same dates, so when the first females arrive, a good number of males are not settled and territories are still being claimed (Alatalo *et al.*, 1984). This could, for example, result in a reduced time to assess male quality (Alatalo *et al.*, 1984) and increase the probability of a female to pair with a male that already has another female. Polygyny is costly for females since the number of unfertilized eggs and chick mortality are higher when a male has more than one female (Lubjuhn *et al.*, 2000). This could be problematic for late-arriving females, as they would supposedly be even more constrained by the increased competition with other females late in the season (Dale *et al.*, 1992). This shorter time span could also decrease the females' body condition and in the long run be detrimental to their breeding success, since egg-laying is particularly costly (Visser & Lessells, 2001). This may have been reflected in the previously increasing clutch size trend, which changed in recent years (although there was no significant decline detected by the broken stick analysis) (Fig. 2a).

The shifts observed in the timing of moult in males may also result in a shortening of the total time available for breeding, similarly to what was reported by Moyes *et al.* (2011). Males advanced their moult onset at a higher rate than the breeding dates, which suggests they are

regressing their gonads earlier as well. Experimental studies in captive starlings (*Sturnus vulgaris*) show that onset of moult is related to the gonadal regression in males (Dawson, 2006); moreover, both gonadal regression and moult advanced in great tits (*Parus major*) experimentally exposed to higher spring temperatures in captivity (Visser *et al.*, 2011). In the present study, there was no effect of temperatures on moult onset, but regardless of the factor that is causing advancements in the timing of moult, an uncoupling between onset and termination of breeding is possible.

Underlying causes of unequal shifts

A curious outcome of our analysis is the faster advancement of moult in relation to breeding. Moult onset in males seems to be determined earlier in the season and not determined by the termination of breeding (Tomotani *et al.*, submitted). If moult is not connected to timing of breeding, it could advance independently. Another possibility is that moult is set when the individual is born. It is known, for example, that the photoperiod when the animal is born can modify the timing of events later in life (Lee & Zucker, 1988; Coppack *et al.*, 2001; Coppack & Pulido, 2009). In particular, it has been proposed that advancements in birth date (as laying dates advance) could modify the timing of migration at the population level (Both, 2010; Gill *et al.*, 2014). If the timing of moult is set when the birds are born, this may explain the observed change in moult onset at the population level (but see Larsson, 1996).

If the dates when birds are born affect both their arrival and moult, we should also observe an advancement in arrival dates, but it was not the case in our study population. Other migratory species, however, have been reported to advance their arrival dates (Usui *et al.*, 2016) and also other populations of pied flycatcher (Ahola *et al.*, 2004; Both *et al.*, 2016; Valtonen *et al.*, 2016). Arrival dates, in comparison to the onset of moult, are much more susceptible to modulations by environmental conditions, such as weather, body condition and wind patterns

(Ahola *et al.*, 2004; Erni *et al.*, 2005; Sinelschikova *et al.*, 2007; Bauer *et al.*, 2008; Eikenaar & Schmaljohann, 2014; Both *et al.*, 2016; Teitelbaum *et al.*, 2016). Moreover, arrival dates can usually only be assigned to individuals that survived the migration and thus there might be a bias if early arrival increases mortality (Brown & Brown, 2000). In comparison to arrival, moult is a relatively cleaner expression of the individuals (endogenous) timing (*i.e.* more directly assessed) (Gwinner, 1996), thus potential changes related to birth effects are supposedly more detectable in timing of moult than timing of arrival.

None of the temperatures important for timing of stages directionally and significantly changed over the past years, with the exception of those explaining the variation in egg-laying dates (Fig. S2). We would expect that the temperatures important for female arrival (nest building) would also have changed since the stage advanced at the same rate as egg-laying dates. However, while the Dutch temperatures that are important for female arrival (nest building) and breeding are almost the same and partly overlap (Table 1), the window size of the temperature important for laying date is 10 days larger. One possibility is that the temperatures important for female arrival (nest building) and egg laying are nearly the same and trends are not detected for the arrival temperatures due to a larger variation of temperatures in the pre-laying period especially in recent years (Fig. S2).

In the present study, while we included multiple annual cycle stages, we were limited to the stages that occur on the breeding grounds – for which we have long-term data available. Even so, there was one stage at the breeding ground that we could not include: the timing of autumn migration. Date of departure in late summer seems to be correlated with the timing of egg-laying, but not so much with the timing of chick hatching (Tomotani *et al.*, submitted). Departure dates also seem to be variable across years. Thus, it is possible that, together with the advancement of breeding and moult, the birds are now also departing earlier. On the other hand, if there is no change in timing of migration, males have more time to moult. It has been

reported that conditions late in the season may improve for some birds, such as for some short-distance migrants (Jenni & Kery, 2003). For now we can only speculate how climate change has affected the timing of autumn migration in our population.

Another aspect that we are unable to assess is the timing of stages at the wintering grounds in Africa. It is unknown if birds remain time constrained at their wintering grounds or if the wintering grounds serve as “time buffers”. Early arrival at the wintering grounds may, for example, be important to secure resources, which could result in improved body reserves that carry-over to the next season (for example in terms of earlier arrival and/or improved individual quality). This is the case for American redstarts (*Setophaga ruticilla*; Marra *et al.*, 1998, 2015a; Norris *et al.*, 2004). In pied flycatchers, an experimental delay of hatching dates imposed a larger moult-breeding overlap and made males from this group winter at a different location than controls and advanced males (Tomotani *et al.*, submitted). This suggests that the selection of wintering territory also depends on what birds experience at the breeding grounds, but whether wintering will later impact breeding remains unknown (Ouwehand & Both, 2017).

Final remarks

Climate change unequally affects the annual cycle stages of bird (Van der Jeugd *et al.*, 2009; Eichhorn *et al.*, 2010; Valtonen *et al.*, 2016) and mammal species (Ozgul *et al.*, 2010; Moyes *et al.*, 2011). Such shifts may lead to positive fitness consequences in some cases, for example, in marmots` offspring that now have more time. But it may also have negative consequences, in the case of the red deer in which not only a mismatch between male and female timing happened, but also a shortening in their breeding window. In the present study, there were also positive and negative fitness consequences of unequal shifts, but they depended on the sex or life stage of the animal. In pied flycatchers, climate change seems to

benefit males and offspring, which have more time available due to the advancement of breeding, but it can also be costly for the females. The impacts of climate change, thus, are not only different for distinct trophic levels (Parmesan, 2006), but also for stages (*e.g.* breeding, moult) sex and life-stages (*e.g.* males, females, offspring) within the same species. It is very possible that such patterns are widespread, especially among organisms with complex annual cycles, meriting a careful look as new and more complete datasets covering the full annual cycle become available for other populations and species.

For a broader understanding of the ecological consequences of climate change, different stages of the annual cycle should be considered (Crozier *et al.*, 2008; Yang & Rudolf, 2010), in particular for organisms with complex cycles, such as migratory birds (Small-Lorenz *et al.*, 2013; Marra *et al.*, 2015b). Our knowledge on climate change impacts on organisms will thus greatly benefit from continued standardised data collection that includes more than one stage.

Acknowledgements

We are grateful to the board of the National Park “De Hoge Veluwe” for the permission to conduct our research on their property. This study was only possible due to the constant effort of multiple researchers, field assistants and students in collecting the field information and managing the database. We also thank all the members of the Vogelwerkgroep Arnhem reporting their phenology data. Finally, we thank Ally Phillimore and three anonymous reviewers for their insightful comments on a previous version of the manuscript. BMT received a doctorate grant from CNPq (237790/2012-2) and a small research grant from Ecology Fonds - KNAW (UPS/375/ECO/J1534).

References

- Ahola M, Laaksonen T, Sippola K, Eeva T, Rainio K, Lehikoinen E (2004) Variation in climate warming along the migration route uncouples arrival and breeding dates. *Global Change Biology*, **10**, 1610–1617.
- Åkesson S, Klaassen R, Holmgren J, Fox JW, Hedenström A (2012) Migration Routes and Strategies in a Highly Aerial Migrant, the Common Swift *Apus apus*, Revealed by Light-Level Geolocators. *PLoS ONE*, **7**, e41195.
- Alatalo RV, Lundberg A, Ståhlbrandt K (1984) Female mate choice in the Pied Flycatcher *Ficedula hypoleuca*. *Behavioral Ecology and Sociobiology*, **14**, 253–261.
- Bauer S, Gienapp P, Madsen J (2008) The Relevance of Environmental Conditions for Departure Decision Changes En Route in Migrating Geese. *Ecology*, **89**, 1953–1960.
- Both C (2010) Flexibility of Timing of Avian Migration to Climate Change Masked by Environmental Constraints En Route. *Current Biology*, **20**, 243–248.
- Both C, te Marvelde L (2007) Climate change and timing of avian breeding and migration throughout Europe. *Climate Research*, **35**, 93–105.
- Both C, Visser ME (2001) Adjustment to climate change is constrained by arrival date in a long-distance migrant bird. *Nature*, **411**, 296–298.
- Both C, Artemyev AV, Blaauw B et al. (2004) Large-scale geographical variation confirms that climate change causes birds to lay earlier. *Proceedings of the Royal Society of London B: Biological Sciences*, **271**, 1657–1662.
- Both C, Turnhout CAMV, Bijlsma RG, Siepel H, Strien AJV, Foppen RPB (2009) Avian population consequences of climate change are most severe for long-distance migrants in seasonal habitats. *Proceedings of the Royal Society of London B: Biological Sciences*, **277**, 1259–1266.

- Both C, Bijlsma RG, Ouweland J (2016) Repeatability in Spring Arrival Dates in Pied Flycatchers Varies Among Years and Sexes. *Ardea*, **104**, 3–21.
- Brown CR, Brown MB (2000) Weather-Mediated Natural Selection on Arrival Time in Cliff Swallows (*Petrochelidon pyrrhonota*). *Behavioral Ecology and Sociobiology*, **47**, 339–345.
- Brown JL, Li S-H, Bhagabati N (1999) Long-term trend toward earlier breeding in an American bird: A response to global warming? *Proceedings of the National Academy of Sciences*, **96**, 5565–5569.
- Charmantier A, Gienapp P (2014) Climate change and timing of avian breeding and migration: evolutionary versus plastic changes. *Evolutionary Applications*, **7**, 15–28.
- Coppack T, Pulido F (2009) Proximate control and adaptive potential of protandrous migration in birds. *Integrative and Comparative Biology*, **49**, 493–506.
- Coppack T, Pulido F, Berthold P (2001) Photoperiodic response to early hatching in a migratory bird species. *Oecologia*, **128**, 181–186.
- Cox DR (1992) Regression models and life-tables. In: *Breakthroughs in statistics*. (eds Kotz S, Johnson NL), pp. 527–541. Springer New York, New York.
- Crick HQP, Dudley C, Glue DE, Thomson DL (1997) UK birds are laying eggs earlier. *Nature*, **388**, 526–526.
- Crozier LG, Hendry AP, Lawson PW et al. (2008) Potential responses to climate change in organisms with complex life histories: evolution and plasticity in Pacific salmon. *Evolutionary Applications*, **1**, 252–270.
- Dale S, Slagsvold T (1995) Female Contests for Nest Sites and Mates in the Pied Flycatcher *Ficedula hypoleuca*. *Ethology*, **99**, 209–222.
- Dale S, Rinden H, Slagsvold T (1992) Competition for a mate restricts mate search of female pied flycatchers. *Behavioral Ecology and Sociobiology*, **30**, 165–176.

- Dawson A (2006) Control of molt in birds: Association with prolactin and gonadal regression in starlings. *General and Comparative Endocrinology*, **147**, 314–322.
- Dawson A, Newton I (2004) Use and Validation of a Molt Score Index Corrected for Primary-Feather Mass. *The Auk*, **121**, 372–379.
- Drost R (1936) Ueber das Brutklei männlicher Trauerfliegenfänger, *Muscicapa hypoleuca*. Nach Untersuchungen an Helgoländer Zugvögeln und an nord- und mitteleuropäischen Stücken. *Der Vogelzug*, **7**, 179–186.
- Easterling DR, Horton B, Jones PD et al. (1997) Maximum and Minimum Temperature Trends for the Globe. *Science*, **277**, 364–367.
- Eichhorn G, van der Jeugd HP, Meijer HAJ, Drent RH (2010) Fueling Incubation: Differential Use of Body Stores in Arctic- and Temperate-breeding Barnacle Geese (*Branta leucopsis*). *The Auk*, **127**, 162–172.
- Eikenaar C, Schmaljohann H (2014) Wind conditions experienced during the day predict nocturnal restlessness in a migratory songbird. *Ibis*, **157**, 125–132.
- Erni B, Liechti F, Bruderer B (2005) The role of wind in passerine autumn migration between Europe and Africa. *Behavioral Ecology*, **16**, 732–740.
- Erni B, Bonnevie BT, Oschadleus H-D, Altwegg R, Underhill LG (2013) moult: An R Package to Analyze Moults in Birds. *Journal of Statistical Software*, **52**, 1–23.
- Gienapp P, Hemerik L, Visser ME (2005) A new statistical tool to predict phenology under climate change scenarios. *Global Change Biology*, **11**, 600–606.
- Gienapp P, Lof M, Reed TE, McNamara J, Verhulst S, Visser ME (2013) Predicting demographically sustainable rates of adaptation: can great tit breeding time keep pace with climate change? *Philosophical Transactions of the Royal Society B: Biological Sciences*, **368**, 20120289.

- Gill JA, Alves JA, Sutherland WJ, Appleton GF, Potts PM, Gunnarsson TG (2014) Why is timing of bird migration advancing when individuals are not? *Proceedings of the Royal Society of London B: Biological Sciences*, **281**, 20132161.
- Gwinner E (1989) Photoperiod as a Modifying and Limiting Factor in the Expression of Avian Circannual Rhythms. *Journal of Biological Rhythms*, **4**, 125–138.
- Gwinner E (1996) Circannual clocks in avian reproduction and migration. *Ibis*, **138**, 47–63.
- Hemborg C, Lundberg A (1998) Costs of Overlapping Reproduction and Moulting in Passerine Birds: An Experiment with the Pied Flycatcher. *Behavioral Ecology and Sociobiology*, **43**, 19–23.
- Jacobs JD, Wingfield JC (2000) Endocrine control of life-cycle stages: a constraint on response to the environment? *The Condor*, **102**, 35–51.
- Jenni L, Kery M (2003) Timing of autumn bird migration under climate change: advances in long-distance migrants, delays in short-distance migrants. *Proceedings of the Royal Society B-Biological Sciences*, **270**, 1467–1471.
- Larsson K (1996) Genetic and environmental effects on the timing of wing moult in the barnacle goose. *Heredity*, **76**, 100–107.
- Lee TM, Zucker I (1988) Vole infant development is influenced perinatally by maternal photoperiodic history. *The American Journal of Physiology*, **255**, R831–838.
- Lubjuhn T, Winkel W, Epplen JT, Brün J (2000) Reproductive Success of Monogamous and Polygynous Pied Flycatchers (*Ficedula hypoleuca*). *Behavioral Ecology and Sociobiology*, **48**, 12–17.
- Marra PP, Hobson KA, Holmes RT (1998) Linking Winter and Summer Events in a Migratory Bird by Using Stable-Carbon Isotopes. *Science*, **282**, 1884–1886.

- Marra PP, Studds CE, Wilson S, Sillett TS, Sherry TW, Holmes RT (2015a) Non-breeding season habitat quality mediates the strength of density-dependence for a migratory bird. *Proceedings of the Royal Society B: Biological Sciences*, **282**, 20150624.
- Marra PP, Cohen EB, Loss SR, Rutter JE, Tonra CM (2015b) A call for full annual cycle research in animal ecology. *Biology Letters*, **11**, 20150552.
- Morrison CA, Baillie SR, Clark JA, Johnston A, Leech DI, Robinson RA (2015) Flexibility in the timing of post-breeding moult in passerines in the UK. *Ibis*, **157**, 340–350.
- Moyes K, Nussey DH, Clements MN et al. (2011) Advancing breeding phenology in response to environmental change in a wild red deer population. *Global Change Biology*, **17**, 2455–2469.
- Norris DR, Marra PP, Kyser TK, Sherry TW, Ratcliffe LM (2004) Tropical winter habitat limits reproductive success on the temperate breeding grounds in a migratory bird. *Proceedings of the Royal Society of London B: Biological Sciences*, **271**, 59–64.
- Ouwehand J, Both C (2017) African departure rather than migration speed determines variation in spring arrival in pied flycatchers. *Journal of Animal Ecology*, **86**, 88–97.
- Ouwehand J, Ahola MP, Ausems ANMA et al. (2016) Light-level geolocators reveal migratory connectivity in European populations of pied flycatchers *Ficedula hypoleuca*. *Journal of Avian Biology*, **47**, 69–83.
- Ozgul A, Childs DZ, Oli MK et al. (2010) Coupled dynamics of body mass and population growth in response to environmental change. *Nature*, **466**, 482–485.
- Parnesan C (2006) Ecological and Evolutionary Responses to Recent Climate Change. *Annual Review of Ecology, Evolution, and Systematics*, **37**, 637–669.
- Parnesan C (2007) Influences of species, latitudes and methodologies on estimates of phenological response to global warming. *Global Change Biology*, **13**, 1860–1872.

- Parmesan C, Yohe G (2003) A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, **421**, 37–42.
- Phillimore AB, Leech DI, Pearce-Higgins JW, Hadfield JD (2016) Passerines may be sufficiently plastic to track temperature-mediated shifts in optimum lay date. *Global Change Biology*, **22**, 3259–3272.
- Plard F, Gaillard J-M, Coulson T, Hewison AJM, Delorme D, Warnant C, Bonenfant C (2014) Mismatch Between Birth Date and Vegetation Phenology Slows the Demography of Roe Deer. *PLOS Biology*, **12**, e1001828.
- Potti J (1998) Arrival Time from Spring Migration in Male Pied Flycatchers: Individual Consistency and Familial Resemblance. *The Condor*, **100**, 702–708.
- R Core Team (2015) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Reed TE, Jenouvrier S, Visser ME (2013a) Phenological mismatch strongly affects individual fitness but not population demography in a woodland passerine. *Journal of Animal Ecology*, **82**, 131–144.
- Reed TE, Grøtan V, Jenouvrier S, Sæther B-E, Visser ME (2013b) Population Growth in a Wild Bird Is Buffered Against Phenological Mismatch. *Science*, **340**, 488–491.
- Serreze MC, Francis JA (2006) The Arctic Amplification Debate. *Climatic Change*, **76**, 241–264.
- Sinelschikova A, Kosarev V, Panov I, Baushev AN (2007) The influence of wind conditions in Europe on the advance in timing of the spring migration of the song thrush (*Turdus philomelos*) in the south-east Baltic region. *International Journal of Biometeorology*, **51**, 431–440.
- Small-Lorenz SL, Culp LA, Ryder TB, Will TC, Marra PP (2013) A blind spot in climate change vulnerability assessments. *Nature Climate Change*, **3**, 91–93.

- Accepted Article
- Stocker TF, Qin D, Plattner GK et al. (2013) *IPCC, 2013: climate change 2013: the physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*. Cambridge University Press, Cambridge.
- Stoks R, Geerts AN, De Meester L (2014) Evolutionary and plastic responses of freshwater invertebrates to climate change: realized patterns and future potential. *Evolutionary Applications*, **7**, 42–55.
- Teitelbaum CS, Converse SJ, Fagan WF, Böhning-Gaese K, O’Hara RB, Lacy AE, Mueller T (2016) Experience drives innovation of new migration patterns of whooping cranes in response to global change. *Nature Communications*, **7**, 12793.
- Thackeray SJ, Henrys PA, Hemming D et al. (2016) Phenological sensitivity to climate across taxa and trophic levels. *Nature*, **535**, 241–245.
- Therneau TM (2015) *A Package for Survival Analysis in S*.
- Tomotani BM, Gienapp P, Beersma DGM, Visser ME (2016) Climate change relaxes the time constraints for late-born offspring in a long-distance migrant. *Proceedings of the Royal Society of London B: Biological Sciences*, **283**, 20161366.
- Usui T, Butchart SHM, Phillimore AB (2016) Temporal shifts and temperature sensitivity of avian spring migratory phenology: A phylogenetic meta-analysis. *Journal of Animal Ecology*, **82**, 250–261.
- Valtonen A, Latja R, Leinonen R, Pöysä H (2016) Arrival and onset of breeding of three passerine birds in Eastern Finland tracks climatic variation and phenology of insects. *Journal of Avian Biology*, n/a-n/a.
- Van der Jeugd HP, Eichhorn G, Litvin KE, Stahl J, Larsson K, Van Der Graaf AJ, Drent RH (2009) Keeping up with early springs: rapid range expansion in an avian herbivore incurs a mismatch between reproductive timing and food supply. *Global Change Biology*, **15**, 1057–1071.

- Accepted Article
- Visser ME (2008) Keeping up with a warming world; assessing the rate of adaptation to climate change. *Proceedings of the Royal Society of London B: Biological Sciences*, **275**, 649–659.
- Visser ME (2016) Phenology: Interactions of climate change and species. *Nature*, **535**, 236–237.
- Visser ME, Both C (2005) Shifts in phenology due to global climate change: the need for a yardstick. *Proceedings of the Royal Society of London B: Biological Sciences*, **272**, 2561–2569.
- Visser ME, Lessells CM (2001) The costs of egg production and incubation in great tits (*Parus major*). *Proceedings of the Royal Society of London B: Biological Sciences*, **268**, 1271–1277.
- Visser ME, Holleman LJM, Gienapp P (2006) Shifts in caterpillar biomass phenology due to climate change and its impact on the breeding biology of an insectivorous bird. *Oecologia*, **147**, 164–172.
- Visser ME, Schaper SV, Holleman LJM, Dawson A, Sharp P, Gienapp P, Caro SP (2011) Genetic variation in cue sensitivity involved in avian timing of reproduction. *Functional Ecology*, **25**, 868–877.
- Visser ME, Gienapp P, Husby A, Morrissey M, de la Hera I, Pulido F, Both C (2015) Effects of Spring Temperatures on the Strength of Selection on Timing of Reproduction in a Long-Distance Migratory Bird. *PLoS Biol*, **13**, e1002120.
- Vose RS, Easterling DR, Gleason B (2005) Maximum and minimum temperature trends for the globe: An update through 2004. *Geophysical Research Letters*, **32**, L23822.
- Walther G-R, Post E, Convey P et al. (2002) Ecological responses to recent climate change. *Nature*, **416**, 389–395.

White GC, Burnham KP (1999) Program MARK: survival estimation from populations of marked animals. *Bird study*, **46**, S120–S139.

Wingfield JC (2008) Organization of vertebrate annual cycles: implications for control mechanisms. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **363**, 425–441.

Yang LH, Rudolf VHW (2010) Phenology, ontogeny and the effects of climate change on the timing of species interactions. *Ecology Letters*, **13**, 1–10.

Zimova M, Mills LS, Nowak JJ (2016) High fitness costs of climate change-induced camouflage mismatch. *Ecology Letters*, **19**, 299–307.

Tables

Table 1: Direction of the significant effects of Dutch and African temperatures on the annual cycle stages of pied flycatchers (red: negative, blue: positive). The values correspond to the coefficients obtained with a cox-proportional hazards model (see Table S4 for the full results), thus high positive values indicate the earlier occurrence of the event with higher temperatures (a higher hazard for the event to occur).

Annual-cycle stage	Sex	Years	Average date	Linear trend	Day length: Temp. NL	Temp. NL	Day length: Temp. Africa (lag)	Temp. Africa (lag)	Temp. Africa (no lag)	Window size (no lag) / Window place (lag Africa)
Arrival (observed)	M	2005-2015	19-Apr	-0.32 (±0.20) n.s.	0.01 (±0.02) n.s.	0.16 (±0.11) n.s.	-0.01 (±0.01) n.s.	-0.77 (±0.50) n.s.	0.9 (±0.49) n.s.	5 days / 10-Mar
Arrival (Calculated)	M	1980-2015	20-Apr	-0.03 (±0.05) n.s.	0.004 (±0.02) n.s.	0.25 (±0.10)	-0.001 (±0.01) n.s.	0.21 (±0.29) n.s.	0.76 (±0.50) n.s.	15 days / 11-Mar
Arrival	F	1980-2015	01-May	-0.30 (±0.06)	-0.01 (±0.01) n.s.	0.23 (±0.07)	0.004 (±0.01) n.s.	0.33 (±0.22) n.s.	-0.18 (±0.38) n.s.	15 days / 22-Mar
Egg laying	F	1959-2015	11-May	-0.30 (±0.04)	0.001 (±0.01) n.s.	0.58 (±0.12)				25 days
Moult	M	2005-2015	13-Jun	-1.11 (±0.29)	-0.08 (±0.06) n.s.	-0.26 (±0.31) n.s.				30 days

Table 2: The top five best models obtained from the mark-recapture analysis that analyzed adult survival in relation to the interval between arrival and breeding and breeding and moult, for both the long and short datasets. In all cases, the best models included the six groups for the recapture component but just sex for the survival component. In both cases, the bootstraps values for c-hat were slightly larger than 1 and deviance and AICc values were adjusted accordingly.

Model name (dataset 1980 to 2015), c-hat = 1.11	Delta AICc	Number of Parameters	Deviance
$\Phi(\text{capture occasion} \times \text{sex} + \text{time}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	0.00	261	1127.24
$\Phi(\text{sex} \times \text{arrival} + \text{capture occasion} \times \text{sex}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	8.71	234	1194.99
$\Phi(\text{capture occasion} \times \text{sex} + \text{capture occasion} \times \text{arrival} + \text{sex} \times \text{arrival}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	10.71	235	1194.81
$\Phi(\text{capture occasion} \times \text{sex} \times \text{arrival}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	14.99	237	1194.75
$\Phi(\text{capture occasion} \times \text{sex} + \text{time} \times \text{sex}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	16.92	287	1086.75
Model name (dataset 2005 to 2015), c-hat = 1.07	Delta AICc	Number of Parameters	Deviance
$\Phi(\text{capture occasion} \times \text{sex}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	0.00	65	305.24
$\Phi(\text{capture occasion} \times \text{sex} \times \text{moult}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	1.64	67	302.56
$\Phi(\text{capture occasion}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	1.84	63	311.40
$\Phi(\text{capture occasion} \times \text{sex} + \text{moult}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	2.16	66	305.24
$\Phi(\text{moult}), P(\text{capture occasion} \times \text{time} \times (\text{sex} \times \text{age first capture}))$	2.50	63	312.06

Figure captions

Figure 1: Long-term trends in annual cycle stages of pied flycatchers. **a)** Schematic illustration of the pied flycatcher annual cycle. **b)** Schematic illustration summarising the observed shifts in timing of annual cycle stages across years. **c)** Year to year variation in the timing of different annual cycle stages of pied flycatchers from 1980 until 2015 (without autumn migration) and **d)** 2005 until 2015 (including departure from the Netherlands and arrival at the African wintering grounds). Each point represents the yearly average date (in April days) of the occurrence of the event and each color represents one stage (gray: observed arrival date of males; yellow: calculated arrival dates of males; green: arrival date of females; blue: laying dates; red: hatching dates; pink: moult onset; purple: departure dates; black: arrival dates at the wintering grounds). **e)** Interval (in days) between arrival and laying dates (grey: observed dataset; yellow: calculated dataset; green: females) or between the calculated chick fledgling date and male moult onset (pink) in relation to year. Solid lines represent significant linear or quadratic trends, the thin yellow line represents the significant regression using a broken stick analysis on the interval between arrival and breeding in the calculated arrival dataset.

Figure 2: Trends across years in fitness components. **a)** Clutch size and **b)** proportion of fledged chicks that recruited in relation to year. Solid lines represent model predictions (logistic in b) and proportion of fledged chicks that recruited in relation to **c)** the interval between calculated male arrival and breeding and **d)** the average laying date. Solid lines represent logistic model predictions.



