



K O N I N K L I J K E N E D E R L A N D S E
A K A D E M I E V A N W E T E N S C H A P P E N

Effects of bio-based residue amendments on greenhouse gas emission from agricultural soil are stronger than effects of soil type with different microbial community composition.

Ho, A.; Ijaz, Umer Zeeshan; Janssens, Thierry; Ruijs, Rienke; Kim, Sang Yoon; De Boer, W.; Termorshuizen, Aad J; van der Putten, W.H.; Bodelier, P.L.E.

published in

Global Change Biology Bioenergy

2017

DOI (link to publisher)

[10.1111/gcbb.12457](https://doi.org/10.1111/gcbb.12457)

document version

Publisher's PDF, also known as Version of record

document license

CC BY

[Link to publication in KNAW Research Portal](#)

citation for published version (APA)

Ho, A., Ijaz, U. Z., Janssens, T., Ruijs, R., Kim, S. Y., De Boer, W., Termorshuizen, A. J., van der Putten, W. H., & Bodelier, P. L. E. (2017). Effects of bio-based residue amendments on greenhouse gas emission from agricultural soil are stronger than effects of soil type with different microbial community composition. *Global Change Biology Bioenergy*, 9(12), 1707-1720. <https://doi.org/10.1111/gcbb.12457>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the KNAW public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the KNAW public portal.


Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

pure@knaw.nl

Effects of bio-based residue amendments on greenhouse gas emission from agricultural soil are stronger than effects of soil type with different microbial community composition

ADRIAN HO^{1,*,#} , UMER Z. IJAZ^{2,#}, THIERRY K. S. JANSSENS^{3,4}, RIENKE RUIJS¹, SANG YOON KIM⁵, WIETSE DE BOER^{1,6}, AAD TERMORSHUIZEN⁷, WIM H. VAN DER PUTTEN^{8,9} and PAUL L. E. BODELIER¹

¹Department of Microbial Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Droevendaalsesteeg 10, Wageningen 6708 PB, The Netherlands, ²School of Engineering, University of Glasgow, Oakfield Avenue, Glasgow G12 8LT, UK, ³Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1085-1087, Amsterdam 1081 HV, The Netherlands, ⁴MicroLife Solutions, Science Park 406, Amsterdam 1098XH, The Netherlands, ⁵Agricultural Microbiology Division, National Institute of Agricultural Sciences (NAS), Rural Development Administration (RDA), Jeollabuk-do 55365, Korea, ⁶Department of Soil Quality, Wageningen University and Research (WUR), P.O.Box 47, Wageningen 6700 AA, The Netherlands, ⁷SoilCares Research, Nieuwe Kanaal 7C, Wageningen 6709 PA, The Netherlands, ⁸Department of Terrestrial Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Droevendaalsesteeg 10, Wageningen 6708 PB, The Netherlands, ⁹Laboratory of Nematology, Wageningen University and Research (WUR), P.O. Box 8123, Wageningen 6700ES, The Netherlands

Abstract

With the projected rise in the global human population, agriculture intensification and land-use conversion to arable fields is anticipated to meet the food and bio-energy demand to sustain a growing population. Moving towards a circular economy, agricultural intensification results in the increased re-investment of bio-based residues in agricultural soils, with consequences for microbially mediated greenhouse gas (GHG) emission, as well as other aspects of soil functioning. To date, systematic studies to address the impact of bio-based residue amendment on the GHG balance, including the soil microorganisms, and nutrient transformation in agricultural soils are scarce. Here, we assess the global warming potential (GWP) of *in situ* GHG (i.e., CO₂, CH₄, and N₂O) fluxes after application of six bio-based residues with broad C : N ratios (5–521) in two agricultural soils (sandy loam and clay; representative of vast production areas in north-western Europe). We relate the GHG emission to the decomposability of the residues in a litter bag assay and determined the effects of residue input on crop (common wheat) growth after incubation. The shift in the bacterial community composition and abundance was monitored using IonTorrent™ sequencing and qPCR, respectively, by targeting the 16S rRNA gene. The decomposability of the residues, independent of C : N ratio, was proportional to the GWP derived from the GHG emitted. The soils harbored distinct bacterial communities, but responded similarly to the residue amendments, because both soils exhibited the highest mean GWP after addition of the same residues (sewage sludge, aquatic plant material, and compressed beet leaves). Our results question the extent of using the C : N ratio alone to predict residue-induced response in GHG emission. Taken together, we show that although soil properties strongly affect the bacterial community composition, microbially mediated GHG emission is residue dependent.

Keywords: 16S rRNA gene diversity, C : N ratio, compost, global warming potential, litter bag, nitrous oxide, soil respiration

Received 13 March 2017; accepted 11 May 2017

Introduction

The ongoing growth of the world population, its increasing demand for feed, food, and fibers, in

combination with climate change necessitates a more sustainable circular economy, imposing a need to recycle residues from bio-based production chains into agricultural land. However, closing the loop in a circular economy may have unwanted side effects, such as increased greenhouse gas (GHG) production. Therefore, the consequences of bio-based residue input in agricultural lands have thus led to the need for a comprehensive organic amendment strategy to attenuate GHG emission, while maintaining the soil carbon storage

*Present address: Institute for Microbiology, Leibniz University Hannover, Herrenhäuserstr. 2, Hannover 30140, Germany

#First co-authorships.

Correspondence: Adrian Ho, tel. + 49 511 762 4385, fax + 49 511 762 5287, e-mail: adrian.ho@ifmb.uni-hannover.de

capacity, quality, and fertility (Paustian *et al.*, 2016). Previous work investigated the response of soil microorganisms to nutrient amendments through addition of specific residues in a particular soil type, or in relation to specific microbial guilds mediating greenhouse gas turnover in agricultural soils as case studies (e.g., Jia & Conrad, 2009; Bannert *et al.*, 2011; Ho *et al.*, 2011, 2013; Harter *et al.*, 2014; Bastida *et al.*, 2015; Pereire *et al.*, 2015; Stempfhuber *et al.*, 2016). Case studies are informative, but the nonuniformity of the methodologies used and variable experimental duration complicate cross-study comparisons (Pan *et al.*, 2010; Shade *et al.*, 2016). Rarely is the response of the soil microorganisms, as well as other aspects of soil functioning defined by a suite of parameters tested simultaneously and systematically.

Addition of organic matter in agricultural soils may alter the abundance, composition (community richness and evenness), and biotic interaction among soil microorganisms (Bannert *et al.*, 2011; Ho *et al.*, 2011; Wagg *et al.*, 2014; Hartmann *et al.*, 2015; Sengupta & Dick, 2015; Chen *et al.*, 2016). For instance, residue input seemingly selects for specific members of the soil microbial community (Kuramae *et al.*, 2011; Kuramae *et al.*, 2012). Moreover, soil carbon turnover and its storage capacity are in part mediated by microbial activity where gaseous carbon dioxide and methane can be released into the atmosphere *via* mineralization of organic matter and associated microbial respiration. Gaseous carbon dioxide and methane release can be further stimulated by amendment of organic matter, resulting from the mineralization of newly added carbon and/or the existing carbon fraction, that is, priming effect (Fontaine *et al.*, 2003; Li *et al.*, 2013; Kim *et al.*, 2014; Bastida *et al.*, 2015; Ho *et al.*, 2015a). Apart from organic matter decomposition, soil-inhabiting microorganisms catalyze a multitude of pivotal ecosystem processes in agricultural soils (e.g., methane oxidation: Ho *et al.*, 2015b; biological nitrogen fixation: Herridge *et al.*, 2008; crop disease suppression: Mendes *et al.*, 2011). Considering that microorganisms are key to ecosystem processes (Wagg *et al.*, 2014; Werling *et al.*, 2014), changes in the microbial community and (a)biotic interactions may thus affect the soil microbial mediated processes, with consequences for GHG emission.

Recognizing the limitations of cross-study comparisons and that the effects of organic matter amendment is likely residue and soil type dependent (Cayuela *et al.*, 2010; Paustian *et al.*, 2016), we aim to determine the response of *in situ* GHG (i.e., CO₂, N₂O, CH₄) emission to derive the global warming potential (GWP) of two representative agricultural soil types (sandy loam and clays soils) in north-western Europe

after amendments with different bio-based residues. In general, it is assumed that the decomposition of a bio-based residue is dependent on its C : N ratio (Cayuela *et al.*, 2010). However, most work on decomposition of residue materials varying in C : N ratios have not considered other effects, such as that of GHG emissions (apart from CO₂) and the effects of the residues may have on crop growth. In order to determine whether GHG emission is dependent on the C : N ratio of bio-based residues, we applied six bio-based residues with a broad C : N coverage (5–521) to the soils. In parallel, a litter bag assay was performed to assess the decomposability of these residues and relate it to the CO₂ flux (microbial respiration) and C : N ratio. Likewise, we anticipate higher N₂O emission with the addition of residues with lower C : N ratios, indicative of less recalcitrant materials with high N content. Targeting the 16s rRNA gene, we determined the response of the soil bacterial community composition and abundance, respectively, *via* IonTorrent™ sequencing and quantitative PCR (qPCR) to the bio-based residue amendments. In addition, we account for crop growth (common wheat; *Triticum aestivum*) performance after incubation of the residue-amended soils over three months under controlled conditions. These efforts were intended to elaborate and assess the impact of bio-based residue amendments on soil functioning notably on the microbially mediated GHG emission and to recommend the optimal use of bio-based residues in agriculture.

Materials and methods

Site description, soil sampling, and residues

Soils were collected from agricultural fields from two research stations of Wageningen University and Research, the Netherlands. Vredepeel (51°32'32"N, 05°50'54"E) and Lelystad (52°31'20"N, 05°34'57"E) represent sandy loam and clay soils, respectively. These are typical agricultural soils for the Netherlands, as well as other temperate regions. These fields were planted with potatoes (*Solanum tuberosum*) and left fallow after harvest before sampling. Soil physicochemical properties have been determined before (Ho *et al.*, 2015b). The upper 10 cm of the soils were collected in October 2013 from 1 m × 1 m plots at random and composited as a single bulk soil. The soil was air-dried at room temperature before being sieved to 2 mm. The residues included in this study were comprised of materials with a broad C : N ratio ranging from sewage sludge (5.5) > aquatic plant material (14.0) > commercial compost (15.3) > lignin-rich organic waste stream (17.1; designated 'wood material') > compressed sugar beet leaves (28.0) > paper pulp (518.3). With the exception of the paper pulp, these residues have been described before (Ho *et al.*, 2015b). The paper pulp was sourced from SCA Hygiene Products SUAMEER BV (the Netherlands) to represent a residue with a high C : N ratio and

is characterized by a total C and N of $399.12 \pm 12.26 \mu\text{g C mg dw sample}^{-1}$ and $0.77 \pm 0.09 \mu\text{g N mg dw sample}^{-1}$, respectively. The residues were air-dried at 30°C , crushed, and sieved to 2 mm prior to use.

Experimental setup, gas flux, and nutrient analyses

Mesocosms comprising of 2.5 kg soil were set up in six replicate per soil and residue amendment as described in detail before (Ho *et al.*, 2015b). Briefly, the residues (dry weight) were added into the soils at a rate of 20 ton ha^{-1} as is the typical range in agricultural practice (Diacono & Montemurro, 2010), and deionized water was added and maintained at 65% soil water retention capacity during the incubation. The mesocosms were incubated at 15°C in the dark for approximately two months (56 days), during which, carbon dioxide, methane, and nitrous oxide flux was measured using an Innova 1412-5i Photoacoustic Infrared gas analyzer (lumaSense Technologies, Ballerup, Denmark) connected to an automated sampler equipped with a moisture trap (Innova 1309 multiplexer, lumaSense Technologies). Prior to the gas flux measurement, the mesocosm was placed in a gas tight chamber (diameter \times height: 24 cm \times 40 cm) for 30 min to equilibrate soil-atmosphere gas exchange. Gas flux rate was determined over an hour from at least four time intervals by linear regression ($R^2 > 0.98$). Following the gas flux measurement, soil was sampled using an auger (diameter \times height: 3 cm \times 10–12 cm), homogenized, and an aliquot of the soil was stored in the -20°C freezer till DNA extraction. The remaining soil was stored in the 4°C refrigerator for not more than 14 days to determine the soil physicochemical parameters. To determine biomass loss from the residues, a litter bag assay was performed in parallel using the same setup and incubation conditions with the exception that the residues were contained in a litter bag (0.4 mm mesh size) buried with soil. After incubation, the soils (control and residue-amended) were homogenized by hand in the same pot and planted with two seedlings of common wheat (*Triticum aestivum*), of which the seeds had been pregerminated in the greenhouse at $\sim 22^\circ\text{C}$. The crop was watered thrice weekly until harvest (100 days). After harvest, the aboveground crop biomass was determined after drying in the oven at 60°C till constant weight (approximately three weeks).

Nutrient concentrations in the soil (NO_x , NH_4^+ , and PO_4^{3-}) were determined in 1 M KCl (1 : 5 dilution) extract using a SEAL QuAAtro SFA autoanalyzer (Beun-de Ronde B.V. Abcoude, the Netherlands). The total carbon and nitrogen contents were determined from oven-dried (40°C for 5 days) and sieved (0.4 mm) samples using the Flash EA1112 CN analyzer (Thermo Fisher Scientific, Bleiswijk, the Netherlands). Ergosterol content, used as a proxy for fungal abundance, was determined as described before using the alkaline extraction method (de Ridder-Duine *et al.*, 2006). As reported before (Ho *et al.*, 2015b), the mean soil pH (1 M KCl), organic matter content (% loss on ignition), and C/N ratio for the sandy loam soil were 5.38, 4.74, and 17.30, respectively, and for the clay soil were 7.64, 4.79, and 15.27, respectively.

DNA extraction and quantitative PCR (qPCR) assay

DNA was extracted from three randomly selected mesocosms out of the six replicate per treatment, soil type, and time using the PowerSoil[®] DNA Isolation Kit (MOBIO, Uden, the Netherlands) according to the manufacturer's instruction. The concentration and quality of the DNA extract was determined using a NanoDrop 2000C Spectrophotometer (ThermoScientific, the Netherlands). The DNA extract was stored in the -20°C freezer till further molecular analyses. A qPCR assay targeting the total 16S rRNA gene copies (EUBAC assay) was performed with the Eub338f/Eub518r primer pair (Fierer *et al.*, 2007) with primer concentrations, PCR thermal profile, and subsequent quality check of the amplicons as described in detail before (Ho *et al.*, 2015b). Plasmid DNA extracted from a pure culture (*Collimonas fungivorans* strain TER331) was used to prepare the calibration curve. The qPCR was performed in duplicate for each DNA extract using a Rotor-Gene Q real-time PCR cyclor (Qiagen, Venlo, the Netherlands).

16S rRNA gene amplicon sequencing and data processing

The V4 region of the 16S rRNA gene was amplified using the bacterial primer pair 515F/806R (Caporaso *et al.*, 2011). The 5'-end of the forward primer consisted of the P1 key (underlined) and a two-bp linker GT (bold) (515-F: CCTCTCTATGGGC **AGT**CGGTGATGTGTGCCAGCMGCCGCGGTAA). The 5'-end of the reverse primer consisted of the A key (underlined), a 12-bp Golay barcode (Caporaso *et al.*, 2012), and the CC linker (bold) (806-R: CCATCTCATCCCTGCGTGTCTCCGACT **AG**-Golay-barcode-CCGGACTACHVGGGTWTCTAAT). The PCR reactions were performed with NEB Phusion High-Fidelity Polymerase (NEB Cat# M0530L). The PCR reactions were carried out on a Biorad iCycler (Bio-Rad Laboratories B.V., Veenendaal, the Netherlands) using the following program: initial cycle 98°C for 30 s, followed by 25 cycles of 98°C for 10 s, 50°C for 30 s, and 72°C for 15 s, followed by a final extension step at 72°C for 7 min. Each PCR reaction (Total volume 50 μL) consisted of 0.5 μL of forward and reverse primers (5 μM of each primer), 10 μL of 5 \times New England BioLabs Phusion High-Fidelity Polymerase reaction buffer, 1 μL of 10 mM DNTPs, 1.5 μL DMSO, 3 μL of template DNA, 0.5 μL Phusion High-Fidelity polymerase, and 33 μL nuclease free. The PCR products were purified with the AMPure XP PCR Purification Kit (Agencourt Cat# A638881) and subsequently quantified with Qubit Fluorometer (Thermo Fisher, Invitrogen, Bleiswijk, Netherlands). The samples were then pooled in equimolar ratios and subjected to IonTorrent[™] sequencing.

The DNA fragments were sequenced unidirectionally using the IonTorrent[™] semiconductor technology (Thermo Fisher Scientific) upon two batches of library prepared from 26 pM of equimolarly pooled samples for the sand and clay samples, respectively. The template was made by emulsion PCR using the Ion OneTouch[™] 2 System with the Ion PGM[™] Template OT2 400 Kit and subsequent enrichment for library containing ion sphere particles. The sequencing was performed using HiQ Ion PGM[™] Sequencing 400 kit on Ion PGM[™] System using two

Ion 318™ Chips v2. For data processing, the forward and reverse primers were removed for each sequence from the sample FASTQ files using FLEXBAR version 2.5 (Dodt *et al.*, 2012). Sequences were trimmed by running in the Sickle tool (Joshi & Fass, 2011) to give a minimum and maximum read length of 25 and 150, respectively, before being converted into FASTA format and concatenated into a single file. Subsequently, all reads were clustered into OTUs employing the UPARSE strategy by dereplication and were sorted by abundance (with at least two sequences) and clustered using the UCLUST algorithm (Edgar, 2010). Next, chimeric sequences were detected using the UCHIME algorithm (Edgar *et al.*, 2011). These steps were implemented in VSEARCH version 1.0.10 (Rognes *et al.*, 2016), an open-source, 64-bit multithreaded compatible alternative to USEARCH. Prior to the dereplication step, all reads were mapped to OTUs using USEARCH global method to create an OTU table and converted to BIOM-Format 1.3.1 (McDonald *et al.*, 2012). The sequences have been processed into the BIOM file using Hydra (de Hollander, 2017). Taxonomic information for each OTU was added to the BIOM file by aligning the sequences to the RDP database (release 11) (Cole *et al.*, 2014) using SINA (Pruesse *et al.*, 2007). These steps were implemented using SNAKEMAKE version 3.1 (Köster & Rahmann, 2012). To determine the phylogenetic relationship between OTUs, the OTUs were first multisequence aligned against each other using MAFFT v7.040 (Katoh & Standley, 2013), and an approximately maximum-likelihood phylogenetic tree was generated using FASTTREE v2.1.7 (Price *et al.*, 2010). The OTU table, phylogenetic tree, taxonomic information, and meta data were used in the multivariate statistical analyses in the context of environmental parameters. Nucleotide sequences found in this study were deposited at the European Molecular Biology Laboratory Sequence Read Archive under the study accession number PRJEB20909.

Statistical analysis

The principal coordinate analysis (PCoA), derived from the genus level assignment of OTUs, was performed in R version 2.15.1. (R Development Core Team, 2012) using the capscale function in the VEGAN package (Oksanen *et al.*, 2015). The ordination was visualized considering the community data (OTUs at 3% divergence) using different distance measures: Bray–Curtis considers the species abundance count and Unweighted Unifrac considers the phylogenetic distance between the branch lengths of OTUs observed in the different samples without accounting for the abundances. The samples were grouped for different conditions, as well as the mean ordination value and spread of points (standard deviations of the weighted averages as ellipse using the Ordiellipse function in Vegan). Noneuclidean distances between objects and group centroids were handled by reducing the original distances (Bray–Curtis or Unweighted Unifrac) to principal coordinates and then, performing ANOVA on these data. Analysis of variance using the distance matrices (Bray–Curtis/Unweighted Unifrac), that is, partitioning distance matrices among sources of variation (both quantitative and qualitative information) was performed using Adonis in vegan. This function, hitherto referred to as

PERMANOVA, fits linear models to distance matrices and uses a permutation test with pseudo-*F* ratios.

The DESeqDataSetFromMatrix function in the DESEQ2 package (Love *et al.*, 2014) was used to find OTUs/genera that were significantly distinct between different conditions with a significant cut-off value of 0.001. The DESeqDataSetFromMatrix function allows negative binomial GLM fitting (as abundance data from metagenomics sequencing is over-dispersed) and Wald statistics for abundance data. After performing these multiple testing corrections, OTUs/genera with log-fold changes between multiple conditions are reported. The same test was applied on the genera table which was obtained by binning the OTUs at genus level, with unresolved OTUs grouped as a single ‘unknown’ category. Next, the OTUs were visualized as boxplots after applying a log-relative transformation on the abundance table and given as Venn diagrams. The scripts and workflows for all the analysis above can be downloaded at: <http://userweb.eng.gla.ac.uk/umer.ijaz#bioinformatics>.

Level of significance between amendments and soils was determined in SIGMAPLOT version 12.5 (Systat Software Inc., USA) using ANOVA or *t*-test.

Results

Greenhouse gas fluxes

Residue amendments increased CO₂ flux in both soils, with the exception of the compost- and wood material-amended soils where CO₂ emission was comparable to the un-amended soils over the 56-day incubation (Fig. 1). In particular, cumulative CO₂ emission appreciably increased after amendment with the aquatic plant material and compressed beet leaves. The spike in CO₂ emission occurred at < 21 days; after 21 days, CO₂ flux remained relatively constant till day 56 (Fig. 1).

N₂O emission appreciably increased after sewage sludge amendment in both soils. N₂O emission also increased after the addition of aquatic plant material, but little or no N₂O could be detected in the un-amended and other residue-amended soils (Fig. 2). The paper pulp had minimal or no effect on all the gas fluxes in both soils (Figs 1 and 2). It should be noted that the gas fluxes were determined in the absence of wheat and may exhibit different emission trends in the presence of the crop.

Litterbag assay, crop biomass, and soil nutrient status

The litterbag assay was performed to determine and relate residue decomposability through weight loss to CO₂ flux; the CO₂ flux was used as proxy for soil respiration. The residues with a relatively high weight loss in the litter bag were the sewage sludge (12–15%), aquatic plant material (40–45%), and compressed beet leaves




Fig. 1 CO₂ flux in the un-amended and residue-amended sandy loam (a) and clay (b) soils (mean ± SD; *n* = 6). Mean cumulative CO₂ emitted over incubation period (56 days) in the un-amended and residue-amended soils (c).

(55–60%) (Fig. 3). This trend was consistent, showing comparable residue weight loss in both soils, except paper pulp, of which the loss was significantly higher ($P < 0.05$) in the clay soil (~16%) than in the sandy loam




Fig. 2 N₂O flux in the un-amended and residue-amended sandy loam (a) and clay (b) soils (mean ± SD; *n* = 6). Mean cumulative N₂O emitted over incubation period (56 days) in the un-amended and residue-amended soils (c). Only soils showing N₂O flux are given; residue-amended soils not shown indicate no detectable N₂O flux during incubation.




Fig. 3 Percentage weight loss of residues (mean \pm SD; $n = 6$) as determined from the litter bag assay (a), and crop (wheat; *Triticum aestivum*) yield (mean \pm SD; $n = 5$) after residue amendments (b). Inset figure in (b) shows the crop yield in a repeated experiment in 2015 after sewage sludge and paper pulp amendments. Soils and residues were collected from the same source, albeit different times, in the repeated mesocosm incubation. Upper and lower case letters indicate the level of significance at $P < 0.05$ (ANOVA) between treatments in the sandy loam and clay soils, respectively. The residues are given in order of increasing C : N ratios on the x -axis.

soil (~4%) (Fig. 3). Compost and wood material showed no or marginal weight loss at 0.5–1.3%, respectively.

After incubation, wheat was planted in the pots to assess the effects of residue amendment on crop performance. Biomass yield, as determined by shoot biomass dry weight, was approximately 50% higher in the clay soil compared to the sandy loam soil (Fig. 3). Significantly higher crop biomass was recorded in the sewage sludge- and aquatic plant material-amended soils,

whereas compressed beet leaves stimulated crop growth only in the sandy loam soil (Fig. 3). While crop yield in the compost- and wood material-amended soils was comparable to the un-amended soils, showing no net effect of these residues on the crop performance, paper pulp amendments significantly suppressed crop growth in both soils. Additionally, the mesocosm study was repeated using the same soils (sampled in 2015) and residues giving contrasting crop yields (i.e., sewage sludge and paper pulp); results confirmed the stimulatory and suppression effects of sewage sludge and paper pulp, respectively, on crop growth, although overall crop yield was lower (Fig. 3; inset figure).

As reported before (Ho *et al.*, 2015b), NH_4^+ concentration was highest after amendment with sewage sludge, but levels gradually decreased, while NO_x concentration increased over time in both soils. This indicated nitrification and is in agreement with the observed appreciable higher N_2O emission after sewage sludge amendment in this study (Fig. 2). This trend was also detected in the aquatic plant material-amended soils, albeit to a lesser extent (Ho *et al.*, 2015b). In other residue-amended soils including the paper pulp, the novel residue in this study, nutrient levels remained relatively constant during incubation.

Ergosterol concentration was determined after 21 and 56 days of incubation and was used as a proxy for fungal abundance. Initial ergosterol concentration in both un-amended soils was comparable (Table 1). Consistent in both soils, ergosterol concentration significantly increased after amendments with the residues, reaching appreciably higher concentrations in the aquatic plant material-, compressed beet leaves-, and paper pulp-amended soils (Table 1). In some of these soils (e.g., both aquatic plant material-amended soils), ergosterol concentration decreased from 21 to 56 days after incubation, while in others, ergosterol concentration increased (e.g., compressed beet leaves-amended sandy loam soil) or remained relatively constant (e.g., compressed beet leaves-, and paper pulp-amended clay soil). Although significantly higher, ergosterol concentration in the sewage sludge-, compost-, and wood oxidation material-amended soils was relatively low (Table 1).

Soil bacterial 16S rRNA gene abundance and composition

Previously, we characterized the methanotrophs quantitatively and qualitatively by targeting the *pmoA* gene (Ho *et al.*, 2015b). Now, we focused on the total bacterial abundance and community composition by targeting the 16S rRNA gene. In the un-amended soils, 16S rRNA gene copies gradually decreased during the incubation, while in the sewage sludge-, aquatic plant material-,

Table 1 Ergosterol concentration (mg kg soil⁻¹) at 21 and 56 days after incubation in the un-amended and residue-amended soils (mean ± SD; *n* = 6). The residues are given in order of increasing C : N ratios. Letters indicate level of significance at *P* < 0.01 between treatments per time and soil

Treatment	Sandy loam soil		Clay soil	
	21 days	56 days	21 days	56 days
Un-amended	0.40 ± 0.04 ^a	0.45 ± 0.07 ^a	0.38 ± 0.09 ^a	0.44 ± 0.06 ^a
Sewage sludge amended	4.33 ± 1.35 ^d	3.48 ± 1.13 ^d	1.55 ± 0.59 ^b	1.10 ± 0.26 ^c
Aquatic plant material amended	10.26 ± 0.97 ^e	3.55 ± 1.01 ^d	7.01 ± 0.66 ^d	3.85 ± 0.95 ^e
Compost amended	0.64 ± 0.17 ^b	1.03 ± 0.19 ^b	0.64 ± 0.28 ^a	0.63 ± 0.03 ^b
Wood oxidation material amended	1.49 ± 0.47 ^c	1.92 ± 0.34 ^c	1.60 ± 0.65 ^b	1.84 ± 0.26 ^d
Compressed beet leaves amended	0.94 ± 0.13 ^c	5.56 ± 1.32 ^e	9.86 ± 2.78 ^d	9.07 ± 2.23 ^g
Paper pulp amended	10.06 ± 1.91 ^e	2.53 ± 0.20 ^d	4.90 ± 0.88 ^c	5.57 ± 0.65 ^f

and compressed beet leaves-amended soils, the 16S rRNA gene copies increased, reaching a peak before values remained relatively constant or decreased (Fig. 4). This trend coincided with the spike in CO₂ flux <21 days (Fig. 1). Although the 16S rRNA gene copies in the sewage sludge-, wood material-, and compressed beet leaves-amended soils were significantly altered after 56 days (pairwise comparison *t*-test; *P* < 0.05), values were not appreciably higher than in the un-amended soils. In the paper pulp-amended soils, mean 16S rRNA gene copies were lower than in the un-amended soil throughout the incubation.

The bacterial community composition was determined after 7 days (for both soils), and 13 days (for the clay soil), or 15 days (for the sandy loam soil), coinciding with the period with the highest CO₂ emission assuming that the strong CO₂ flux beyond background levels (un-amended soils) is caused by residue-induced microbial activity. The principal coordinate analysis (PCoA) revealed that the bacterial community composition was distinct in both soils, even after amendments with the same residues (Fig. 5). More pronounced in the sandy loam soil, the bacterial community composition was more similar following aquatic plant material and compressed beet leaves amendments than in the un-amended soils (Fig. 5). Indeed, we identified treatments (residue amendments), soil types, time (days), and pH to be highly significant (*P* = 0.001), respectively, accounting for 29.9%, 24.5%, 5.7%, and 3% of the variability influencing the community composition when we performed PERMANOVA of the community data against the different environmental parameters. Among the soil bacteria, identifiable predominant members fall into the Proteobacteria, Bacteroidetes, Actinobacteria, Acidobacteria, and Firmicutes at the phylum level, together forming the vast majority of the bacterial community composition (>70% relative abundance) in the sandy loam and clay soils (Fig. S1); differences in the

community composition in these soils became apparent at the genus level (Figs S1–S3).

To detect discrepancies in the bacterial community composition between and within soils and treatments, we considered only OTUs showing strong changes (i.e., log-fold), termed as ‘significant’ OTUs/genera. Although the sequence length allows the identification of OTUs to the genus level, some OTUs could only be assigned to the family level (Fig. S1). Hence, the significant members of the community were represented here at the family level (Figs S2 and S3). After genus level assignment of the OTUs (boxplot; Figs S4 and S5), the significant genera between treatments were composed of community members belonging to the phylum Actinobacteria, Bacteroidetes, Firmicutes, Proteobacteria, and Plantomycetes in both soils; undefined genera identified as *incertae sedis* in the Ribosomal Database Project (RDP) database were not considered (Figs S2 and S3). Besides, significant genera belonging to the Verrucomicrobia and Nitrospirae were only detected in the clay soil after sewage sludge and compost amendment, respectively. Unique to the sandy loam soil are members of Armatimonadates and Thaumarchaeota in the compost-amended soil and Acidobacteria in the beet leaves-amended soil.

Both soils harbored similar significant amounts of the members of the Bacteroidetes (i.e., families Flexibacteraceae, Sphingobacteriaceae, Chitinophagaceae, Flavobacteriaceae, and Cytophagaceae) although they do not necessarily occur in the same residue amendments in both soils (Figs S2 and S3). Actinobacteria was represented by some families (e.g., Conexibacteraceae, Glycomycetaceae, Promicromonosporaceae, Streptomyetaceae, and Acidimicrobiaceae) which were not common to both soils (Figs S2 and S3). Some Actinobacterial families only emerged after specific residue amendments (e.g., Acidimicrobiaceae in compost-amended clay soil, and all families in the residue-amended sandy loam soil,




Fig. 4 Change in the total bacterial 16S rRNA gene abundance using qPCR in the un-amended and residue-amended sandy loam (a) and clay (b) soils (mean \pm SD; $n = 6$). The 16S rRNA gene copy numbers were determined in duplicate for each DNA extract ($n = 3$) per treatment and time, giving a total of six replicate 16S rRNA gene quantification. Shifts in the 16S rRNA gene copy in the un-amended, and sewage sludge-, aquatic plant material-, and compost-amended incubations are as given in Ho *et al.* (2015b). Results were reanalyzed to include shifts in the 16S rRNA gene copy in the wood material-, compressed beet leaves-, and paper pulp-amended soils in this study. The inset figures show linear changes in the 16S rRNA gene copy numbers.

as no Actinobacterial genus was considered significant in the un-amended soil; Figs S2 and S3). Although Planococcaceae, Paenibacillaceae, and Bacillaceae belonging to Firmicutes are common in both soils, members of the family Turicibacteraceae, Leuconostocaceae, and Clostridiaceae are unique significant genera detected in the wood oxidation material-, beet leaves-, and compost-/paper pulp-amended sandy loam soil, respectively (Fig. S2). Within Proteobacteria, significant genera fell into the subphylum Alpha-, Beta-, Delta-, and

Gammaproteobacteria. Members of Gamma- and Deltaproteobacteria showed the highest variability without any shared significant genera between the soils and amendments (Figs S2 and S3). In contrast, with the exception of the wood oxidation material-amended clay soil, Oxalobacteraceae (subphylum Betaproteobacteria) was found to be significant in all residue-amended sandy loam and clay soils, including the un-amended clay soil (Figs S2 and S3). Within Alphaproteobacteria, Caulobacteraceae was significant in all residue-amended sandy loam soil, with the exception of the wood oxidation material-amended soil, but was not found to be significantly abundant in the clay soil (Figs S2 and S3).

Discussion

Response of microbial activity to bio-based residue amendment

The total CO₂ emission trend was in agreement with the percentage residue loss as determined in the litter bag assay, in that, amendments with aquatic plant material and compressed beet leaves emitted the highest cumulative CO₂ levels. This is in line with the documented highest residue weight loss in both soils (up to 60% during the 56-day incubation; Figs 1 and 3). In contrast, compost and wood oxidation material exhibited marginal weight loss in the litter bag assay (<2%) and also showed the lowest total CO₂ emissions in both soils. Therefore, the aquatic plant material, compressed beet leaves, and sewage sludge can be regarded as readily degradable residues despite of their rather broad C : N ratios (5–28). The C : N ratio is typically used to assess the recalcitrance of organic matter, with a high C : N ratio (>18) indicating the presence of more complex organic material (Huang *et al.*, 2004; Cayuela *et al.*, 2010). Hence, our results question the extent of using C : N ratios alone to predict the soil response to residue input and consequences for greenhouse gas flux. While we cannot completely exclude a priming effect causing increased soil respiration following residue amendments (Fontaine *et al.*, 2003; Cong *et al.*, 2015), significant CO₂ release coupled to a relatively higher biomass loss of the same residues suggest that the newly added carbon source, rather than the existing carbon fraction in the soil was mineralized, in which case, priming effect would be minimal.

As anticipated based on the relatively low C : N ratio (5.5), N₂O emission was highest after sewage sludge addition. On the other hand, CH₄ flux showed a more variable trend over time, depending on the residue, but independent of the residue C : N ratio (Ho *et al.*, 2015b). We derived the global warming potential (GWP) in mg CO₂ equivalent per kg soil (IPCC, 2007) by




Fig. 5 Principal coordinate analysis (PCoA) colored by treatments (i.e., un-amended and residue-amended sandy loam and clay soils) using Bray–Curtis distance showing the first two principal axes along with the percentage variability explained. The ellipse shows 95% confidence interval of the standard errors. The symbols depict the time (d; days) of incubation.

re-analyzing the cumulative CH_4 flux (Ho *et al.*, 2015b) in combination with the cumulative CO_2 and N_2O fluxes (Figs 1 and 2). The GWP value for CH_4 and N_2O is, respectively, considered to be 25 and 298 over a hundred-year time frame, while the GWP value for CO_2 is regarded as 1 (IPCC, 2007). With the exception of the sewage sludge-amended sandy loam soil where N_2O was the main contributor to the GWP, CO_2 is the main contributor to the GWP for the other amendments. All residue amendments increased the mean GWP. Amendments with sewage sludge and compressed beet leaves, respectively, recorded a four- to ninefold and four- to fivefold GWP increase, whereas amendments with compost and wood oxidation material showed marginal GWP change (0.4- to 0.6-fold higher) when compared to the un-amended soil (Fig. 6). As the residues are used

to replace or in conjunction with mineral fertilizers, future work could consider the GWP caused by amendments of a combination of residue and mineral fertilizers under field conditions.

The CO_2 , CH_4 , and N_2O emission patterns and rates responded similarly to the different residue additions despite the soils (sandy loam and clay soils) possessing different physicochemical parameters and soil texture (e.g., pH, C : N ratio; Ho *et al.*, 2015b) and harbored distinct bacterial community composition (Fig. 5), indicating that residue input exerts a stronger impact on microbially mediated GHG emission than soil properties during the controlled incubation. Alternatively, although the bacterial community composition in both soils was distinct, they apparently shared similar traits, presumably as a result of a shared history (i.e.,




Fig. 6 Mean global warming potential (GWP) in the un-amended and residue-amended soils derived from the cumulative CO₂ (Fig. 1), N₂O (Fig. 2), and CH₄ (Ho *et al.*, 2015b) fluxes over 56 days. The residues are given in order of increasing C : N ratios on the x-axis.

comparable agricultural regime), in mediating C- and N-cycling.

Trade-off between crop production and greenhouse gas emissions

In contrast to previous work solely determining greenhouse gas emissions and/or the response of the soil microbial community composition to (in)organic nutrient amendments (e.g., Huang *et al.*, 2004; Ceja-Navarro *et al.*, 2010; Serrano-Silva *et al.*, 2011; Galvez *et al.*, 2012; Odlare *et al.*, 2012; Li *et al.*, 2013; Sengupta & Dick, 2015), common wheat, a cereal grain crop, was planted in the mesocosms to assess crop growth and productivity (aboveground dry weight biomass) after residue amendments. Amendments with sewage sludge and aquatic plant material yielded higher crop biomass in both soils, in line with the higher N (NO_x and NH₄⁺) and P concentrations in the residue-amended soils prior to wheat addition (Ho *et al.*, 2015b). This indicates nutrient limitation for crop, as well as for soil microorganisms. Using measured environmental parameters as constraints and qPCR data for a correspondence analysis revealed a positive correlation of NH₄⁺ concentration and CO₂ flux (Ho *et al.*, 2015b), indicating a relief of N limitation for organic matter mineralization by soil microorganisms after residue input. Interestingly, paper pulp (in)directly suppressed crop growth. Repeating the experiment with the same soils (collected in 2015) and residues further confirmed our results (Fig. 3). The

bacterial community composition in the paper pulp- and un-amended incubations was relatively similar as revealed by the principal coordinate analysis (Fig. 5), yet wheat biomass was significantly lower in the paper pulp-amended soils. Hence, the significantly adverse effect on plant growth is unlikely associated to the soil bacterial composition. Rather, the immobilization of available nitrogen caused by the high C:N ratio of paper pulp may lead to acute N-limiting condition for the crop. Therefore, not all bio-based residues are suitable soil additives. Although some residues may contribute to soil carbon (paper pulp C : N ratio: 518.3) and nutrients as in this study and others (e.g., Galvez *et al.*, 2012; Kirkby *et al.*, 2013; Pereire *et al.*, 2015), the positive effects may be offset by detrimental effects on crop growth.

Among the residues tested, compost and wood oxidation material impose the lowest GWP, comparable to the un-amended soils. However, these residues did not yield higher crop biomass, indicating an apparent net zero effect with regard to overall greenhouse gas turnover (CO₂, CH₄, and N₂O) and crop production, while causing minimal change in the bacterial community composition. Although seemingly having no net effect in the short term, the use of these residues is not without merit and may benefit other aspects of soil functioning. In particular, compost mineralization is slower than in fresh organic residues; hence, the extraneous carbon pool is sequestered in the soil for longer periods or is accumulated with recurring compost amendments (Ryals *et al.*, 2015). Previously, we showed a significant stimulation of soil methane uptake with compost addition (Ho *et al.*, 2015b). Although transient, compost-induced methane uptake offsets up to 16% of total carbon dioxide emitted during the incubation (Ho *et al.*, 2015b). Therefore, agricultural soils may benefit from repeated compost amendments as means of a carbon sequestration strategy or by stimulating soil methane consumption, thereby reducing the GWP in agricultural lands.

Although crop yield was compromised in favor for reduced GWP, crop productivity, and properties may improve considering compost amendment complemented with other (in)organic soil additives (Ye *et al.*, 2016). Application of inorganic fertilizer and organic residue mixtures is indeed common agricultural practice, which has also been shown to increase organic matter and nutrient retention (Kirkby *et al.*, 2013; Chen *et al.*, 2016). Hence, future studies under field conditions determining the optimum proportions of inorganic fertilizer and organic soil additives (i.e., to achieve optimum C : N : P ratio through combinations of these materials), as well as the application rate to minimize GWP while allowing sustainably high crop yields

warrant further attention. More generally, our results reinforce the need for a more comprehensive approach when assessing 'climate-smart' agricultural soil management practices (Paustian *et al.*, 2016; Ye *et al.*, 2016).

Bacterial 16S rRNA gene copies and fungal abundances in residue-amended soils

Given that intensive Dutch agricultural soils are dominated by bacteria (Van der Wal *et al.*, 2006), we followed the shift in the bacterial abundance over the incubation period and considered the community composition during the strong CO₂ efflux. However, as certain fungi can respond rapidly to organic matter additions in agricultural soils (Van der Wal *et al.*, 2013), the fungal biomass was determined after 21 and 56 days.

Despite a general trend showing the gradual decrease in the 16S rRNA gene copy numbers in the un-amended and residue-amended soils, values were higher

(7–21 days) after reaching a peak at day 7 following sewage sludge and aquatic plant material amendments. The concomitant increase in the 16S rRNA gene copy numbers and abrupt CO₂ increase (<21 days) beyond levels exhibited in the un-amended soils suggest residue-induced stimulation of bacterial abundance or growth (Figs 1 and 4). However, this trend was not observed in the compressed beet leaves-amended soils as would be anticipated if decomposition was bacteria mediated. Unexpectedly, the 16S rRNA gene copy numbers in the compressed beet leaves-amended soils remained relatively low and were comparable to values in the un-amended soils, albeit a significantly higher CO₂ emission as well as a higher residue weight loss were detected. Instead, the ergosterol concentration in the compressed beet leaves-amended soils was appreciably higher than in the un-amended soils ≥ 21 days, indicating the relatively important role of fungal degradation in this amendment.

Ergosterol concentration was significantly higher (ANOVA; $P < 0.05$; Table 1) in all residue-amended soils than in the controls, but was appreciably higher after aquatic plant material, compressed beet leaves, and paper pulp amendments in both soils. This suggests the relative dominance and stimulation of the fungal population after amendments with specific residues. Even in bacteria-dominated agricultural soils, fungi can respond quickly to addition of organic materials (Van der Wal *et al.*, 2006). This appears to be the case in the current study. Therefore, fungi have contributed to the production of CO₂. However, their possible involvement in the production or consumption of other GHG gases is not well-documented (Thomson *et al.*, 2012).

Bacterial community response to bio-based residue amendments

The shift in the bacterial community composition following substrate addition had been documented before and could be related to the rapid CO₂ efflux in soils (Fig. 1; Padmanabhan *et al.*, 2003; Cleveland *et al.*, 2007). Both soils harbored distinct bacterial communities, as revealed by the principal coordinate analysis with the plots depicting Bray–Curtis distances between the sandy loam and clay soils (Fig. 5). Phyla-level bacterial community response provides an overview, but may mask differential response among members of the same phylum (Ranjan *et al.*, 2015; Morrissey *et al.*, 2016; Ho *et al.*, 2017). Therefore, we monitored the response of significant genera represented at the family level to bio-based residue amendments. The significant genera identified belonged to the phyla (e.g., Actinobacteria, Bacteroidetes, Firmicutes, and Proteobacteria) common in agricultural soils from diverse geographical origins (Noll *et al.*, 2005; Kuramae *et al.*, 2012; Hartmann *et al.*, 2015).

Interestingly, with the exception of the compressed beet leaves-amended sandy loam soil and sewage sludge-amended clay soil, members of Verrucomicrobia (families Opiritaceae and Verrucomicrobiaceae) and Acidobacteria (family Acidobacteriaceae) were not found to be significant in all other treatments, including the un-amended soils despite being shown to be ubiquitous and abundant across widespread agricultural and pristine environments (Jones *et al.*, 2009; Bergmann *et al.*, 2011; Kuramae *et al.*, 2012; Ranjan *et al.*, 2015; Kielak *et al.*, 2016; Ho *et al.*, 2017). The dominance of Opiritaceae and Verrucomicrobiaceae in the sewage sludge-amended clay soil thus suggests that these bacteria may have been introduced along with the residue and persisted in the soil, but their dominance was not observed in the sandy loam soil after sewage sludge addition. Although members of Verrucomicrobia and Acidobacteria are physiologically and taxonomically heterogeneous, studies consistently indicate that members of these phyla were more competitive under oligotrophic conditions (Fierer *et al.*, 2007; Ho *et al.*, 2017). In agreement with this presumption, bacteria assigned to the Verrucomicrobia and Acidobacteria appear to represent only a relatively minor fraction among the soil bacterial community after amendments, as well as in the un-amended soils. This suggests that the agricultural soils did not remain substrate/nutrient impoverished, as would be anticipated with ongoing agriculture where substrate/nutrient addition is a recurring practice.

Conversely, some bacterial families become important after residue input. These include Oxalobacteraceae of Betaproteobacteria and Chitinophagaceae of

Bacteroidetes in the sandy loam soils after residue amendments (Fig. S2). In the clay soil, these bacteria also became significant, but only after amendments with certain residues (i.e., sewage sludge, aquatic plant material, and compost for Chitinophagaceae), as well as appearing in the un-amended incubations (i.e., Oxalobacteraceae) (Fig. S3). This suggests that members of Chitinophagaceae were more responsive to the residue amendments, and appreciably increased in relative abundance, more evident in the sandy loam soil. Hence, some members of particular phyla may indeed be disproportionately important to labile C-decomposition, but they are by no means definitely copiotrophs (please see review Ho *et al.*, 2017). Tracking their persistence or cessation with labeled substrates and teasing apart other confounding factors (e.g., nutrient co-limitation) in prolonged incubation following the consumption of labile materials may reveal their ecological role.

The dominance of some bacterial families after residue amendments in the different soil types, but not in both soils after the same residue amendments, suggests that the soil properties override the effect of residue addition in altering the bacterial community composition. The PCoA supports this hypothesis, depicting a divergence in the relative abundance of taxons in both soils after residue amendments (Fig. 5) with some members contributing to the variability between the community compositions in these soils. For instance, significant genera constituting Planctomycetes appeared only in the clay soil, and those belonging to Gammaproteobacteria and Deltaproteobacteria are almost exclusive to either soils but not both (Figs S2 and S3). Nevertheless, the community composition after amendments with the more easily degradable residues (i.e., aquatic plant material, compressed beet leaves, and to a lesser extent sewage sludge) assuming higher respiration rate and residue weight loss as an indicator for degradability tended to cluster together (Fig. 5). This trend was more evident in the sandy loam soil where the bacterial community composition after amendments with the aquatic plant material and compressed beet leaves was more similar than in the un-amended soil. Consistent with previous work, soil properties (e.g., C : N ratio, pH) have been documented to be strong determinants and take precedence over land-use type and management in selecting for soil microorganisms (Lauber *et al.*, 2008; Kuramae *et al.*, 2012; Delmont *et al.*, 2014), as well as their activity (Li *et al.*, 2013).

Taken together, we conclude that soil properties are important determinants of the soil microbial community composition which take precedence over the effects of bio-based residue amendments. However, residue type, independent of C : N ratio, elicits a stronger response in the functioning of these communities with respect to GHG emission.

Acknowledgements

We are grateful to Iris Chardon, Marion Meima-Franke, and Tirza Andrea for excellent technical assistance. We extend our gratitude to Mattias de Hollander for assistance during the sequence analysis. UZI is supported by NERC Independent Research Fellowship NE/L011956/1. AH is financially supported by the BE-Basic Grant F03.001 (SURE/SUPPORT). This publication is publication nr. 6296 of the Netherlands Institute of Ecology.

Conflict of interest

The authors have no conflict of interests.

References

- Bannert A, Kleineidam K, Wissing L *et al.* (2011) Changes in diversity and functional gene abundances of microbial communities involved in nitrogen fixation, nitrification, and denitrification in a tidal wetland versus paddy soils cultivated for different time periods. *Applied and Environmental Microbiology*, **77**, 6109–6116.
- Bastida F, Selevsek N, Torres IF, Hernandez T, Garcia C (2015) Soil restoration with organic amendments: linking cellular functionality and ecosystem processes. *Scientific Reports*, **5**, 15550.
- Bergmann GT, Bates ST, Eilers KG *et al.* (2011) The under-recognized dominance of Verrucomicrobia in soil bacterial communities. *Soil Biology and Biochemistry*, **43**, 1450–1455.
- Caporaso JG, Lauber CL, Costello EK *et al.* (2011) Moving pictures of the human microbiome. *Genome Biology*, **12**, R50. <http://genomebiology.com/2011/12/5/R50>
- Caporaso JG, Lauber CL, Walters WA *et al.* (2012) Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME Journal*, **6**, 1621–1624.
- Cayuela ML, Oenema O, Kuikman PJ, Bakker RR, van Groenigen JW (2010) Bioenergy by-products as soil amendments? Implications for carbon sequestration and greenhouse gas emissions. *Global Change Biology Bioenergy*, **2**, 201–213.
- Ceja-Navarro JA, Rivera-Orduña FN, Patiño-Zúñiga L, Vila-Sanjurjo A, Crossa J, Govaerts B, Dendooven L (2010) Phylogenetic and multivariate analyses to determine the effects of different tillage and residue management practices on soil bacterial communities. *Applied and Environmental Microbiology*, **76**, 3685–3691.
- Chen C, Zhang J, Lu M *et al.* (2016) Microbial communities of an arable soil treated for 8 years with organic and inorganic fertilizers. *Biology and Fertility of Soils*, **52**, 445–467.
- Cleveland CC, Nemergut DR, Schmidt SK, Townsend AR (2007) Increases in soil respiration following labile carbon additions linked to rapid shifts in soil microbial community composition. *Biogeochemistry*, **82**, 229–240.
- Cole JR, Wang Q, Fish JA *et al.* (2014) Ribosomal database project: data and tools for high throughput rRNA analysis. *Nucleic Acid Research*, **42**, 633–642.
- Cong WF, Hoffland E, Li L, Janssen BH, van der Werf W (2015) Intercropping affects the rate of decomposition of soil organic matter and root litter. *Plant and Soil*, **391**, 399–411.
- Delmont TO, Francioli D, Jacquesson S *et al.* (2014) Microbial community development and unseen diversity recovery in inoculated sterile soil. *Biology and Fertility of Soils*, **50**, 1069–1076.
- de Ridder-Duine AS, Smant W, van der Wal A, van Veen JA, de Boer, W. (2006) Evaluation of a simple, non-alkaline extraction protocol to quantify soil ergosterol. *Pedobiologia*, **50**, 293–300.
- Diacono M, Montemurro F (2010) Long-term effects of organic amendments on soil fertility: a review. *Agronomy for Sustainable Development*, **30**, 401–422.
- Doti M, Roehr JT, Ahmed R, Dieterich C (2012) FLEXBAR – Flexible barcode and adapter processing for next-generation sequencing platforms. *Biology*, **1**, 895–905.
- Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, **26**, 2460–2461.
- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics*, **27**, 2194–2200.

- Fierer N, Bradford MA, Jackson RB (2007) Toward an ecological classification of soil bacteria. *Ecology*, **88**, 1354–1364.
- Fontaine S, Mariotti A, Abbadie L (2003) The priming effect of organic matter: a question of microbial competition? *Soil Biology and Biochemistry*, **35**, 837–843.
- Galvez A, Sinicco T, Cayuela ML, Mingorance MD, Fornasier F, Mondini C (2012) Short term effects of bioenergy by-products on soil C and N dynamics, nutrient availability and biochemical properties. *Agricultural, Ecosystems and Environment*, **160**, 3–14.
- Harter J, Krause H-M, Schuetzler S *et al.* (2014) Linking N₂O emissions from biochar-amended soil to the structure and function of the N-cycling microbial community. *ISME Journal*, **8**, 660–674.
- Hartmann M, Frey B, Mayer J, Mäder P, Widmer F (2015) Distinct soil microbial diversity under long-term organic and conventional farming. *ISME Journal*, **9**, 1177–1194.
- Herridge DF, Peoples MB, Boddey RM (2008) Global inputs of biological nitrogen fixation in agricultural systems. *Plant and Soil*, **311**, 1–18.
- Ho A, Lüke C, Cao Z, Frenzel P (2011) Ageing well: methane oxidation and methane-oxidizing bacteria along a chronosequence of 2000 years. *Environmental Microbiology Reports*, **3**, 738–743.
- Ho A, Lüke C, Reim A, Frenzel P (2013) Selective stimulation in a natural community of methane oxidizing bacteria: effects of copper on *pmoA* transcription and activity. *Soil Biology and Biochemistry*, **65**, 211–216.
- Ho A, El-Hawwary A, Kim SY, Meima-Franke M, Bodelier PLE (2015a) Manure-associated stimulation of soil-borne methanogenic activity in agricultural soils. *Biology and Fertility of Soils*, **51**, 511–516.
- Ho A, Reim A, Kim SY *et al.* (2015b) Unexpected stimulation of soil methane uptake as emergent property of agricultural soils following bio-based residue application. *Global Change Biology*, **21**, 3864–3879.
- Ho A, Di LP, Bodelier PLE (2017) Revisiting life strategy concepts in environmental microbial ecology. *FEMS Microbiology Ecology*, **93**, 1–14: <https://doi.org/10.1093/femsec/fix006>.
- de Hollander M (2017) nioo-knaw/hydra: 1.1. <https://doi.org/10.5281/zenodo.556539>.
- Huang Y, Zou J, Zheng X, Wang Y, Xu X (2004) Nitrous oxide emissions as influenced by amendment of plant residues with different C: N ratios. *Soil Biology and Biochemistry*, **36**, 973–981.
- IPCC (2007) Summary for policymakers. In: *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Parry ML, Canziani OF, Palutikof JP, Van der Linden PJ, Hanson CE), pp. 81–82. Cambridge University Press, Cambridge.
- Jia Z, Conrad R (2009) Bacteria rather than Archaea dominate microbial ammonia oxidation in an agricultural soil. *Environmental Microbiology*, **11**, 1658–1671.
- Jones RT, Robeson MS, Lauber CL, Hamady M, Knight R, Fierer N (2009) A comprehensive survey of soil acidobacterial diversity using pyrosequencing and clone library analyses. *ISME Journal*, **3**, 442–453.
- Joshi NA, Fass JN (2011) Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files (version 1.33) Software. Available at: https://github.com/na_joshi/sickle. (accessed 15 July 2016).
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, **30**, 772–780.
- Kielak AM, Barreto CC, Kowalchuk GA, van Veen JA, Kuramae EE (2016) The ecology of *Acidobacteria*: moving beyond genes and genome. *Frontiers in Microbiology*, **7**, 744.
- Kim SY, Pramanik P, Gutierrez J, Hwang HY, Kim PJ (2014) Comparison of methane emission characteristics in air-dried and composted cattle manure amended paddy soil during rice cultivation. *Agriculture, Ecosystems and Environment*, **197**, 60–67.
- Kirkby CA, Richardson AE, Wade LJ, Batten GD, Blanchard C, Kirkegaard JA (2013) Carbon-nutrient stoichiometry to increase soil carbon sequestration. *Soil Biology and Biochemistry*, **60**, 77–86.
- Köster J, Rahmann S (2012) Snakemake – a scalable bioinformatics workflow engine. *Bioinformatics*, **28**, 2520–2522.
- Kuramae E, Gamper H, van Veen J, Kowalchuk G (2011) Soil and plant factors driving the community of soil-borne microorganisms across chronosequences of secondary succession of chalk grasslands with a neutral pH. *FEMS Microbiology Ecology*, **77**, 285–294.
- Kuramae EE, Yergeau E, Wong LC, Pijl AS, van Veen JA, Kowalchuk GA (2012) Soil characteristics more strongly influence soil bacterial communities than land-use type. *FEMS Microbiology Ecology*, **79**, 12–24.
- Lauber CL, Strickland MS, Bradford MA, Fierer N (2008) The influence of soil properties on the structure of bacterial and fungal communities across land-use types. *Soil Biology Biochemistry*, **40**, 2407–2415.
- Li F, Cao X, Zhao L *et al.* (2013) Short-term effects of raw rice straw and its derived biochar on greenhouse gas emission in five tropical soils in China. *Soil Science and Plant Nutrition*, **59**, 800–811.
- Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, **15**, 550.
- McDonald D, Clemente JC, Kuczynski J *et al.* (2012) The biological observation matrix (BIOM) format or: how I learned to stop worrying and love the ome-ome. *GigaScience*, **1**, 7.
- Mendes R, Kruijt M, de Bruijn I *et al.* (2011) Deciphering the rhizosphere micro-biome for disease-suppressive bacteria. *Science*, **332**, 1097–1100.
- Morrissey EM, Mau RL, Schwartz E *et al.* (2016) Phylogenetic organization of bacterial activity. *ISME Journal*, **10**, 2336–2340.
- Noll M, Matthies D, Frenzel P, Derakhshani M, Liesack W (2005) Succession of bacterial community structure and diversity in a paddy soil oxygen gradient. *Environmental Microbiology*, **7**, 382–395.
- Odlare M, Abubaker J, Lindmark J, Pell M, Thorin E, Nehrenheim E (2012) Emissions of N₂O and CH₄ from agricultural soils amended with two types of biogas residues. *Biomass and Bioenergy*, **44**, 112–116.
- Oksanen J, Blanchet F, Kindt R *et al.* (2015) Vegan: community ecology package. R package version 2.2-1, Available at: <http://cran.r-project.org/package=vegan> (accessed 1 July 2016).
- Padmanabhan P, Padmanabhan S, De Rito C *et al.* (2003) Respiration of 13C-labeled substrates added to soil in the field and subsequent 16S rRNA gene analysis of 13C-labeled soil DNA. *Applied and Environmental Microbiology*, **69**, 1614–1622.
- Pan Y, Bodrossy L, Frenzel P *et al.* (2010) Impacts of inter- and intralaboratory variations on the reproducibility of microbial community analyses. *Applied and Environmental Microbiology*, **76**, 7451–7458.
- Paustian K, Lehmann J, Ogle S, Reay D, Robertson GP, Smith P (2016) Climate-smart soils. *Nature*, **532**, 49–57.
- Pereire ESP, Suddick EC, Mansour I *et al.* (2015) Biochar alters nitrogen transformations but has minimal effects on nitrous oxide emissions in an organically managed lettuce mesocosm. *Biology Fertility of Soils*, **51**, 573–582.
- Price MN, Dehal PS, Arkin AP (2010) FastTree 2 – Approximately maximum-likelihood trees for large alignments. *PLoS ONE*, **5**, 9490.
- Pruesse E, Quast C, Knittel K *et al.* (2007) SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acid Research*, **35**, 7188–7196.
- Ranjan K, Paula FS, Mueller RC *et al.* (2015) Forest-to-pasture conversion increases the diversity of the phylum Verrucomicrobia in Amazon rainforest soils. *Frontiers Microbiology*, **6**, 779. <https://doi.org/10.3389/fmicb.2015.00779>.
- R Development Core Team (2012) R: a language and environment for statistical computing. 2.15.1. R Foundation for statistical computing, Vienna, Austria.
- Rognes T, Flouri T, Nichols B, Quince C, Mahé F (2016) VSEARCH: versatile open source tool for metagenomics. *PeerJ*, **4**, 2584.
- Ryals R, Hartman MD, Parton WJ, DeLonge MS, Silver WL (2015) Long-term climate change mitigation potential with organic matter management on grasslands. *Ecological Applications*, **25**, 531–545.
- Shade A. (2016). Diversity is the question, not the answer. *ISME Journal*, Advanced online publication, <https://doi.org/10.1038/ismej.2016.118>.
- Serrano-Silva N, Luna-Guido M, Fernández-Luqueño F, Marscha R, Dendooven L (2011) Emission of greenhouse gases from an agricultural soil amended with urea: A laboratory study. *Applied Soil Ecology*, **47**, 92–97.
- Sengupta A, Dick WA (2015) Bacterial community diversity in soil under two tillage practices as determined by pyrosequencing. *Microbial Ecology*, **70**, 853–859.
- Stempfhuber B, Richter-Heitmann T, Regan KM *et al.* (2016) Spatial interaction of archaeal ammonia-oxidizers and nitrite-oxidizing bacteria in an unfertilized grassland soil. *Frontiers in Microbiology*, **6**, 1567.
- Thomson AJ, Giannopoulos G, Pretty J, Baggs EM, Richardson DJ (2012) Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philosophical Transactions of the Royal Society B*, **367**, 1157–1168.
- Van der Wal A, van Veen JA, Pijl AS, Summerbell RC, de Boer W (2006) Constraints on development of fungal biomass and decomposition processes during restoration of arable sandy soils. *Soil Biology and Biochemistry*, **38**, 2890–2902.
- Van der Wal A, Geydan TD, Kuypers TW, de Boer W (2013) A thready affair: linking fungal diversity and community dynamics to terrestrial decomposition processes. *FEMS Microbiology Reviews*, **37**, 477–494.
- Wagg C, Bender SF, Widmer F, van der Heijden MGA (2014) Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings of the National Academy of Science of the United States of America*, **111**, 5266–5270.

Werling BP, Dickson TL, Isaacs R *et al.* (2014) Perennial grasslands enhance biodiversity and multiple ecosystem services in bioenergy landscapes. *Proceedings of the National Academy of Science of the United States of America*, **111**, 1652–1657.

Ye J, Zhang R, Nielsen S, Joseph SD, Huang D, Thomas T (2016) A combination of biochar-mineral complexes and compost improves soil bacterial processes, soil quality, and plant properties. *Frontiers in Microbiology*, **7**, 372.

Supporting Information

Additional Supporting Information may be found online in the supporting information tab for this article:

Figure S1. Distribution of bacterial community composition in the un-amended and residue-amended soils at the phylum (a), family (b), and genus (c) levels.

Figure S2. Venn diagram showing only the significant genera with log-fold changes (comparing the different amendments, as well as the un-amended soil) in the sandy loam soil represented at the family level, which grouped into the phyla Actinobacteria (a), Bacteroidetes (b), Firmicutes (c), and Proteobacteria (d–g). Abbreviations: R1, sewage sludge; R2, aquatic plant material; R3, compost; R4, wood material; R5, compressed beet leaves; R6, paper pulp.

Figure S3. Venn diagram showing only the significant genera with log-fold changes (comparing the different amendments, as well as the un-amended soil) in the clay soil represented at the family level, which grouped into the phyla Actinobacteria (a), Bacteroidetes (b), Firmicutes (c), and Proteobacteria (d–g). Abbreviations: R1, sewage sludge; R2, aquatic plant material; R3, compost; R4, wood material; R5, compressed beet leaves; R6, paper pulp.

Figure S4. Boxplot showing significant genera with log-fold changes in the sandy loam soil after 7 and 13 days incubation in the un-amended soil (S1; a), and soil amended with sewage sludge (R1; b), aquatic plant material (R2; c), compost (R3; d), wood material (R4; e), compressed beet leaves (R5; f), and paper pulp (R6; g).

Figure S5. Boxplot showing significant genera with log-fold changes in the clay soil after 7 and 15 days incubation in the un-amended soil (S2 Control; a), and soil amended with sewage sludge (R1; b), aquatic plant material (R2; c), compost (R3; d), wood material (R4; e), compressed beet leaves (R5; f), and paper pulp (R6; g).