



K O N I N K L I J K E N E D E R L A N D S E  
A K A D E M I E V A N W E T E N S C H A P P E N

## Spatial configuration of plant-soil feedbacks promotes plant species coexistence

Xue, W.; Berendse, Frank; Bezemer, T.M.

2018

### **document version**

Publisher's PDF, also known as Version of record

### **document license**

CC BY

[Link to publication in KNAW Research Portal](#)

### **citation for published version (APA)**

Xue, W., Berendse, F., & Bezemer, T. M. (2018). *Spatial configuration of plant-soil feedbacks promotes plant species coexistence: lay summary*. (Functional Ecology; No. 20180503).  
<https://fesummaries.wordpress.com/2018/05/03/spatial-configuration-of-plant-soil-feedbacks-promotes-plant-species-coexistence/>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the KNAW public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the KNAW public portal.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### **E-mail address:**

[pure@knaw.nl](mailto:pure@knaw.nl)

## Spatial configuration of plant-soil feedbacks promotes plant species coexistence

*Wei Xue, Frank Berendse and T. Martijn Bezemer*

Plant-soil feedbacks refer to how plants change the soil's abiotic and biotic properties where they grow, which can in turn influence the growth of the same or other plant species that grow later in this soil. Each plant influences the soil differently, hence plant-soil feedbacks are spatially varied in the field. Ecologists are beginning to realize the importance of spatial variation of plant-soil feedbacks for plant species coexistence, but we lack empirical evidence.

Here we conducted a two-phase experiment. First, in an experimental garden, we planted sweet vernal grass (*Anthoxanthum odoratum*) and the broad-leaved herb (forb), brown knapweed (*Centaurea jacea*), in monocultures in separate experimental plots. After three years, we collected soils from these plots, hence we had two types of soil: one that had been influenced by the grass (grass soil) and the other one by the forb (forb soil). Then, in a greenhouse we created heterogeneous soils consisting of patches of the two soils in a chessboard manner, and homogeneous soils where the same amount of the two soils were evenly mixed. We planted the grass and the forb together in the heterogeneous and homogeneous soils to examine how the spatial configuration of plant-soil feedbacks may influence plant species coexistence.

We found that the growth difference between the two species was reduced in



*Greenhouse experiment.*

heterogeneous soil compared to homogeneous soil, indicating that heterogeneous soil can promote species coexistence by reducing growth inequalities.

But how? When we looked at the growth of the two plant species in the heterogeneous soil, we found that the grass grew better in forb soil patches than in grass soil patches while the reverse was true for the forb, which may contribute to the more equal growth of the two species in the heterogeneous soil. Future study should examine whether this phenomenon is generally true in the field.